

1  
2  
3  
4

**TITLE PAGE**

**- Korean Journal for Food Science of Animal Resources -**

**Upload this completed form to website with submission**

<b>ARTICLE INFORMATION</b>	<b>Fill in information in each box below</b>
<b>Article Title</b>	Effect of incorporation of pomegranate peel and bagasse powder and their extracts on quality characteristics of chicken meat patties
<b>Running Title (within 10 words)</b>	Effect of pomegranate byproducts on quality of chicken patties
<b>Author</b>	Priyanka Sharma <sup>1</sup> and S. Yadav <sup>1</sup>
<b>Affiliation</b>	1 LUVAS, Hisar, Haryana, India-125004
<b>Special remarks – if authors have additional information to inform the editorial office</b>	-
<b>ORCID (All authors must have ORCID)</b> <b><a href="https://orcid.org">https://orcid.org</a></b>	Priyanka Sharma ( <a href="https://orcid.org/0000-0002-8798-0413">https://orcid.org/0000-0002-8798-0413</a> ) Sanjay Yadav ( <a href="https://orcid.org/0000-0002-3569-6963">https://orcid.org/0000-0002-3569-6963</a> )
<b>Conflicts of interest</b>  List any present or potential conflicts of interest for all authors.  (This field may be published.)	The authors declare no potential conflict of interest.
<b>Acknowledgements</b>  State funding sources (grants, funding sources, equipment, and supplies). Include name and number of grant if available.  (This field may be published.)	-
<b>Author contributions</b>  (This field may be published.)	Conceptualization: Yadav S  Data curation: Sharma Priyanka  Formal analysis: Sharma Priyanka, Yadav S

	<p>Methodology: Sharma Priyanka</p> <p>Software: Sharma Priyanka, Yadav S</p> <p>Validation: Sharma Priyanka</p> <p>Investigation: Sharma Priyanka</p> <p>Writing - original draft: Sharma Priyanka, Yadav S</p> <p>Writing - review &amp; editing: Yadav S</p>
<p><b>Ethics approval (IRB/IACUC)</b></p> <p>(This field may be published.)</p>	<p>This manuscript does not require IRB/IACUC approval because there are no human and animal participants.</p>

5

**6 CORRESPONDING AUTHOR CONTACT INFORMATION**

For the <u>corresponding</u> author (responsible for correspondence, proofreading, and reprints)	Fill in information in each box below
First name, middle initial, last name	SANJAY YADAV
Email address – this is where your proofs will be sent	SYADAV_LPT123@YAHOO.CO.IN
Secondary Email address	
Postal address	DEPTT. OF LPT, LUVAS, HISAR
Cell phone number	91-9254051339
Office phone number	
Fax number	

7

8

9 **Abstract**

10 A study was conducted to develop chicken meat patties by incorporating pomegranate peel and  
11 bagasse powders and their extracts. Patties were developed by incorporating pomegranate peel  
12 powder (PPP, 2 g), pomegranate aril bagasse powder (PABP, 4 g), pomegranate peel powder  
13 aqueous extract (PPAE, 6 g) and pomegranate aril bagasse powder aqueous extract (PABAE, 9 g)  
14 individually per 100 g of minced meat.

15 Both types of powders and extracts treated patties had significantly higher total phenolic content  
16 than control and butylated hydroxytoluene (BHT) treated patties. Both types of powder (PPP and  
17 PABP) treated patties had significantly higher water holding capacity, ash, crude fibre content,  
18 and hardness values, and significantly lower moisture content and lightness values in comparison  
19 to control patties. Emulsion stability and cooking yield of PABP treated patties were  
20 significantly higher than control. Addition of extracts and BHT did not influence the physico-  
21 chemical properties and proximate composition of chicken patties. Both types of powders and  
22 extracts provided better protection to chicken meat patties against oxidative rancidity and  
23 microbial proliferation in comparison to control and BHT treated patties during refrigerated  
24 storage. It is concluded that pomegranate fruit byproducts in the form of peel powder, aril  
25 bagasse powder and their extracts can be successfully utilised in development of healthier  
26 chicken meat patties and these byproducts can also be effectively used as a replacement of  
27 synthetic antioxidants such as BHT.

28 **Key words:** Pomegranate peel powder, pomegranate aril bagasse powder, antioxidants, aqueous  
29 extract, chicken patties

30

31

## 32 **Introduction**

33 The function of diet in the avoidance and treatment of various diseases has been extensively  
34 acknowledged. The idea of functional foods is attaining recognition nowadays as consumers are  
35 trying to improve their health by natural means. It has been observed that such foods promote  
36 one or more functions in the body apart from regular dietary effects. Studies related to  
37 development of healthier and functional meat and meat products would prove beneficial to the  
38 meat industry and eventually the consumers as the relation involving diet and chronic disease  
39 avoidance continues to rise (Kandeepan et al., 2007).

40 Meat and its products supply good quality proteins, essential fatty acids, minerals, vitamins, and  
41 additional nutrients, but they are deficient in dietary fibre. Consistent eating of meat products is  
42 being linked with a range of health problems like colon cancer, obesity, and cardiovascular  
43 ailments (Larsson and Wolk, 2006; Tarrant, 1998).

44 Lipid oxidation, autoxidation, and microbial proliferation are the main reasons for quality decline  
45 in meat and meat products. Changes in poultry meat due to oxidative rancidity may differ  
46 considerably from slight decrease in freshness to considerable flavour changes, colour losses,  
47 and structural degradation of proteins. Rancidity process is started due to vulnerability to the  
48 enzyme lipoxygenase, metalloprotein catalysts, heat, ionizing radiation, light, and metal ions  
49 (Daker et al., 2008). Meat processing procedures like mincing and cooking distort membranes of  
50 muscle cells promoting the interaction of unsaturated lipids with prooxidant substances like non-  
51 haem iron. This escalates lipid oxidation resulting in rancidity which deteriorates the quality of  
52 meat products (Tichivangana and Morrissey, 1985).

53 Synthetic antioxidants have been used in meat and poultry products to inhibit lipid oxidation. But  
54 they have come under watch because of their possible toxicological effects (Naveena et al.,

55 2008; Nunez de Gonzalez et al., 2008). Therefore, significance of natural source of antioxidants  
56 for application in meat products has increased in recent years.

57 Pomegranate peel and pomegranate seed/aril bagasse are main byproducts of pomegranate juice  
58 industry (Jalal et al., 2018). Pomegranate peel is major source of bioactive compounds like  
59 phenolics, flavonoids, ellagitannins, and proanthocyanidin compounds. Antioxidant and  
60 antibacterial quality of peel of pomegranate in in-vitro experimental models have been  
61 documented. Pomegranate peel extracts curb the growth of many food borne pathogens (Agourram  
62 et al., 2013; Al-Zoreky, 2009). Pomegranate seed/bagasse have a variety of nutraceutical  
63 compounds like sterols,  $\gamma$ -tocopherol, punicic acid, and hydroxyl benzoic acids (Liu et al., 2009).  
64 Pomegranate seed extracts exhibited antidiarrheal and antioxidant bioactivities (Singh et al.,  
65 2002). Bhol and Bosco (2013) found that pomegranate aril bagasse and pomegranate whole fruit  
66 bagasse are rich source of dietary fibre. Due to presence of valuable pharmaceutical and  
67 nutritional compounds, these byproducts can be better utilized in food industry in place of being  
68 exploited as feed for animals or marketable cosmetic products (Liu et al., 2009).

69 Poultry industry is attaining more significance worldwide in present times due to its better  
70 consumer recognition and freedom from religious hindrances. Huge growth of fast food market  
71 has escalated ready to eat snack foods demand. Chicken patty is one of the favourite comminuted  
72 products which have a noticeable position because of its distinctive flavour and palatability (Raut  
73 et al., 2011). In light of above information, this study was undertaken to develop chicken meat  
74 patties by incorporating powder and aqueous extracts of pomegranate peel and aril bagasse, and  
75 study the effect of their addition on quality characteristics and shelf life of developed products.

## 76 **Materials and methods**

### 77 **Preparation of pomegranate peel and bagasse powder and their extracts**

78 Pomegranate peels were collected from local market juice shop. Pomegranate aril bagasse was

79 obtained after extraction of juice from pomegranate fruit. Both the byproducts were separately  
80 washed with clean water. They were squeezed through muslin cloth to remove extra water and  
81 dried in hot air drier at a temperature of about 50-55°C for about 2 d. After complete drying,  
82 both the dried products were ground to fine powder in a grinder, packed in polythene bags  
83 separately and stored at  $-18\pm 2^{\circ}\text{C}$  for further use.

84 For preparing extracts, 10 g of each type of dried powder was mixed in 100 mL of distilled water  
85 separately. Both the mixtures were incubated for overnight at room temperature. Each type of  
86 mixture was filtered through muslin cloth and the filtrate was retained for use in chicken meat  
87 patties. Fresh extracts were prepared each time for use during product development and test  
88 procedures.

89 **Slaughtering and dressing of chicken** Birds were slaughtered and dressed following the  
90 established procedure in the experimental slaughter house of the Department. The dressed chickens  
91 were deboned manually, washed and packaged in low density polyethylene bags and stored at  
92  $-18\pm 2^{\circ}\text{C}$  till further use. The frozen meat chunks were drawn according to necessity and thawed  
93 overnight in a refrigerator ( $4\pm 2^{\circ}\text{C}$ ) for further use.

94 **Preparation of control and treated meat patties** Deboned frozen meat was cut into small  
95 pieces and minced in an electrical mincer (3 mm plate) (Mado Primus Meat Mincer MEW-613,  
96 Dr. Froeb India Pvt. Ltd., Noida, India). In control meat patties, 100 g of minced meat was taken  
97 to which sodium chloride, sodium tripolyphosphate, sodium nitrite, spice mix, condiments  
98 (ginger and garlic in ratio of 1:1), bread crumbs, water, egg liquid, and fat were added in suitable  
99 proportion (Table 1) and blended with the minced meat in a mixer (Select 600; Morphyrichards  
100 food processor, Mumbai, India) for 4 to 5 min. Treatments consisted of addition of BHT,  
101 pomegranate peel and bagasse powder and their aqueous extracts separately to minced meat.

102 Approximately 60 g of meat emulsion was hand moulded into patty shape using a petri dish.  
103 Patties were prepared by baking in a preheated oven at a temperature of 160°C for 35 min (20  
104 min first side and 15 min second side). Both control and treated patties were packaged in low  
105 density polythene bags and subjected to physico-chemical, nutritional, instrumental colour, and  
106 texture analysis. Patties were also stored under refrigerated storage ( $4\pm 2^{\circ}\text{C}$ ) conditions and  
107 physico-chemical and microbiological quality of the products were analysed at a regular interval  
108 of 4 d upto 16 d of storage.

109 **Total Phenolic content** Folin Ciocalteu's technique was followed to determine total phenolic  
110 content. Absorbance was measured at 750 nm by UV visible spectrophotometer (G 10 S UV-VIS;  
111 Thermo Fisher Scientific India Pvt. Ltd., Mumbai, India). For standard, gallic acid was used and  
112 results were calculated as mg of gallic acid equivalent (GAE)/100 g of dry mass (Bhalodia et al.,  
113 2011).

114 **Physico-chemical parameters** The pH of chicken patties was estimated with pH meter (Cyber  
115 Scan pH 510; Eutech Instruments, Thermo Fisher Scientific India Pvt. Ltd., Mumbai, India)  
116 following the procedure of Trout et al. (1992). Control and treated emulsions stability were  
117 estimated using the method of Baliga and Madaiah (1970). For cooking yield, the weight of raw  
118 and cooked patties was measured and yield was expressed as percentage. Water holding  
119 capacity (WHC) was determined as per the procedure of Wardlaw et al. (1973). TBA value of  
120 patties was estimated as per the method of Witte et al. (1970). Trichloroacetic acid extract of  
121 meat samples was mixed with thiobarbituric acid reagent. The contents were placed in boiling  
122 water bath and optical density was determined at 540 nm. TBA value was calculated as mg  
123 malonaldehyde/kg of sample.

124 **Proximate composition** Moisture, protein, fat, ash, and crude fibre content of chicken meat

125 patties were estimated by standard procedure of AOAC (2005). Finally chopped sample of meat  
126 was dried in hot air oven (JSGW, Ambala, India), cooled in desiccator and loss in weight was  
127 expressed as moisture content of the sample. Protein was estimated using Kjeldahl digestion  
128 method. The fat content of samples was determined by solvent extraction method using  
129 petroleum ether (60 to 80°C) as solvent. Ashing was done in a muffle furnace (Yorco, Yorco  
130 sales Pvt. Ltd., New Delhi, India) set at 550°C to determine ash content of samples. For crude  
131 fibre content, fat free samples were subjected to acid and alkali digestion. The residue remaining  
132 after digestion was weighed and subjected to ashing. Difference in weight was calculated as  
133 crude fibre.

134 **Texture profile analysis** The textural properties of patties were evaluated by the method of  
135 Bourne (1978) using TA HD plus twin column texture analyser (Stable Micro Systems Ltd.,  
136 Vienna Court, Lammas Road, Godalming, Surrey GU7 1YL, United Kingdom) equipped with  
137 the exponent software (version 5,1,1,0 Lite).

138 Samples of 2 cm<sup>3</sup> size were compressed (by 70 mm compression plate, 50 kg load cell and the  
139 test speed of 2 mm/s) to 50% of their initial height. Between two compression cycles, 5 s time  
140 interval was given to obtain force time deformation curves. Hardness (N), cohesiveness,  
141 springiness, gumminess (N), and chewiness (N) of the samples were analyzed.

142 **Firmness and toughness** The force required to shear a 1cm<sup>3</sup> thick sample of cooked chicken  
143 meat patties transversely was analysed using warner-bratzler shear probe of texture analyser  
144 (TA.HD plus; Stable Micro Systems Ltd., Vienna Court, Lammas Road, Godalming, Surrey  
145 GU7 1YL, United Kingdom). A force time curve was obtained. Firmness (N) was the maximum  
146 shear force required to cut the sample. Toughness (N-s) was the total shear energy (i.e. work of  
147 shear) calculated as the area under force time curve from start to the end of the shear test.

148 **Instrumental colour analysis** The colour scores of chicken patties were measured as CIE Lab,  
149 L\* (lightness), a\* (redness), and b\* (yellowness) using a chroma meter (Konica Minolta Sensing,  
150 Inc., Japan) with 8 mm orifice for measurement. The equipment was standardized with a white  
151 standard plate before measurement.

152 **Microbiological evaluation** Total plate count (TPC), psychrotrophic, and thermophilic counts of  
153 chicken patties were estimated during refrigerated storage (APHA, 2001).

154 **Statistical analysis** The data acquired were evaluated by analysis of variance. For fresh products,  
155 one-way analysis of variance and for refrigerated stored products, two-way analysis of variance  
156 were performed. Duncan's test was performed using SPSS version 16 (SPSS Inc., Chicago, USA)  
157 to determine the significant difference in the mean values at 5% significance level.

## 158 **Results and discussion**

159 **Total phenolic content** A small quantity of phenolic compounds were observed in control raw  
160 emulsion and patties which was due to polyphenols provided by spices and condiments added to  
161 chicken meat emulsion (Table 2). Polyphenolic content and antioxidant capacity of different spices  
162 have been well established (Pellegrini et al., 2006; Zheng and Wang, 2001). Addition of BHT,  
163 pomegranate peel and bagasse and their extracts to chicken meat resulted in an appreciable  
164 increase in total phenolic content. Also, pomegranate by products and their extracts treated patties  
165 had significantly higher phenolic content than BHT treated patties. PPP treatment had significantly  
166 higher phenolic content in comparison to PABP treatments. Similarly, peel extract treated  
167 emulsion and patties had significantly higher phenolic content than bagasse extract treated  
168 emulsion and patties, respectively. Higher phenolic content in peel treated emulsion and patties in  
169 comparison to bagasse treated emulsion and patties was due to higher total phenolic content in PPP  
170 (52.3 mg GAE/g) in comparison to PABP (30.1 mg GAE/g). A number of research workers have

171 demonstrated higher phenolic content in pomegranate peel in comparison to its pulp and bagasse.  
172 Findings of present research work are in agreement with previous studies. Li et al. (2006) revealed  
173 that peel tissues of pomegranate normally contain higher quantity of phenolics than its pulp.  
174 Sultana et al. (2008) reported that total phenol and total flavonoid content of pomegranate peel  
175 powder were higher in comparison to pomegranate aril bagasse and whole fruit  
176 bagasse powder. Devatkal and Naveena (2010) also reported higher total phenolic content in  
177 pomegranate rind powder (PRP) in comparison to pomegranate seed powder (PSP). Within a  
178 particular byproduct, powder treatment had higher phenolic content in comparison to respective  
179 extract treatment. Higher amount of phenolics in powder treated emulsion and patties in  
180 comparison to respective extract treated emulsion and patties might be due to less extraction of  
181 phenolics in aqueous extract. Separation of phenolic compounds from their innate structure is  
182 complex due to their heterogenicity and vulnerability to oxidation and hydrolysis (Naczka and  
183 Shahidi, 2004). Kind of solvent, proportion of solid to liquid, extraction temperature, and peel  
184 particles size significantly affect antioxidant extraction (Ismail et al., 2012).

185 **Physico-chemical qualities of chicken meat emulsion and patties** The pH of raw emulsion and  
186 cooked chicken patties did not vary significantly between control and treatments, although a non-  
187 significantly lower pH was noticed in pomegranate byproducts and their extract incorporated patties  
188 (Table 2). WHC increased significantly in pomegranate by products i.e. PPP and PABP  
189 incorporated patties in comparison to control. PABP treated patties had significantly highest  
190 water holding capacity value among all treatments. Higher WHC in PPP and PABP treated  
191 patties might be due to presence of dietary fibre in pomegranate peel and bagasse powder which  
192 has been reported to increase the WHC (Cofrades et al., 2000). Viuda-Martos et al. (2012)  
193 reported that pomegranate bagasse powder co-product exhibited WHC equal to 4.86 times of its

194 own weight. Akhtar et al. (2015) found that incorporation of 3% PRP in raw beef sausage  
195 improved their WHC.

196 Emulsion stability and cooking yield of PABP treated patties was significantly higher than  
197 control. As discussed earlier, higher cooking yield of PABP treated patties was due to presence  
198 of dietary fibres which increases cooking yield because of their water and fat binding attributes  
199 (Cofrades et al., 2000). PPP treated patties also had dietary fibre, but it did not result in  
200 significant increase in cooking yield, although a non-significantly higher yield than control was  
201 observed in PPP treated patties. This might be due to lower level of PPP used in patties. Abdel  
202 Fattah et al. (2016) also reported better cooking yield in beef burgers incorporated with PPP.  
203 Addition of BHT and pomegranate by product aqueous extracts (PPAE and PABAE) did not  
204 result in any significant effect on WHC, emulsion stability, and cooking yield of chicken meat  
205 patties.

206 **Proximate composition** Addition of pomegranate peel and aril bagasse powder contributed to  
207 significant decline in moisture content of patties (Table 3). This was due to replacement of meat  
208 with dried peel and aril bagasse containing very low moisture. The moisture content of dried peel  
209 powder and dried aril bagasse powder was 11.06% and 5.55%, respectively. El-Nashi et al.  
210 (2015) revealed that addition of PPP to beef sausage contributed to significant decline in  
211 moisture content. Powder treated patties had significantly lower moisture content in comparison  
212 to respective extract treated patties.

213 Protein content of control and treated patties did not varied remarkably. Addition of pomegranate  
214 aril bagasse powder contributed to significant increase in fat content and treatment PABP had  
215 significantly higher fat content than control. Higher fat content in PABP treated patties was due  
216 to more retention of fat and presence of higher amount of fat in pomegranate aril bagasse powder

217 (16.86%). Earlier, Ozgul-Yucel (2005) revealed that pomegranate seeds are a plentiful source of  
218 total lipids. Ash content of pomegranate peel and bagasse powder treated patties was  
219 significantly higher in comparison to control which was due to higher ash content in PPP (4.03%)  
220 and PABP (3.22%). A small quantity of crude fibre was observed in control chicken patties  
221 which was attributable to dietary fibre provided by spices and condiments incorporated in raw  
222 emulsion. A significant increase in crude fibre content in PPP and PABP patties was observed  
223 which was because of higher crude fibre content in pomegranate peel and aril bagasse powder.  
224 PABP patties also had significantly higher crude fibre content in comparison to PPP patties. This  
225 was due to more amount of PABP (4 g) incorporated in chicken meat in comparison to PPP (2 g).  
226 Also, higher amount of crude fibre in PABP (32.32%) in comparison to PPP (15.80%)  
227 contributed to significantly higher crude fibre content in PABP treated patties. Rowayshed et al.  
228 (2013) revealed that the PPP and PSP are considered as good source of crude fibre. Bhol and  
229 Bosco (2013) observed a substantial increase in fibre content in bread with rise in level of  
230 pomegranate bagasse powder. Addition of BHT, PPAE, and PABAE did not influence the  
231 proximate composition of chicken patties.

232 **Instrumental colour analysis** Incorporation of pomegranate by products and their extracts  
233 contributed to a decline in lightness and significant decline in relation to control was noticed in  
234 PPP and PABP treated patties (Table 4). Significant decrease in lightness in these treatments  
235 might be due to dark colour contributed by pomegranate peel and bagasse. Monsalve Gonzalez et  
236 al. (1994) reported that colour is influenced by various factors, but drying process in particular  
237 influence the colour. Pulp is exposed to high temperature during its drying, which leads to  
238 enzymatic and non-enzymatic browning resulting in darkening of the product. Addition of BHT,  
239 powders, and extracts did not influence the redness and yellowness values of chicken patties.

240 **Texture profile, firmness and toughness of chicken meat patties** Texture profile analysis of  
241 control and treated patties revealed a significant increase in hardness of PPP and PABP treated  
242 patties (Table 5). Higher hardness values in these treatments might be due to hardness and  
243 rigidity contributed by pomegranate peel and bagasse. Lower moisture content in these  
244 treatments might have contributed to higher hardness. Yadav et al. (2016) also documented an  
245 increase in hardness scores of dried apple pomace added chicken sausage. PPP patties had  
246 significantly higher hardness than PABP treated patties. Gumminess score of PPP treated patties  
247 were significantly higher in comparison to control, which was due to their higher hardness values.  
248 Chewiness values of both the extract treated patties were significantly lower in comparison to  
249 PPP treated patties, which was due to their significantly lower hardness value and non-  
250 significantly lower springiness values. Pomegranate byproducts powder addition resulted in an  
251 increase in firmness and toughness, and significant rise in relation to control was observed in  
252 PPP incorporated patties. Addition of BHT, PPAAE, and PABAAE did not affect texture profile,  
253 firmness, and toughness of chicken patties.

254 **Physico-chemical and microbiological quality of pomegranate fruit byproducts treated**  
255 **chicken patties during refrigerated storage**

256 **pH values** pH of control and treated patties did not vary considerably during refrigerated storage  
257 (Table 6). There was a modest decrease in pH of chicken meat patties during initial days of  
258 refrigeration. Subsequently, pH increased on 12 d and 16 d of storage. Lactic acid bacteria  
259 multiply in the beginning, which leads to disintegration of sugar into acids. Subsequently,  
260 bacterial deamination of proteins occurs which raises the pH of the product (Jay, 1996).

261 **Thiobarbituric acid (TBA) value** TBA value of control and treated patties increased  
262 significantly during refrigerated storage. However, TBA values of PPP and PABP treated

263 chicken patties were significantly lower than control throughout the storage period. TBA values  
264 of BHT, PPAE, and PABAE were also lower than control during storage. Significant difference  
265 was observed from 8 d onward in PPAE and PABAE treated patties and 12 d onward in BHT  
266 treated patties. Bioactive compounds like phenols and flavonoids present in pomegranate peel  
267 and bagasse provided antioxidant effect and inhibited lipid oxidation in chicken patties resulting  
268 in lower rise in TBA values in these treatments. The results indicate that pomegranate peel,  
269 bagasse, and extracts provided better protection against rise in TBA value in comparison to BHT,  
270 and can be used as natural antioxidant sources in chicken meat patties in place of synthetic  
271 antioxidants. Decrease in TBA value of meat and meat products due to antioxidant effect of  
272 pomegranate fruit byproducts, and their extracts has been reported previously by various  
273 research workers. El-Gharably and Ashoush (2011) and Naveena et al. (2008) observed that PPP  
274 improved the storage stability of meat products during refrigeration by reducing the rate of lipid  
275 oxidation. Devatkal et al. (2010) also observed a significant reduction in TBARS values of goat  
276 meat patties treated with the extracts of PRP and PSP as compared to control during refrigerated  
277 storage. Abdel Fattah et al. (2016) reported lower TBARS value in PPP incorporated beef  
278 burgers in comparison to control on 12 d of storage.

279 **Microbiological status** TPC, psychrotrophic count, and thermophilic count of control and treated  
280 patties increased significantly with increase in storage duration in all the treatments (Table 7). The  
281 rate of increase was less in pomegranate byproducts and their extract treated patties. Inhibitory  
282 effect of bioactive and phenolic compounds present in pomegranate peel, bagasse, and their  
283 extracts resulted in significantly lower TPC in treated patties in comparison to control at the end of  
284 storage. Maximum inhibitory effect on growth of microbes was noticed in PPP treated patties  
285 resulting in significantly lowest TPC at the end of storage. PPP treated patties also had

286 significantly lower psychrotrophic and thermophilic counts in comparison to control patties at the  
287 end of storage. Results indicate that pomegranate fruit by products such as peel, aril bagasse, and  
288 their extracts possess antimicrobial activity. Alzoreky (2009) observed that pomegranate extract  
289 intervened with production of bacterial proteins. Chandralekha et al. (2012) observed a significant  
290 decrease in standard plate count of 5% rind powder incorporated chicken meatball in comparison  
291 to control during refrigerated storage.

## 292 **Conclusion**

293 Incorporation of pomegranate fruit byproducts in the form of peel powder, aril bagasse powder,  
294 and their aqueous extracts improved nutritional value of chicken patties by increasing their total  
295 phenolics content. TBA values of powder and extract treated patties were lower in comparison to  
296 control and BHT treated patties during refrigerated storage. This study indicates that  
297 pomegranate fruit byproducts and their extracts can be used as a replacement of synthetic  
298 antioxidants such as butylated hydroxytoluene for development of chicken meat patties. Both the  
299 powders provided additional nutritional benefit by increasing crude fibre content of patties while  
300 bagasse powder also increased their cooking yield. Microbial counts in powder and extract  
301 treated patties were lower than control during refrigerated storage. Best antimicrobial effect was  
302 observed in pomegranate peel powder treated patties.

303

304 **References**

- 305 Abdel Fattah AA, Abdel-Rahman NR, El-Razik MA, Hafssa B. 2016. Utilization of  
306 Pomegranate Peels for Improving Quality Attributes of Refrigerated Beef Burger. *Curr*  
307 *Sci Int* 05: 427- 441.
- 308 Agourram A, Ghirardello D, Rantsiou K, Zeppa G, Belviso S, Romane A. 2013. Phenolic  
309 content, antioxidant potential and antimicrobial activities of fruit and vegetable by-  
310 product extracts. *Int J Food Prop* 16: 1092-1104.
- 311 Akhtar S, Ismail T, Fraternali D, Sestili P. 2015. Pomegranate peel and peel extracts: Chemistry  
312 and food features. *Food Chem* 174: 417-425.
- 313 Al-Zoreky NS. 2009. Antimicrobial activity of pomegranate (*Punica granatum* L.) fruit peels.  
314 *Intl J Food Microbiol* 134: 244-248.
- 315 AOAC 2005. Official Methods of Analysis, 16<sup>th</sup> edition, Association of Official Analytical  
316 Chemists, Washington, DC.
- 317 APHA 2001. Recommended methods for microbiological examination of foods. Washington,  
318 DC.
- 319 Baliga BR, Madaiah N. 1970. Quality of sausage emulsion prepared from mutton. *J Food Sci* 35:  
320 383-385.
- 321 Bhalodia N, Nariya P, Acharya R, Shukla V. 2011. Evaluation of in vitro antioxidant activity of  
322 flowers of *Cassia fistula* Linn. *Int J Pharmtech Res* 3: 589-599.
- 323 Bhol S, Bosco JD. 2013. Enrichment of yeast leavened bread by pomegranate bagasse  
324 powder. *Int J Sci Res* 2(5): 298-300.
- 325 Bourne MC. 1978. Texture profile analysis. *Food Technol* 33: 62-66, 72.
- 326 Chandralekha S, Angalakuditi JB, Sreenivasa PR, Balakrishnan K. 2012. Studies on the effect of  
327 pomegranate rind powder extract as natural antioxidant in chicken meat balls during  
328 refrigerated storage. *J Adv Vet Res* 2: 107-112.
- 329 Cofrades S, Guerra MA, Carballo J, Fernandez-Martin F, Jimenez-Colmenero F. 2000. Plasma  
330 protein and soy fibre content effect on bologna sausage properties as influenced by fat  
331 level. *J Food Sci* 65:281-287.
- 332 Daker M, Abdullah N, Vikineswary S, Goh PC, Kuppusamy UR. 2008. Antioxidant from maize  
333 and maize fermented by *Marasmiellus* sp. as stabiliser of lipid-rich foods. *Food Chem*  
334 107: 1092-1098.
- 335 Devatkal SK, Narsaiah K, Borah A. 2010. Anti-oxidant effect of extracts of kinnow rind,  
336 pomegranate rind and seed powders in cooked goat meat patties. *Meat Sci* 85: 155-159.

- 337 Devatkal SK, Naveena BM. 2010. Effect of salt, kinnow and pomegranate fruit by-product  
338 powders on color and oxidative stability of raw ground goat meat additive LWT-Food Sci  
339 Technol 44: 307-311.
- 340 El-Gharably AMA, Ashoush IS. 2011. Utilization impact of adding pomegranate rind powder  
341 and red beet powder as natural antioxidant on quality characteristics of beef sausages.  
342 World J Dairy Food Sci 6: 86-97.
- 343 El-Nashi HB, Fattah AFAKA, Rahman NRA, El-Razik MA. 2015. Quality characteristics of  
344 beef sausage containing pomegranate peels during refrigerated storage. Ann Agric Sci 60:  
345 403-412.
- 346 Ismail Tariq, Sestili Piero, Akhtar Saeed. 2012. Pomegranate peel and fruit extracts: A review of  
347 potential anti inflammatory and anti infective effects. J Ethnopharmacol 143: 397-405.
- 348 Jalal H, Pal MA, Ahmad SR, Rather M, Andrabi M, Hamdani S. 2018. Physico-chemical and  
349 functional properties of pomegranate peel and seed powder. J Pharm Innov 7: 1127-1131.
- 350 Jay JM. 1996. Modern food Microbiology. 4th (Edn). CBS Publishers and Distributors, New  
351 Delhi, India.
- 352 Kandeepan G, Anjaneyuly ASR, Gadekar YP. 2007. Meat as a functional food. Processed Food  
353 Ind 5: 30-37.
- 354 Larsson SC, Wolk A. 2006. Meat consumption and risk of colorectal cancer: a meta-analysis of  
355 prospective studies. Inter J Cancer 119: 2657-2664.
- 356 Li Y, Guo C, Yang J, Wei J, Xu J, Cheng S. 2006. Evaluation of antioxidant properties of  
357 pomegranate peel extract in comparison with pomegranate pulp extract. Food Chem 96:  
358 254-260.
- 359 Liu G, Xu X, Hao Q, Gao Y. 2009. Supercritical CO<sub>2</sub> extraction optimization of pomegranate  
360 (*Punica granatum L.*) seed oil using response surface methodology. LWT- Food Sci  
361 Technol 42: 1491-1495
- 362 Monsalve Gonzalez R, Enríquez AV. 1994. Sugar cane forage for milk production. Effect of the  
363 inclusion of star grass (*Cynodon nlemfuensis*) forage on rumen liquor turnover. Cuban J  
364 Agric Sci 28: 285-287.
- 365 Naczki M, Shahidi F. 2004. Extraction and analysis of phenolics in food. J Chromatogra A 1054:  
366 95-111.
- 367 Naveena BM, Sen AR, Vaithyanathan S, Babji Y, Kondaiah N. 2008. Comparative efficacy of  
368 pomegranate juice, pomegranate rind powder and BHT in cooked chicken patties. Meat  
369 Sci 80:1304-1308.

- 370 Nunez de Gonzalez MT, Hafley BS, Boleman RM, Miller RK, Rhee KS, Keeton JT. 2008.  
371 Antioxidant properties of plum concentrates and powder in precooked roast beef to  
372 reduce lipid oxidation. *Meat Sci* 80: 997-1004
- 373 Ozgul-Yucel S. 2005. Determination of conjugated linolenic acid content of selected oil seeds  
374 grown in Turkey. *J Am Oil Chem Soc* 82:893-897.
- 375 Pellegrini N, Serafini M, Salvatore S, Del Rio D, Bianchi M, Brighenti F. 2006. Total  
376 antioxidant capacity of spices, dried fruits, nuts, pulses, cereals and sweets consumed in  
377 Italy assessed by three different in vitro assays. *Mol Nutr Food Res* 50: 1030-1038.
- 378 Raut KA, Raziuddin M, Zanzad PN, Manwar SJ. 2011. Physico-chemical properties and sensory  
379 qualities of chicken patties prepared with cereal and non-cereal extender. *J Anim Sci* 5:  
380 123-127.
- 381 Rowayshed G, Salama A, Abul-Fadl M, Akila-Hamza S, Emad A Mohamed. 2013. Nutritional  
382 and chemical evaluation for pomegranate (*Punica granatum L.*) fruit peel and seeds  
383 powders by products. *Middle East J Appl Sci* 3:169-179.
- 384 Singh RP, Murthy KNC, Jayaprakasha GK. 2002. Studies on the antioxidant activity of  
385 pomegranate (*Punica granatum*) peel and seed extracts using in vitro models. *J Agric*  
386 *Food Chem* 50: 81-86.
- 387 Sultana B, Anwar F, Asi MR, Chatha SAS. 2008. Antioxidant potential of extracts from different  
388 agro wastes: Stabilization of corn oil. *Grasas Aceites* 59: 205-217.
- 389 Tarrant PV. 1998. Some recent advances and future priorities in research for the meat industry.  
390 *Meat Sci* 49: S1-S16.
- 391 Tichivangana JZ, Morrissey PA. 1985. Metmyoglobin and inorganic metals as prooxidantes in  
392 raw and cooked muscle systems. *Meat Sci* 15: 107-116.
- 393 Trout ES, Hunt MC, Johson DE, Clans JR, Castner CL, Kroff DH. 1992. Characteristics of low  
394 fat ground beef containing texture modifying ingredients. *J Food Sci* 57:19-24.
- 395 Viuda-Martos M, Ruiz-Navajas Y, Martin-Sanchez A, Sanchez-Zapata E, Fernandez-Lopez J,  
396 Sendra E et al 2012 Chemical, physico-chemical composition, functional properties of  
397 pomegranate (*Punica granatum L.*) bagasse powder coproduct. *J Food Eng* 10:220- 224.
- 398 Wardlaw FR, McCaskill LH, Acton JC. 1973. Effects of postmortem changes on poultry meat  
399 loaf properties. *J Food Sci* 38: 421-423.
- 400 Witte VC, Krouze GF, Bailey ME. 1970. A new extraction method for determining 2-  
401 thiobarbituric acid values of pork and beef during storage. *J Food Sci* 35: 582-585.

402 Yadav S, Malik A, Pathera A, Islam RU, Diwakar S. 2016. Development of dietary fibre  
403 enriched chicken sausages by incorporating corn bran, dried apple pomace and dried  
404 tomato pomace. *Nutr Food Sci* 46: 16-29.

405 Zheng W, Wang SY. 2001. Antioxidant activity and phenolic compounds in selected herbs. *J*  
406 *Agric Food Chem* 49: 5165- 5170.

407

ACCEPTED

408

409 **Table 1. Composition of meat emulsion for preparation of chicken meat patties**

Treatments	Control	BHT	PPP	PABP	PPAE	PABAE
<b>Ingredients</b>						
Meat	100	100	100	100	100	100
Sodium tripolyphosphate	0.4	0.4	0.4	0.4	0.4	0.4
Sodium chloride	1.9	1.9	1.9	1.9	1.9	1.9
Spice mix	2.0	2.0	2.0	2.0	2.0	2.0
Condiments (Ginger: Garlic) 1:1	3	3	3	3	3	3
Fat	15	15	15	15	15	15
Egg liquid	5	5	5	5	5	5
Water	10	10	10	10	10	10
Sodium nitrite	0.015	0.015	0.015	0.015	0.015	0.015
Bread crumbs	2	2	2	2	2	2
Butylated hydroxytoluene	-	0.01	-	-	-	-
Dried pomegranate peel powder	-	-	2	-	-	-
Dried pomegranate aril bagasse powder	-	-	-	4	-	-
Dried pomegranate peel powder aqueous extract	-	-	-	-	6	-
Dried pomegranate aril bagasse powder aqueous extract	-	-	-	-	-	9

410

411 PPP, PABP, PPAE and PABAE- chicken meat patties incorporated with 2 g dried pomegranate  
412 peel powder, 4 g dried pomegranate aril bagasse powder, 6 g dried pomegranate peel powder  
413 aqueous extract and 9 g dried pomegranate aril bagasse powder aqueous extract, respectively.

414 **Table 2. Effect of pomegranate peel powder, pomegranate aril bagasse powder, and their aqueous extracts on total phenolic**  
 415 **content and physico-chemical properties of chicken meat emulsion and patties**

Treatments	Total phenols (Raw) (mg GAE/g)	Total phenols (Cooked) (mg GAE/g)	pH (Raw emulsion)	pH (Cooked)	Water holding capacity (%)	Emulsion stability (%)	Cooking yield (%)
Control	0.11±0.04 <sup>e</sup>	0.12±0.07 <sup>e</sup>	6.13±0.06 <sup>a</sup>	6.30±0.10 <sup>a</sup>	43.60±1.42 <sup>c</sup>	92.56±0.77 <sup>b</sup>	83.65±0.69 <sup>b</sup>
BHT	0.27±0.06 <sup>d</sup>	0.28±0.04 <sup>d</sup>	6.10±0.10 <sup>a</sup>	6.25±0.08 <sup>a</sup>	44.13±1.94 <sup>bc</sup>	93.40±1.47 <sup>b</sup>	83.36±2.37 <sup>b</sup>
PPP	1.36±0.19 <sup>a</sup>	1.49±0.08 <sup>a</sup>	6.07±0.13 <sup>a</sup>	6.21±0.14 <sup>a</sup>	46.24±1.16 <sup>b</sup>	93.78±1.28 <sup>b</sup>	85.37±2.70 <sup>ab</sup>
PABP	0.84±0.06 <sup>b</sup>	0.88±0.02 <sup>b</sup>	6.09±0.06 <sup>a</sup>	6.23±0.09 <sup>a</sup>	48.78±2.12 <sup>a</sup>	95.00±0.47 <sup>a</sup>	86.03±1.50 <sup>a</sup>
PPAE	0.88±0.07 <sup>b</sup>	0.91±0.06 <sup>b</sup>	6.07±0.10 <sup>a</sup>	6.23±0.12 <sup>a</sup>	44.15±2.20 <sup>bc</sup>	92.98±0.77 <sup>b</sup>	84.15±1.58 <sup>ab</sup>
PABAE	0.66±0.12 <sup>c</sup>	0.69±0.15 <sup>c</sup>	6.08±0.08 <sup>a</sup>	6.24±0.10 <sup>a</sup>	44.36±1.49 <sup>bc</sup>	93.62±0.87 <sup>b</sup>	84.20±0.77 <sup>ab</sup>

416 (n=6, Mean ± SD)

417 Means with different superscripts in a column differ significantly (p≤0.05).

418 BHT, 100 ppm BHT; PPP, 2 g pomegranate peel powder; PABP, 4 g pomegranate aril bagasse powder; PPAE, 6 g pomegranate peel  
 419 powder aqueous extract; PABAE, 9 g pomegranate aril bagasse powder aqueous extract.

420

421

422

423

424 **Table 3. Effect of pomegranate peel powder, pomegranate aril bagasse powder, and their**  
 425 **aqueous extracts on proximate composition of chicken meat patties**

Treatments	Moisture (%)	Protein (%)	Fat (%)	Ash (%)	Crude fibre (%)
Control	60.08±0.97 <sup>a</sup>	17.08±0.80 <sup>a</sup>	16.58±0.39 <sup>b</sup>	1.73±0.24 <sup>b</sup>	0.26±0.13 <sup>c</sup>
BHT	60.06±0.99 <sup>a</sup>	16.76±0.99 <sup>a</sup>	16.68±0.56 <sup>b</sup>	1.70±0.12 <sup>b</sup>	0.29±0.11 <sup>c</sup>
PPP	58.61±1.21 <sup>bc</sup>	16.61±0.95 <sup>a</sup>	17.04±0.75 <sup>ab</sup>	2.10±0.15 <sup>a</sup>	0.60±0.10 <sup>b</sup>
PABP	57.64±1.20 <sup>c</sup>	17.15±1.36 <sup>a</sup>	17.60±0.57 <sup>a</sup>	2.15±0.14 <sup>a</sup>	1.58±0.10 <sup>a</sup>
PPAE	59.93±0.99 <sup>a</sup>	16.89±1.18 <sup>a</sup>	16.83±0.88 <sup>ab</sup>	1.84±0.18 <sup>ab</sup>	0.34±0.08 <sup>c</sup>
PABAE	59.74±0.85 <sup>ab</sup>	16.33±0.63 <sup>a</sup>	16.96±0.63 <sup>ab</sup>	1.85±0.26 <sup>ab</sup>	0.35±0.08 <sup>c</sup>

426 (n=6, Mean ± SD)

427 Means with different superscripts in a column differ significantly (p≤0.05).

428 BHT, 100 ppm BHT; PPP, 2 g pomegranate peel powder; PABP, 4 g pomegranate aril bagasse  
 429 powder; PPAE, 6 g pomegranate peel powder aqueous extract; PABAE, 9 g pomegranate aril  
 430 bagasse powder aqueous extract.

431 Moisture, crude fat, crude protein, crude fibre and ash content for pomegranate fruit peel powder  
 432 were 11.03%, 1.42%, 6.83%, 15.80%, respectively.

433 Moisture, crude fat, crude protein, crude fibre and ash content for pomegranate aril bagasse  
 434 powder were 5.55%, 5.38%, 16.86%, 17.42%, 32.32% and 3.22%, respectively.

435 **Table 4. Effect of pomegranate peel powder, pomegranate aril bagasse powder, and their**  
 436 **aqueous extracts on instrumental colour analysis of chicken meat patties**

Treatments	Lightness (L*)	Redness (a*)	Yellowness (b*)
Control	56.46±1.09 <sup>a</sup>	4.05±0.67 <sup>a</sup>	11.84±0.93 <sup>a</sup>
BHT	55.49±0.64 <sup>a</sup>	4.47±0.44 <sup>a</sup>	11.87±0.50 <sup>a</sup>
PPP	50.08±3.07 <sup>b</sup>	4.58±0.74 <sup>a</sup>	11.14±0.35 <sup>a</sup>
PABP	51.35±4.55 <sup>b</sup>	4.76±0.40 <sup>a</sup>	11.06±1.09 <sup>a</sup>
PPAE	52.03±2.06 <sup>ab</sup>	4.65±0.76 <sup>a</sup>	11.24±0.76 <sup>a</sup>
PABAE	52.20±2.38 <sup>ab</sup>	4.51±0.88 <sup>a</sup>	11.63±1.07 <sup>a</sup>

437 (n=6, Mean ± SD)

438 Means with different superscripts in a column differ significantly (p≤0.05).

439 BHT, 100 ppm BHT; PPP, 2 g pomegranate peel powder; PABP, 4 g pomegranate aril bagasse  
 440 powder; PPAE, 6 g pomegranate peel powder aqueous extract; PABAE, 9 g pomegranate aril  
 441 bagasse powder aqueous extract.

442 **Table 5. Effect of pomegranate peel powder, pomegranate aril bagasse powder, and their aqueous extracts on texture profile,**  
 443 **firmness (N), and toughness (N-s) of chicken meat patties**

Treatments	Hardness (N)	Springiness	Cohesiveness	Gumminess (N)	Chewiness (N)	Firmness (N)	Toughness (N-s)
Control	38.87±4.78 <sup>c</sup>	0.89±0.06 <sup>a</sup>	0.54±0.15 <sup>a</sup>	20.68±5.33 <sup>b</sup>	18.59±5.92 <sup>ab</sup>	8.59±1.02 <sup>b</sup>	52.91±3.55 <sup>b</sup>
BHT	39.57 ±3.98 <sup>c</sup>	0.87±0.04 <sup>a</sup>	0.54±0.10 <sup>a</sup>	21.39±3.55 <sup>b</sup>	18.58±3.59 <sup>ab</sup>	9.99±1.77 <sup>b</sup>	48.06±4.48 <sup>b</sup>
PPP	54.72±2.76 <sup>a</sup>	0.87±0.04 <sup>a</sup>	0.50±0.10 <sup>a</sup>	27.26±5.54 <sup>a</sup>	23.79±5.92 <sup>a</sup>	13.55±2.36 <sup>a</sup>	73.71±3.23 <sup>a</sup>
PABP	48.31 ±4.70 <sup>b</sup>	0.87±0.04 <sup>a</sup>	0.46±0.02 <sup>a</sup>	22.19±2.47 <sup>b</sup>	19.30±2.80 <sup>ab</sup>	11.33±3.35 <sup>ab</sup>	55.97±3.00 <sup>b</sup>
PPAE	37.55 ±3.65 <sup>c</sup>	0.85±0.03 <sup>a</sup>	0.50±0.07 <sup>a</sup>	18.69±3.35 <sup>b</sup>	15.90±3.10 <sup>b</sup>	10.39±2.00 <sup>b</sup>	50.82±5.22 <sup>b</sup>
PABAE	36.25 ±3.07 <sup>c</sup>	0.84±0.03 <sup>a</sup>	0.48±0.05 <sup>a</sup>	17.54±1.50 <sup>b</sup>	14.76±1.51 <sup>b</sup>	9.29±2.69 <sup>b</sup>	50.23±5.87 <sup>b</sup>

444 (n=6, Mean ± SD)

445 Means with different superscripts in a column differ significantly (p≤0.05).

446 BHT, 100 ppm BHT; PPP, 2 g pomegranate peel powder; PABP, 4 g pomegranate aril bagasse powder; PPAE, 6 g pomegranate peel  
 447 powder aqueous extract; PABAE, 9 g pomegranate aril bagasse powder aqueous extract.

448

449

450

451 **Table 6. Effect of pomegranate peel powder, pomegranate aril bagasse powder, and their**  
 452 **aqueous extracts on pH and TBA value of chicken meat patties packaged under**  
 453 **aerobic conditions and stored at 4±2 °C**

Treatments	0 day	4 <sup>th</sup> day	8 <sup>th</sup> day	12 <sup>th</sup> day	16 <sup>th</sup> day
pH					
Control	6.30±0.10 <sup>aBC</sup>	6.21±0.13 <sup>aC</sup>	6.25±0.04 <sup>aBC</sup>	6.35±0.08 <sup>aAB</sup>	6.42±0.09 <sup>aA</sup>
BHT	6.25±0.08 <sup>aBC</sup>	6.14±0.09 <sup>aC</sup>	6.24±0.07 <sup>aBC</sup>	6.30±0.10 <sup>aB</sup>	6.42±0.09 <sup>aA</sup>
PPP	6.21±0.14 <sup>aBC</sup>	6.11±0.11 <sup>aC</sup>	6.20±0.08 <sup>aBC</sup>	6.27±0.06 <sup>aAB</sup>	6.36±0.09 <sup>aA</sup>
PABP	6.23±0.09 <sup>aBC</sup>	6.17±0.07 <sup>aC</sup>	6.20±0.08 <sup>aBC</sup>	6.28±0.08 <sup>aB</sup>	6.41±0.10 <sup>aA</sup>
PPAE	6.23±0.12 <sup>aBC</sup>	6.16±0.04 <sup>aC</sup>	6.23±0.09 <sup>aBC</sup>	6.28±0.07 <sup>aAB</sup>	6.39±0.11 <sup>aA</sup>
PABAE	6.24±0.10 <sup>aB</sup>	6.18±0.06 <sup>aB</sup>	6.23±0.13 <sup>aB</sup>	6.30±0.10 <sup>aAB</sup>	6.40±0.10 <sup>aA</sup>
TBA value (mg malonaldehyde/kg)					
Control	0.56±0.08 <sup>aD</sup>	0.63±0.26 <sup>aD</sup>	1.05±0.03 <sup>aC</sup>	1.64±0.27 <sup>aB</sup>	1.95±0.15 <sup>aA</sup>
BHT	0.53±0.10 <sup>aC</sup>	0.61±0.22 <sup>abC</sup>	0.87±0.04 <sup>abB</sup>	1.16±0.24 <sup>baA</sup>	1.40±0.32 <sup>baA</sup>
PPP	0.24±0.07 <sup>cC</sup>	0.35±0.10 <sup>cC</sup>	0.72±0.14 <sup>baA</sup>	0.78±0.20 <sup>caA</sup>	0.81±0.11 <sup>caA</sup>
PABP	0.30±0.10 <sup>cB</sup>	0.40±0.12 <sup>bcB</sup>	0.75±0.29 <sup>baA</sup>	0.83±0.22 <sup>caA</sup>	0.84±0.06 <sup>caA</sup>
PPAE	0.41±0.07 <sup>bbB</sup>	0.50±0.07 <sup>abcB</sup>	0.75±0.27 <sup>baA</sup>	0.81±0.05 <sup>caA</sup>	0.83±0.10 <sup>caA</sup>
PABAE	0.46±0.13 <sup>abC</sup>	0.56±0.12 <sup>abBC</sup>	0.81±0.16 <sup>baB</sup>	0.81±0.05 <sup>caB</sup>	1.00±0.39 <sup>caA</sup>

454 (n=6, Mean ± SD)  
 455 BHT, 100 ppm BHT; PPP, 2 g pomegranate peel powder; PABP, 4 g pomegranate aril bagasse  
 456 powder; PPAE, 6 g pomegranate peel powder aqueous extract; PABAE, 9 g pomegranate aril  
 457 bagasse powder aqueous extract.  
 458 Means with different small superscripts within a column and capital superscripts within a row for  
 459 a particular parameter differ significantly (p≤0.05).

466 **Table 7. Effect of pomegranate peel powder, pomegranate aril bagasse powder, and their**  
 467 **aqueous extracts on microbial counts of chicken meat patties packaged in aerobic**  
 468 **conditions and stored at 4±2 °C**

Treatments	0 day	4 <sup>th</sup> day	8 <sup>th</sup> day	12 <sup>th</sup> day	16 <sup>th</sup> day
Total plate count (Log CFU/g)					
Control	2.43±0.48 <sup>aE</sup>	3.46±0.56 <sup>aD</sup>	4.09±0.35 <sup>aC</sup>	5.09±0.37 <sup>aB</sup>	5.72±0.36 <sup>aA</sup>
BHT	2.51±0.22 <sup>aD</sup>	3.36±0.58 <sup>aC</sup>	3.81±0.41 <sup>abC</sup>	4.80±0.72 <sup>abB</sup>	5.46±0.59 <sup>abA</sup>
PPP	2.35±0.65 <sup>aD</sup>	2.96±0.41 <sup>aC</sup>	3.40±0.40 <sup>bcC</sup>	4.07±0.49 <sup>cbB</sup>	4.70±0.23 <sup>caA</sup>
PABP	2.43±0.63 <sup>aC</sup>	2.96±0.54 <sup>aC</sup>	3.60±0.55 <sup>abB</sup>	4.28±0.42 <sup>bcA</sup>	4.85±0.41 <sup>caA</sup>
PPAE	2.24±0.12 <sup>aE</sup>	3.06±0.56 <sup>aD</sup>	3.70±0.49 <sup>abC</sup>	4.51±0.46 <sup>abcB</sup>	5.11±0.45 <sup>bcA</sup>
PABAE	2.28±0.09 <sup>aE</sup>	3.04±0.38 <sup>aD</sup>	3.72±0.43 <sup>abC</sup>	4.60±0.55 <sup>abcB</sup>	5.16±0.37 <sup>bcA</sup>
Psychrotrophic count (Log CFU/g)					
Control	ND	1.30±0.69 <sup>aC</sup>	1.96±0.28 <sup>aB</sup>	2.51±0.57 <sup>aAB</sup>	3.10±0.42 <sup>aA</sup>
BHT	ND	1.12±0.61 <sup>aC</sup>	1.79±0.39 <sup>aB</sup>	2.46±0.63 <sup>aB</sup>	2.96±0.36 <sup>aA</sup>
PPP	ND	0.91±0.75 <sup>aC</sup>	1.60±0.74 <sup>aBC</sup>	2.08±0.41 <sup>aAB</sup>	2.40±0.28 <sup>baA</sup>
PABP	ND	1.01±0.76 <sup>aC</sup>	1.69±0.97 <sup>aBC</sup>	2.17±0.30 <sup>aAB</sup>	2.65±0.26 <sup>abA</sup>
PPAE	ND	1.02±0.37 <sup>aD</sup>	1.69±0.38 <sup>aC</sup>	2.24±0.36 <sup>aB</sup>	2.75±0.37 <sup>abA</sup>
PABAE	ND	1.13±0.67 <sup>aC</sup>	1.70±0.44 <sup>aBC</sup>	2.20±0.40 <sup>aAB</sup>	2.80±0.47 <sup>abA</sup>
Thermophilic count (Log CFU/g)					
Control	1.36±0.39 <sup>aE</sup>	1.98±0.31 <sup>aD</sup>	2.61±0.61 <sup>aC</sup>	3.09±0.30 <sup>aA</sup>	3.63±0.32 <sup>aA</sup>
BHT	1.23±0.54 <sup>aC</sup>	1.81±0.49 <sup>aC</sup>	2.49±0.64 <sup>aB</sup>	2.91±0.36 <sup>abAB</sup>	3.42±0.40 <sup>abA</sup>
PPP	1.21±0.74 <sup>aD</sup>	1.68±0.81 <sup>aCD</sup>	2.14±0.38 <sup>aBC</sup>	2.60±0.41 <sup>baB</sup>	2.96±0.22 <sup>caA</sup>
PABP	1.25±0.75 <sup>aC</sup>	1.69±0.41 <sup>aC</sup>	2.37±0.58 <sup>aB</sup>	2.81±0.38 <sup>abAB</sup>	3.10±0.42 <sup>bcA</sup>
PPAE	1.26±0.74 <sup>aC</sup>	1.71±0.38 <sup>aBC</sup>	2.41±0.56 <sup>aB</sup>	2.82±0.38 <sup>abA</sup>	3.25±0.26 <sup>abcA</sup>
PABAE	1.26±0.74 <sup>aD</sup>	1.72±0.97 <sup>aCD</sup>	2.42±0.52 <sup>aBC</sup>	2.84±0.38 <sup>abAB</sup>	3.30±0.25 <sup>abcA</sup>

469 (n=6, Mean ± SD)

470 ND; Not detected.

471 BHT, 100 ppm BHT; PPP, 2 g pomegranate peel powder; PABP, 4 g pomegranate aril bagasse  
 472 powder; PPAE, 6 g pomegranate peel powder aqueous extract; PABAE, 9 g pomegranate aril  
 473 bagasse powder aqueous extract.

474 Means with different small superscripts within a column and capital superscripts within a row for  
 475 a particular parameter differ significantly (p≤0.05).