

1
2
3
4

TITLE PAGE
- Food Science of Animal Resources -
Upload this completed form to website with submission

| ARTICLE INFORMATION | Fill in information in each box below | |
|--|--|--|
| Article Title | Current Issues and Technical Advances in Cultured Meat Production: A Review | |
| Running Title (within 10 words) | A Review of Cultured Meat Production | |
| Author | Tae Kyung Hong ^{1,†} , Dong-Min Shin ^{2,†} , Joonhyuk Choi ^{1,†} , Jeong Tae Do ^{1,*} , and Sung Gu Han ^{2,*} | |
| Affiliation | ¹ Department of Stem Cell and Regenerative Biotechnology, KU Institute of Science and Technology, Konkuk University, Seoul 05029, Republic of Korea ² Department of Food Science and Biotechnology of Animal Resources, Konkuk University, Seoul 05029, Republic of Korea | |
| Special remarks – if authors have additional information to inform the editorial office | [†] These authors contributed equally to this work. [*] Corresponding authors | |
| ORCID (All authors must have ORCID) https://orcid.org | Tae Kyung Hong (0000-0002-5237-2266) Dong-Min Shin (0000-0003-2755-433X) Joonhyuk Choi (0000-0001-7677-5978) Jeong Tae Do (0000-0001-6721-1441) Sung Gu Han (0000-0002-1485-861X) | |
| Conflicts of interest List any present or potential conflicts of interest for all authors. (This field may be published.) | The authors declare no potential conflict of interest. | |
| Acknowledgements State funding sources (grants, funding sources, equipment, and supplies). Include name and number of grant if available. (This field may be published.) | This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) of the Republic of Korea (grant nos. 2016M3A9B6946835 and 2020R1A2C3007562). | |
| Author's contributions (This field may be published.) | Conceptualization: Do JT, Han SG. Writing - original draft: Hong TK, Shin D-M, Choi J. Do JT, Han SG. Writing - review & editing: Do JT, Han SG. | |
| Ethics approval (IRB/IACUC) (This field may be published.) | This manuscript does not require IRB/IACUC approval because there are no human and animal participants. | |

5
6

CORRESPONDING AUTHOR CONTACT INFORMATION

| For the <u>corresponding</u> author (responsible for correspondence, proofreading, and reprints) | Fill in information in each box below | Fill in information in each box below |
|---|--|--|
| First name, middle initial, last name | Jeong Tae Do | Sung Gu Han |

| | | |
|--|---|---|
| Email address – this is where your proofs will be sent | dojt@konkuk.ac.kr | hansg@konkuk.ac.kr |
| Secondary Email address | | |
| Postal address | 120 Neungdong-ro, Gwangjin-gu, #418 Animal Science Building, Konkuk University, Seoul 05029, Korea | 120 Neungdong-ro, Gwangjin-gu, #618 Animal Science Building, Konkuk University, Seoul 05029, Korea |
| Cell phone number | | |
| Office phone number | 82-2-450-3673 | 82-2-450-0526 |
| Fax number | 82-2-455-1044 | 82-2-4556-1044 |

7

8

ACCEPTED

9 **Abstract**

10 As the global population grows, we need a stable protein supply to meet the demands.
11 Although plant-derived protein sources are widely available, animal meat maintains its
12 popularity as a high-quality and savory protein source. Recently, cultured meat, also known as
13 *in vitro* meat, has been suggested as a meat analog produced through *in vitro* cell culture
14 technology. Cultured meat has several advantages over conventional meat, such as
15 environmental protection, disease prevention, and animal welfare. However, cultured meat
16 manufacturing is an emerging technology; thus, its further and dynamic development would be
17 pivotal. Commercialization of cultured meat to the public will take a long time but cultured
18 meat undoubtedly will come to our table someday. Here, we discuss the social and economic
19 aspects of cultured meat production as well as the recent technical advances in cultured meat
20 technology.

21

22 Key words: cultured meat, *in vitro* meat, livestock farming, myogenic satellite cells,
23 alternative protein sources



24 **Introduction**

25 The current global population is 7.3 billion and is estimated to reach 10 billion by 2050 (UN,
26 2019). Consequently, such an increase might result in a protein demand twice as much as the
27 current protein production (Godfray et al., 2019). Since conventional meat production systems
28 such as animal agriculture are no longer sustainable, scientists have been searching for
29 alternative protein sources (Goodwin and Shoulders, 2013). Early attempts for meat
30 alternatives were focused on plant-based meat analogues with the use of soy-, wheat-, or fungi-
31 based protein sources (Hoek et al., 2004; Sadler, 2004). Only recently researchers have tried to
32 use cultured muscle cells as alternatives to real meat. Cultured meat, also known as *in vitro*
33 meat, is a meat analog produced using *in vitro* cell culture technology where the animal cells
34 are primarily skeletal muscle-derived cells isolated through muscle biopsy and from
35 slaughtered livestock (Choi et al., 2021; Datar and Betti, 2010).

36 Cultured meat technologies have received a lot of attention because many people think that
37 this technology could supplement or partially replace conventional animal production systems
38 (Post et al., 2020). In fact, conventional animal production system has been the most important
39 part of agriculture. Nonetheless, during last few decades, people and researchers have raised
40 concerns about the conventional animal production system because it may cause several
41 problems, including environmental and social concerns, and animal welfare issues (Post, 2012).

42 The first cultured meat was produced in 2013 by Mark Post from the Maastricht University,
43 Netherlands, from primary bovine skeletal muscle cells. Since then, several university
44 laboratories and companies have entered this research field (Stephens et al., 2018). Later,
45 another US-based start-up company, Memphis Meats, produced several forms of cultured meat
46 products such as meatballs, beef fajita, chicken, and duck (Stephens et al., 2018). In addition,

47 Just, Inc., a vegan cookie dough and mayonnaise company, announced that they would debut
48 cultured chicken nuggets. Further, a start-up company, Modern Meadow, developed a steak
49 chip made of cultured meat combined with a hydrogel (Marga, 2016; Stephens et al., 2018).
50 Since the introduction of the first cultured meat patty in 2013, several private companies have
51 been founded and focusing on cultured meat production (Choudhury et al., 2020).

52 Although there are many technological difficulties associated with cultured meat area, at
53 least some of the global problems could be potentially solved through the successful
54 development of this technology (Table 1). Therefore, in this review, we summarized the current
55 issues and technological development about cultured meat production, particularly focusing on
56 three areas: 1) social and economical aspects of cultured meat, 2) biological basis underlying
57 the meat culture of various livestock, and 3) technological approaches for cultured meat
58 production.

59

60 **1. Social and economic aspects of cultured meat production systems**

61 **1.1 Economic sustainability of cultured meat**

62 Cultured meat system requires less use of water, land, feed grain, and energy compared with
63 traditional livestock system (Tuomisto and Teixeira de Mattos, 2011). In addition, cultured
64 meat system may exhibit a higher conversion rate transformed into edible meat than traditional
65 livestock system that exhibits 5-25% conversion rate (Alexander, 2011; Bhat and Hina, 2011).
66 Thus, cultured meat could be an ideal alternative due to its potential sustainability and limited
67 environmental effects. For example, a 20 m³ bioreactor, the largest size for cultured meat
68 production today, could produce 25,600 kg of cultured meat per year (Van der Weele and
69 Tramper, 2014). Assuming no loss during the cultured meat production process, this represents

70 an estimated supply of cultured meat for 2,560 people per year (Van der Weele and Tramper,
71 2014). The calculation on feeding 2,560 people is based on Van der Weele and Tramper (2014)
72 who assumed that everybody in the world will eat 25–30 grams of cultured meat per person
73 per day (10 kg/year). Considering that such production requires only a few hours of labor per
74 day to maintain the bioreactor, cultured meat production is a potentially low-cost alternative to
75 the current livestock system for meat production (Bhat et al., 2014). In addition, it was reported
76 that the price of cultured meat burger decreased from \$325,000 to \$11.36 per burger or \$80 per
77 kilogram of meat within 2 years (Crew, 2015). Another economic benefits could be found in
78 the distribution of cultured meat. By locating cultured meat production facilities close to the
79 cities, the transport cost can be largely decreased (Bhat et al., 2015). Additionally, in terms of
80 food waste, traditional meat industry has big problem in waste management because whole
81 carcass cannot be used for consumption. However, culture meat system can provide prime cut
82 alone for consumption and further processing and that will be an substantial economical benefit
83 (Stephens et al., 2018).

84

85 **1.2 Environmental sustainability of cultured meat**

86 The current livestock system negatively influences the environment, causing environmental
87 sustainability concerns. Although the water used by livestock farming mostly returns to the
88 environment, a significant part of it becomes polluted or evaporates (Melvin, 1995). This
89 pollution is caused by livestock and feed production, as well as product processing, in turn
90 increasing the demand for water (Steinfeld et al., 2006). In order to produce 1 kg of beef, 15,495
91 liters of water would be required, and 99% of such water consumption is used for the growth
92 of grain and roughages (e.g., pasture, dry hay and silage) (Hoekstra and Chapagain, 2006).

93 Only 1% of water (about 155 liters) is used for drinking and servicing to livestock. The demand
94 is mostly attributed to the drinking water requirement for the animals, as well as crop and plant
95 growth (Chriki and Hocquette, 2020). Both water pollution and consumption might lead to the
96 destruction of biodiversity through destruction of wildlife habitats (Steinfeld et al., 2006).
97 However, cultured meat technology uses approximately 82–96% less water than traditional
98 livestock farming (Tuomisto and Teixeira de Mattos, 2011).

99 In general, livestock production requires 30% of the total land surface—33% of cultivated
100 land for livestock feed and 26% for pasture (Steinfeld et al., 2006). However, cultured meat
101 production systems use only 1% of the land required for traditional livestock production
102 systems (Alexander et al., 2017; Tuomisto and Teixeira de Mattos, 2011). Nevertheless, this
103 assumption is restricted to the production of an algae-based culture medium biomass, and the
104 expense and efficiency of producing different culture media are therefore uncertain. Although
105 cultured meat production systems require lesser land than traditional livestock systems, the
106 cultured meat production system requires at least four times more energy than traditional
107 livestock (Alexander et al., 2017). In detail, cultured meat requires 18-25 GJ/t of direct energy
108 (Tuomisto and Teixeira de Mattos, 2011), while 4.5 GJ/t of direct energy is required to produce
109 traditional meat (MacLeod et al., 2013).

110 Livestock production consumes direct energy, such as lighting, heating, and cooling, while
111 cultured meat production systems require energy for muscle cell culture, as well as for the
112 sterilization and hydrolysis of biomass material required in the cell culture media (Tuomisto
113 and Teixeira de Mattos, 2011).

114 Livestock provides a quarter of all the protein content (and 15% of energy) consumed in
115 food, and also contributes to 18% of the global greenhouse gas and 37% of methane emissions
116 into the atmosphere, the values of which are higher than those of global transportation (FAO,

117 2012; Steinfeld et al., 2006). Cultured meat production would assumably affect less the
118 environment compared to conventional farming. In particular, reducing greenhouse gas
119 emissions would be a significant advantage of cultured meat production. Another potential
120 environment-related advantage of cultured meat production could be the lower land use
121 compared to conventional livestock farming, especially in the case of ruminants (Chriki and
122 Hocquette, 2020).

123

124 **1.3 Animal welfare and cultured meat**

125 Recently, approximately 56 billion animals are slaughtered for their meat every year
126 (Dorovskikh, 2015). Hence, the traditional livestock production-related animal welfare is a
127 major worldwide ethic agenda. Cultured meat production systems have been raised as good
128 alternatives to the current meat production systems (Post, 2012). Cultured meat could be an
129 attractive option for vegetarians, vegans, and opponents who reject meat consumption for
130 ethical reasons (Hopkins and Dacey, 2008). According to a previous article, we could expect
131 the following effects of widespread cultured meat production: 1) a significant reduction in
132 animal use, 2) a great reduction in animal suffering, and 3) a variety of cultured meat sources,
133 including those of wild animals (Bhat et al., 2014).

134

135 **1.4 Cultured meat-related consumer acceptance and ethical issues**

136 Despite the potential animal welfare- and environment-related merits of cultured meat, the
137 mercantile success of cultured meat greatly depends on consumer perception and various
138 societal concerns, including naturalness, food safety and security issues, framing effect,
139 legislation, religion, and ethics (Chriki and Hocquette, 2020; Mancini and Antonioli, 2020).

140 Hence, the consumer acceptance of cultured meat is highly important but could be controversial.
141 One of the most common cultured meat-related hurdles is its artificial nature. Consumers
142 usually do not easily accept new technologies, such as genetically modified organisms, when
143 they have limited information about the given technology (Bánáti, 2011). In addition, framing
144 effects on cultured meat significantly contribute to consumer attitude, beliefs, and behavioral
145 intention to cultured meat (Bryant and Dillard, 2019). However, changes in consumer
146 perception by providing positive information could make consumers try, buy, and pay for
147 cultured meat. Continuous evaluation of the changes in consumer perception over time would
148 thus be necessary.

149 The regulatory structures are important for building consumers' trust towards cultured meat
150 production and cultured meat itself, including safety and nutritional composition (Laestadius
151 and Caldwell, 2015). Several reports focus on the regulation of cultured meat in the United
152 States and the European Union (Petetin, 2014; Schneider, 2012). However, it is difficult to
153 establish cultured meat-related regulations due to the currently available insufficient
154 information and incomplete technology for cultured meat (Stephens et al., 2018).

155 There is controversy concerning cultured meat in several religious communities, including
156 Jews, Muslims, and Hindus, due to its nebulous status (Chriki and Hocquette, 2020). In a
157 cultured meat-related consumer acceptance survey targeting 3,030 participants, including Jews,
158 Muslims, and Hindus, most participants responded that they would be willing to eat cultured
159 meat (Bryant et al., 2019). However, religious duties, such as dietary laws (Kosher, Halal, beef-
160 eating restrictions in Hinduism), still need to be discussed (Bryant, 2020).

161 In the case of food choices, ethical issues become increasingly important. Although cultured
162 meat technology gets closer to actual commercial availability, it is obvious that ethical concerns
163 of cultured meat is not completely solved yet (Dilworth et al., 2015). There are some arguments

164 amongst consumers regarding the ethical issues of cultured meat. Advocates believe that
165 cultured meat systems demand significantly fewer animals for meat production than traditional
166 livestock and could also contribute to stop animal suffering, such as confining in tight space or
167 slaughtering under cruel conditions (Chriki and Hocquette, 2020). In addition, cultured meat
168 might be preferred by people who are interested in reducing their meat consumption for ethical
169 reasons, including vegetarians and vegans (Hopkins and Dacey, 2008). According to a previous
170 report, cultured meat could have a positive impact on a carbon footprint, and this makes a
171 potentially effective strategy to improve awareness of cultured meat (Tomiyama et al., 2020).
172 However, despite of potential advantages of introducing cultured meat, many people concern
173 about food safety regarding unnaturalness perception of cultured meat (Laestadius, 2015;
174 Verbeke et al., 2015). Moreover, some have concerned that cultured meat may aggravate
175 consumer inequality between the rich and the poor (Bonny et al., 2015; Cole and Morgan, 2013;
176 Stephens et al., 2018).

177

178 **2. Biological basis underlying the cultured meat production of various livestock**

179 Currently, 32 cultured meat companies exist worldwide, focusing on cultured beef (25%),
180 poultry (22%), pork (19%), seafood (19%), and other exotic meats (15%), such as mouse,
181 kangaroo, and horse (Choudhury et al., 2020). Most of these companies are based in North
182 America (40%), followed by Asia (31%) and Europe (25%). Substantial amount of capital has
183 been invested in cultured meat-related research and development in the past 5 years.
184 Approximately \$320 million have presumably been invested in beef and pork (75%) as well as
185 in seafood production (25%) (Choudhury et al., 2020).

186

187 **2.1 Characteristics of satellite cells**

188 Meat from industrial animals, including cattle, pigs, poultry, and fish, consists mainly of
189 skeletal muscles, fibroblasts, and adipose cells (Dodson et al., 2015). In addition, meat can also
190 provide vitamin B12 and heme iron, which are essential for human nutrition. Skeletal muscle
191 cells are multinucleated and striated cells, which fulfill the basic function of muscle contraction.
192 Moreover, skeletal muscles are able to regenerate and recover minor damage in the muscle
193 tissue (Laumonier and Menetrey, 2016). Their self-renewal ability is due to stem cells, i.e.,
194 satellite cells that reside within the skeletal muscle tissue. As the number of satellite cells
195 reportedly remains constant after multiple injuries, these cells are considered stem cells that
196 could most certainly be maintained by self-renewal (Shi and Garry, 2006). Under normal
197 conditions, satellite cells are quiescent but could be activated by intrinsic or extrinsic cues, such
198 as muscle injury. The quiescent state of satellite cells is maintained by the negative cell cycle
199 and growth factor regulation and the expression of tumor suppressors, such as retinoblastoma
200 protein (Rb) (Dumont et al., 2015). Up-regulated Notch signaling is also a quiescent satellite
201 cell marker. Therefore, Notch down-regulation is a prerequisite for myogenic differentiation
202 (Brack et al., 2008). Moreover, myogenic factor 5 (Myf5), myogenic determination (MyoD),
203 and myogenin (Myog) are critical factors that are expressed from activated satellite cell under
204 muscle stimulus, and therefore, they are committed myogenic progenitor markers (Dumont et
205 al., 2015). Active proliferating satellite cells (quiescent cells) – expressing high levels of paired
206 box 7 (Pax7), and concurrently negative for Myf5 and MyoD – are crucial for maintaining
207 stemness (Figure 1).

208 Satellite cells were first isolated *in vitro* by Richard Bischoff in 1974 (Bischoff, 1974). Since
209 the discovery of muscle satellite cell isolation and proliferation methods (Bischoff, 1975),

210 various modified protocols have been developed to isolate satellite cells more efficiently from
211 multiple livestock, such as chicken (Yablonka-Reuveni et al., 1987), horse (Greene and Raub,
212 1992), cow (Dodson et al., 1987), sheep (Dodson et al., 1986), fish (Greenlee et al., 1995), and
213 pig (Doumit and Merkel, 1992). Using isolated satellite cells, researchers were able to
214 understand further the underlying processes of muscle formation and development (Allen et al.,
215 1979). Recently, scientists have used stem cells and muscle culture technology to develop lab-
216 grown meat, cultured in a laboratory incubator using isolated skeletal muscle and satellite cells
217 (Bischoff, 1975).

218 Although not yet on the market and much more expensive than farmed meat, cultured meat
219 offers multiple advantages over conventional meat. Cultured meat is a clean meat, free of
220 possible pathogens (Kadim et al., 2015), environmentally friendly due to its lack of need for
221 large space to raise livestock, and significantly less global gas emission compared to
222 conventional livestock farming (Tuomisto and Teixeira de Mattos, 2011). Several startup
223 companies are currently emerging around the world, and research on the production of high-
224 quality, low-cost culture meat production is underway (Table 2).

225

226 **2.2 Chicken meat**

227 Chicken muscle satellite cell *in vitro* isolation and differentiation was described in 1983 by
228 Matsuda et al. (Matsuda et al., 1983). Yablonka-Reuveni et al obtained chicken pectoralis cells
229 differentiated from satellite cells, isolated by centrifugation through a Percoll density gradient
230 (Yablonka-Reuveni et al., 1987). Satellite cells play a crucial role in the muscle growth of post-
231 hatch broiler chicken and in muscle maintenance and repair after muscle injury. Since the stem
232 cell properties of muscle satellite cells, the proliferation and differentiation potential of chicken

233 satellite cells have been evaluated in detail. In general, when skeletal muscle is damaged, new
234 muscle fibers derived from pre-existing quiescent satellite cells replace the damaged area and
235 reconstruct the muscle structure (Feldman and Stockdale, 1991). Feldman and Stockdale et al.
236 suggested that chicken satellite cells isolated from the fast muscle (pectoralis major) part would
237 be differentiated only into fast fibers, whereas satellite cells isolated from the slow muscle
238 (anterior latissimus dorsi) part could mostly differentiate into fast muscles but, to a small extent,
239 also into slow muscles. (Feldman and Stockdale, 1991).

240 Cultured meat has not yet been formally commercialized and sold, but many companies have
241 promoted it as various prototype foods such as hamburgers, bacon, and nuggets. Artificial
242 chicken meat was presented by JUST, a vegan food company, in 2018, through a promotional
243 video (JUST, 2018). They showed a footage of clean chicken meat that was created using cell
244 cultures (JUST, 2018). Moreover, JUST successfully manufactured a cell-cultured chicken
245 nugget product in 2019 at the cost of 50 dollars per nugget (Savvides, 2020). A food technology
246 company, Memphis Meats (Berkeley, California), published a similar promotion video
247 introducing the concept of a cultured meat product in 2016 (Meats, 2016a). In the following
248 year, Memphis Meats was able to successfully manufacture and introduce a cultured chicken
249 meat product (Meats, 2017). Future Meat Technologies, a start-up company founded in 2018
250 and based in Israel, also created cell-cultured chicken meat. This company managed to reduce
251 production costs to 150 dollars per pound of chicken (Lucas, 2019). However, even these small
252 pieces of foods, such as artificial nuggets, require FDA and USDA approval (Savvides, 2020).
253 The commercialization of these products has not yet realized.

254

255 **2.3 Duck meat**

256 During embryonic development, proliferating myoblasts differentiate into myotubes,
257 followed by further maturation and differentiation into mature muscle fibers (Braun and Gautel,
258 2011). Adal and Cheng studied the structure of duck muscle cells as early as 1980 and showed
259 that the duck muscle spindle consists of several muscle fibers and a capsule surrounding them
260 (Adal and Cheng, 1980). In 1986, stromal mesenchymal cells in the iris of a duck reportedly
261 migrated towards the muscle of the iris and became iridial skeletal muscles (Yamashita and
262 Sohal, 1986). Muscle-specific microRNAs, called MyomiRs, are expressed in the muscle cells,
263 although they are also expressed in several other tissues (McCarthy, 2008). Li et al. found
264 detected 279 novel miRNAs in the breast muscle of ducks, indicating the importance of
265 miRNAs in muscle development and maturation (Li et al., 2020). Among these, miRNA-1 and
266 miRNA-133 have been suggested to be crucial factors for duck skeletal muscle proliferation
267 and differentiation. miRNA-1 reportedly promoted myogenesis by targeting the transcriptional
268 repressor histone deacetylase 4 (HDAC4), and miRNA-133 reportedly inhibited serum
269 response factor (SRF) and TGFBR1 expression, increasing myoblast proliferation (Wu et al.,
270 2019).

271 During duck embryonic development, MyoD expression in both the breast and leg muscles
272 tended to increase gradually, and MyoD expression level in the breast muscle was higher than
273 that in the leg muscle (Li et al., 2014; Li et al., 2010). However, Li et al. suggested that MyoD
274 expression in the breast muscle was consistent but decreased in leg muscle during early
275 embryonic development (Li et al., 2014; Li et al., 2010). They also showed that the MyoD and
276 Myf6 gene expressions correlated with that in the leg muscle. However, insulin-like growth
277 factor-1 (IGF-1) induced the expression of MyoD and Myf5 and increased muscle hypertrophy
278 (Liu et al., 2012). IGF-1 is known to stimulate skeletal muscle (Musaro et al., 2001).

279 Similarly to cultured chicken meat, Memphis Meats also produced cultured duck meat,

280 which was cooked and presented, followed by product tasting (Meats, 2017). Moreover, a
281 French start-up company Gourmey, was able to cultivate duck egg cells with slightly adjusted
282 nutrients to mimic the effect of force-feeding in order to create artificial foie gras, which they
283 refer to as ‘ethical foie gras’ (Gourmey; Southey, 2020). In 2020, the vegan food company
284 JUST managed to produce duck chorizo and pâté completely based on cultured duck cells
285 (Purdy, 2020).

286

287 **2.4 Beef**

288 Beef has long been studied in various ways. Several biological aspects of muscles have been
289 studied for the basic understanding of the mechanisms underlying cellular proliferation, and
290 many scientific findings have been reported related to muscle development and proliferation
291 in meat animals (Allen et al., 1979; Wojtczak, 1979; Dayton and White, 2008). In meat animals,
292 the fetal stage of muscle development is crucial since the number of muscle fibers does not
293 change after birth (Zhu et al., 2004). Therefore, the postnatal muscle develops by enlarging the
294 muscle fiber size (Karunaratne et al., 2005; Stickland, 1978). Satellite cells located under the
295 basal lamina of the muscle fibers are crucial for muscle growth after birth. Major satellite cells
296 differentiate into the myogenic lineage, but a small population of satellite cells could also
297 differentiate into fibroblasts or adipocytes, which comprise the skeletal muscle tissue.
298 Understanding the mechanisms underlying satellite cell-related muscle growth and
299 differentiation would enable further improvements in cultured meat production (Rubin, 2019).

300 Controlling nutrient supplementation and several signaling factors is important for skeletal
301 muscle growth and marbling. For example, skeletal muscle growth is enhanced by the
302 activation of the Wnt signaling, while it inhibits adipogenesis (Du et al.,

2010). β -catenin, which is stabilized by Wnt signaling, positively regulates myogenic genes, such as Pax3, MyoD, and Myf5 (Ridgeway and Skerjanc, 2001). For commercial applications, marbling could be controlled by the activation and repression of the Wnt/ β -catenin signaling during culture, in order to produce higher-quality meat.

Mosa Meat, a Dutch start-up company, was the first to promote cultured beef in public. This beef was generated by culturing and differentiating stem cells obtained from a cow and was formulated into muscle strips. Mosa Meat cooked the cultured meat at a conference, then organized a tasting party (BBC news, 2013). Mosa Meat now creates cell-cultured meats that are more cost-effective than before, and it has now developed a bovine serum-free medium (Kateman, 2020). Memphis Meats, a start-up company based in California, showed the first meatballs made from cell-cultured beef in 2016. The company is now building a pilot plant for cultured beef and chicken meat (Meats, 2016; Shaffer, 2020)

2.5 Pork

Doumit and Merkel suggested that porcine myogenic satellite cells could be isolated from porcine skeletal muscle and developed an optimized medium for porcine satellite cells (Doumit and Merkel, 1992). This culture condition has been improved with slight modifications (Mau et al., 2008; Metzger et al., 2020). For the *in vitro* culture of skeletal muscle, satellite cells or muscle fibers could be isolated from muscle tissues to induce growth and differentiation (Mau et al., 2008; Metzger et al., 2020). Pax7 is a critical marker for functional satellite cells in several species, including mice, humans, cattle, and pigs (Ding et al., 2017). IGF-1 also affects pig satellite cells through the mTOR pathway (Han et al., 2008). CD56 and CD34 have been suggested as myogenic cell markers in swine skeletal muscles (Perruchot et al., 2013). LDHA

326 and COPB1 were also suggested to be involved in pig muscle development (Qiu et al., 2010).
327 RNA-seq analysis using the pig longissimus dorsi muscle revealed that long non-coding RNAs
328 are involved in muscle growth and fat deposition (Chen et al., 2019). As shown in mice, the
329 number of satellite cells decreases with age and during long-term culturing due to the loss of
330 their self-renewal and differentiation potentials (Ding et al., 2017).

331 Meatable, a Dutch start-up company, produced cell-cultured pork meat using stem cell
332 technology, which allowed the company to easily extract specific cell types required to produce
333 meat (Brodwin, 2018a). Another startup company in San Francisco, New Age Meats,
334 successfully produced prototype pork sausage made from fat and muscle cell culture from a
335 live pig sample (Brodwin, 2018b).

336

337 **3. Technical approaches for cultured meat production**

338 Tissue engineering-based cultured meat production largely depends on large-scale cell
339 culture technologies, which could provide a significant amount of cells, allowing meat
340 production (Verbruggen et al., 2018). Large-scale cell production systems also aim at
341 producing as many cells as possible with the least of the required resources. Minimal handling
342 and a short culture period for a sufficient number of harvested cells are also commonly
343 considered factors for efficient cell mass production (Moritz et al., 2015). Several cell types
344 are potentially viable options for cultured meat production, including myogenic satellite cells,
345 embryonic stem cells, and induced pluripotent stem cells (Kadim et al., 2015). Among these
346 various cell types, myogenic satellite cells are widely used as the promising option due to their
347 efficient differentiation into myotubes (Arshad et al., 2017). A variety of methods and
348 bioreactors are used to expand anchorage-dependent cells (Merten, 2015). Each technology has

349 its own merits, but in common, these platforms provide an attachment surface area for the cells
350 while assuring gas and nutrient exchange in parallel (Tavassoli et al., 2018).

351

352 **3.1 Multi-tray systems**

353 As cell culture is a major step in the production of cultured meat, the choice of the
354 appropriate culture dish or vessel is pivotal. T-flasks commonly used in cell culture provide a
355 surface area of 20–225 cm². In the case of large-scale cultures that require a significantly larger
356 surface area than that, multiple T-flasks could be used. A multi-tray system has been developed
357 as an alternative high-surface area provided within a single unit. Although this system has
358 multiple trays that provide multiple cell attachment surfaces, handling multiple T-flasks might
359 be labor-intensive as each T-flask must be managed individually (Rafiq et al., 2013).

360

361 **3.2 Roller bottles**

362 Roller bottles were devised by Gey in 1933, aiming at low-cost maintenance of a large
363 number of cell populations while using less culture medium (Melero-Martin and Al-Rubeai,
364 2007) (Gey, 1933). Roller bottles, placed in a gas-tight chamber or a case with no chamber,
365 could be sealed to keep the cells and medium from drying. This system also requires a slow
366 driving mechanism, allowing the bottles with cells to slowly rotate, enabling the medium to
367 cover cells evenly, and allowing greater gas exchange (Melero-Martin and Al-Rubeai, 2007).
368 Roller bottles could offer a surface attachment area of up to 350,000 cm². Compared to T-flasks
369 or multi-tray cultures, roller bottles provide superior applications for anchorage (Rafiq et al.,
370 2013). However, the real-time monitoring of the roller bottle system is difficult, and handling
371 several roller bottles simultaneously is laborious (Tavassoli et al., 2018). To overcome these

372 shortcomings, relevant efforts have been made to automate the roller bottle-based culture
373 process (Kunitake et al., 1997). Roller bottles have been used to culture chicken muscle cells
374 in large scale (Wesson et al., 1949) which may be applied for chicken meat production.
375 According to the United States Department of Agriculture, roller bottle incubator systems
376 drastically improved swine muscle cell production output, providing enough cells for 3D
377 fabrication of cellular sheets for *in vitro* meat engineering (Marga, 2012).

378

379 **3.3 Microcarriers**

380 Culturing cells in suspension provide more output than monolayer culture systems, but
381 adhesion to a specific culture surface is crucial for the anchorage-dependent cells to proliferate
382 without losing their cellular properties (Grinnell, 1978). In order to mass-culture anchorage-
383 dependent cells, microcarriers are used to establish suspension cultures (Rafiq et al., 2013). In
384 1967, Van Wezel described the concept of “micro-carriers” using dextran particles for
385 developing large-scale cell cultures in a stirred suspension. These dextran particles are micro-
386 sized beads that display positively-charged surfaces and attract animal cells that contain
387 negatively charged membranes (Van Wezel, 1967). Various materials could be used as
388 microcarriers, including dextran, cellulose, gelatin, and plastic (Stanbury et al., 2013). These
389 microcarriers might be solid or porous, and the materials could be selected according to culture
390 intention and cell type (Tavassoli et al., 2018). Food compliance should be considered in
391 situations where microcarriers are used in the production of edible meat. Although several
392 researchers have focused on the development of microcarriers suitable for human stem cells,
393 microcarriers for myoblast expansion or cultured meat production are yet to be developed
394 (Bodiou et al., 2020). The separation process of the cultured cells from the microcarriers is the

395 final step, in which cultured cells are used for subsequent applications (Nienow et al., 2014).
396 Microcarriers could be separated from the cultured cells by using enzymes or mechanical forces,
397 known to be a challenging procedure (Verbruggen et al., 2018). Microcarriers made of thermo-
398 responsive materials could temperature-dependently change their surface properties and
399 dislodge the attached cells, which could be subsequently filtered (Bodiou et al., 2020).
400 Biodegradable microcarriers are also being widely used as they do not require fastidious
401 harvesting procedures (Lam et al., 2017). Various edible polymers and hydrogels might be used
402 as bases for edible microcarriers (Ali and Ahmed, 2018). Edible microcarriers might not need
403 a cell dissociation step as the whole structure is safe for ingestion (Bodiou et al., 2020).
404 Myogenic satellite cells could be cultured in suspension with biodegradable or edible
405 microcarriers (Moritz et al., 2015). Recently, satellite cells have been grown and differentiated
406 in suspension culture systems using biodegradable scaffolds for the development of cultured
407 meat. This process requires cells to be anchored to scaffold surfaces, which could be provided
408 by tissue engineering constructs (Post, 2012).

409

410 **3.4 Scaffolding**

411 Obtaining tissue structure from muscle cell would be efficient way for creating cultured meat.
412 However, normally growing cells in a dish to get tissue-like structure is very challenging. For
413 cells to form an appropriate structure, scaffolds are utilized. Scaffolds molded into desirable
414 shape may provide physical support for muscle cell anchorage (Ben-Arye and Levenberg,
415 2019). Cells are highly niche dependent, and scaffolds aim to provide cells propriate niche-
416 resembling environment for growth (Zeltinger et al., 2001). Hydrogel is often used as scaffold
417 base material to mimic cell niche. Hydrogel engineered into porous structure mimics ECM as

418 it provides cells with permeable anchorage fit for water, gas, and nutrient exchange. Such 3D
419 scaffolds can be utilized by simply seeding the cells onto finished structure or mixing cells into
420 bioink and 3D printing cell encapsulated mixture to form cell-laden scaffold (Hwang et al.,
421 2010). Several types of base materials are used for tissue engineering. Collagen, fibrin, and
422 alginate are utilized as hydrogel, but to make gels more biologically like actual tissues, bioinks
423 using decellularized extracellular matrix(dECM) are introduced (Choi et al., 2016b). Bioinks
424 made with dECM contains more tissue-specific factors including growth factors, adhesive
425 proteins, compared to general hydrogels, and is believed to be more fit for tissue engineering
426 (Kim et al., 2020). Though no case of producing cultured meat by scaffolding cells have been
427 reported, but research show cell-laden 3D printed structures could be used for tissue
428 transplantation (Liu et al., 2019), and myoblasts are also capable of being 3D printed and
429 cultured (Choi et al., 2019). Decellularizing plant tissues for 3D cellulose scaffolds are also
430 viable. Plant tissues are abundant, easy to obtain and economically cheap. Culturing muscle
431 cells on decellularized plant scaffold stimulate growth, proliferation, and differentiation, while
432 providing myotube alignment due to natural plant cellulose patterns (Cheng et al., 2020).

433

434 **4. Future perspectives**

435 The ultimate goal of cultured meat is to produce edible meat products without directly
436 involving animals, not to obtain and proliferate the meat taken from livestock. To do this,
437 pluripotent stem cells might offer the best option as they could differentiate into muscle, fat,
438 and other cell types that could enhance the real meat flavor. Among the two pluripotent stem
439 cell types, embryonic and induced pluripotent stem cells (ESCs and iPSCs, respectively), iPSCs
440 seem to be more suitable as they are easy to establish and offer the advantage of a non-embryo-

441 based alternative. To date, iPSCs from various livestock have been established, including cattle
442 (Han et al., 2011), pigs (Wu et al., 2009), and chicken (Choi et al., 2016a). Although human
443 and mouse iPSCs exhibit limitless self-renewal potential, livestock iPS cells lose stemness
444 during long-term culture in the present culture system (Choi et al., 2016). Therefore, the culture
445 medium should be improved for long-term livestock iPSC culture. Since muscle tissue is a
446 complex structure of multiple different cell types, reliable muscle, fat, myoglobin, etc.,
447 differentiation protocols should be established, as well as a technique for forming a three-
448 dimensional (3D) structure for multiple cell types (Figure 2). Using the tissue engineering
449 technology or bioprinting system, muscle cells and various supportive cell types could be
450 cultured on the same 3D scaffold to form complex tissues that mimic *in vivo* skeletal muscle
451 structure (Krieger et al., 2018). Recently, a 3D engineered scaffold was used for bovine satellite
452 cells, which were proliferated on the 3D scaffold by submerging them into a myogenic growth
453 medium. Bovine smooth muscle cells and endothelial cells are differentiated on the scaffold to
454 form cell-based meat products, which are reported to be suitable for consumption as food
455 products (Ben-Arye et al., 2020).

456

457 **Acknowledgments**

458 This work was supported by the National Research Foundation of Korea (NRF) grant
459 funded by the Korea government (MSIT) of the Republic of Korea (grant nos.
460 2016M3A9B6946835 and 2020R1A2C3007562).

461

462 **References**

463 Adal MN, Cheng SB. 1980. Capsules of duck muscle spindles. Cell Tissue Res 211:465-474.

464 Alexander P, Brown C, Arneth A, Dias C, Finnigan J, Moran D, Rounsevell MD. 2017. Could
465 consumption of insects, cultured meat or imitation meat reduce global agricultural land
466 use? *Global Food Security* 15:22-32.

467 Alexander R. 2011. In vitro meat: A vehicle for the ethical rescaling of the factory farming industry
468 and in vivo testing or an intractable enterprise? *Intersect: The Stanford Journal of Science,
469 Technology, and Society* 4:42-47.

470 Ali A, Ahmed S. 2018. Recent advances in edible polymer based hydrogels as a sustainable
471 alternative to conventional polymers. *Journal of agricultural food chemistry* 66:6940-6967.

472 Allen RE, Merkel RA, Young RB. 1979. Cellular aspects of muscle growth: Myogenic cell
473 proliferation. *J Anim Sci* 49:115-127.

474 Arshad MS, Javed M, Sohaib M, Saeed F, Imran A, Amjad ZJCF, Agriculture. 2017. Tissue
475 engineering approaches to develop cultured meat from cells: A mini review. 3:1320814.

476 Bánáti D. 2011. Consumer response to food scandals and scares. *Trends in Food Science &
477 Technology* 22:56-60.

478 BBC news. 2013. *World's first lab-grown burger is eaten in london* [Online]. BBC. Available:
479 <https://www.bbc.com/news/science-environment-23576143>.

480 Ben-Arye T, Levenberg S. 2019. Tissue engineering for clean meat production. *Frontiers in
481 Sustainable Food Systems* 3:46.

482 Ben-Arye T, Shandalov Y, Ben-Shaul S, Landau S, Zagury Y, Ianovici I, Lavon N, Levenberg SJNF.
483 2020. Textured soy protein scaffolds enable the generation of three-dimensional bovine
484 skeletal muscle tissue for cell-based meat. 1:210-220.

485 Bhat Z, Bhat H, Pathak V. 2014. Prospects for in vitro cultured meat—a future harvest. In *Principles
486 of tissue engineering*. Elsevier.

487 Bhat ZF, Fayaz H. 2011. Prospectus of cultured meat—advancing meat alternatives. *Journal of
488 Food Science and Technology* 48:125-140.

489 Bhat ZF, Hina B. 2011. Animal-free meat biofabrication. *American Journal of Food Technology*
490 6:441-459.

491 Bhat ZF, Kumar S, Fayaz H. 2015. In vitro meat production: Challenges and benefits over
492 conventional meat production. *Journal of Integrative Agriculture* 14:241-248.

493 Bhat ZF, Morton JD, Mason SL, Bekhit AEDA, Bhat HFJCRIFS, Safety F. 2019. Technological,
494 regulatory, and ethical aspects of in vitro meat: A future slaughter-free harvest. 18:1192-
495 1208.

496 Bischoff R. 1974. Enzymatic liberation of myogenic cells from adult rat muscle. *Anat Rec* 180:645-
497 661.

498 Bischoff R. 1975. Regeneration of single skeletal muscle fibers in vitro. *Anat Rec* 182:215-235.

499 Bodiou V, Moutsatsou P, Post MJ. 2020. Microcarriers for upscaling cultured meat production.
500 *Frontiers in nutrition* 7:10.

501 Bonny SP, Gardner GE, Pethick DW, Hocquette J-F. 2015. What is artificial meat and what does it
502 mean for the future of the meat industry? *Journal of Integrative Agriculture* 14:255-263.

503 Brack AS, Conboy IM, Conboy MJ, Shen J, Rando TA. 2008. A temporal switch from notch to wnt
504 signaling in muscle stem cells is necessary for normal adult myogenesis. *Cell Stem Cell*
505 2:50-59.

506 Braun T, Gautel M. 2011. Transcriptional mechanisms regulating skeletal muscle differentiation,
507 growth and homeostasis. *Nat Rev Mol Cell Biol* 12:349-361.

508 Brodwin E. 2018a. *A new lab-grown meat startup says it's overcome a key barrier to making meat*
509 *without slaughter* [Online]. Business Insider. Available:
510 [https://www.businessinsider.com/lab-grown-meat-startup-solving-barrier-meat-without-](https://www.businessinsider.com/lab-grown-meat-startup-solving-barrier-meat-without-slaughter-meatable-2018-9)
511 [slaughter-meatable-2018-9](https://www.businessinsider.com/lab-grown-meat-startup-solving-barrier-meat-without-slaughter-meatable-2018-9).

512 Brodwin E. 2018b. *The startup behind the first lab-grown pork links let us see how their sausage*
513 *gets made – and said it slashed the cost from \$2,500 to \$216 in a month* [Online].
514 Business Insider. Available: [https://www.businessinsider.com/taste-test-lab-grown-meat-](https://www.businessinsider.com/taste-test-lab-grown-meat-sausage-cost-2018-11)
515 [sausage-cost-2018-11](https://www.businessinsider.com/taste-test-lab-grown-meat-sausage-cost-2018-11).

516 Bryant C, Dillard C. 2019. The impact of framing on acceptance of cultured meat. *Frontiers in*
517 *Nutrition* 6:103.

518 Bryant C, Szejda K, Parekh N, Desphande V, Tse B. 2019. A survey of consumer perceptions of
519 plant-based and clean meat in the USA, india, and china. *Frontiers in Sustainable Food*
520 *Systems* 3:11.

521 Bryant CJ. 2020. Culture, meat, and cultured meat. *Journal of Animal Science* 98:skaa172.

522 Chen G, Cheng X, Shi G, Zou C, Chen L, Li J, Li M, Fang C, Li C. 2019. Transcriptome analysis
523 reveals the effect of long intergenic noncoding rnas on pig muscle growth and fat
524 deposition. *Biomed Res Int* 2019:2951427.

525 Cheng Y-W, Shiwarski DJ, Ball RL, Whitehead KA, Feinberg AWJaBS, Engineering. 2020.
526 Engineering aligned skeletal muscle tissue using decellularized plant-derived scaffolds.
527 *6:3046-3054*.

528 Choi HW, Kim JS, Choi S, Ju Hong Y, Byun SJ, Seo HG, Do JTJSC, Development. 2016a.
529 Mitochondrial remodeling in chicken induced pluripotent stem-like cells. *25:472-476*.

530 Choi YJ, Jun YJ, Kim DY, Yi HG, Chae SH, Kang J, Lee J, Gao G, Kong JS, Jang J, Chung WK, Rhie
531 JW, Cho DW. 2019. A 3d cell printed muscle construct with tissue-derived bioink for the
532 treatment of volumetric muscle loss. *Biomaterials* 206:160-169.

533 Choi YJ, Kim TG, Jeong J, Yi HG, Park JW, Hwang W, Cho DW. 2016b. 3d cell printing of functional
534 skeletal muscle constructs using skeletal muscle-derived bioink. *Adv Healthc Mater*
535 *5:2636-2645*.

536 Choudhury D, Tseng TW, Swartz E. 2020. The business of cultured meat. *Trends in Biotechnology*
537 *38:573-577*.

538 Chriki S, Hocquette J-F. 2020. The myth of cultured meat: A review. *Frontiers in nutrition* 7:7.

539 Cole M, Morgan K. 2013. Engineering freedom? A critique of biotechnological routes to animal
540 liberation. *Configurations* 21:201-229.

541 Crew B. 2015. Cost of lab-grown burger patty drops from \$325,000 to \$11.36. *Science Alert* 2:-.

542 Datar I, Betti M. 2010. Possibilities for an in vitro meat production system. *Innovative Food
543 Science & Emerging Technologies* 11:13-22.

544 Dayton WR, White ME. 2008. Cellular and molecular regulation of muscle growth and
545 development in meat animals. *J Anim Sci* 86:E217-225.

546 Dilworth T, Mcgregor AJJOA, Ethics E. 2015. Moral steaks? Ethical discourses of in vitro meat in
547 academia and australia. 28:85-107.

548 Ding S, Wang F, Liu Y, Li S, Zhou G, Hu P. 2017. Characterization and isolation of highly purified
549 porcine satellite cells. *Cell Death Discov* 3:17003.

550 Dodson M, Mcfarland D, Martin E, Brannon M. 1986. Isolation of satellite cells from ovine skeletal
551 muscles. *Journal of tissue culture methods* 10:233-237.

552 Dodson MV, Allen RE, Du M, Bergen WG, Velleman SG, Poulos SP, Fernyhough-Culver M, Wheeler
553 MB, Duckett SK, Young MR, Voy BH, Jiang Z, Hausman GJ. 2015. Invited review: Evolution
554 of meat animal growth research during the past 50 years: Adipose and muscle stem cells.
555 *J Anim Sci* 93:457-481.

556 Dodson MV, Martin EL, Brannon MA, Mathison BA, Mcfarland DC. 1987. Optimization of bovine
557 satellite cell-derived myotube formation in vitro. *Tissue Cell* 19:159-166.

558 Dorovskikh A. 2015. Killing for a living: Psychological and physiological effects of alienation of
559 food production on slaughterhouse workers. Undergraduate Honors Theses.

560 Doumit ME, Merkel RA. 1992. Conditions for isolation and culture of porcine myogenic satellite
561 cells. *Tissue Cell* 24:253-262.

562 Du M, Tong J, Zhao J, Underwood KR, Zhu M, Ford SP, Nathanielsz PW. 2010. Fetal programming
563 of skeletal muscle development in ruminant animals. *J Anim Sci* 88:E51-60.

564 Dumont NA, Wang YX, Rudnicki MA. 2015. Intrinsic and extrinsic mechanisms regulating satellite
565 cell function. *Development* 142:1572-1581.

566 Fao. 2012. Agriculture organization of the united nations. 2012. *FAO statistical yearbook*.

567 Feldman JL, Stockdale FE. 1991. Skeletal muscle satellite cell diversity: Satellite cells form fibers of
568 different types in cell culture. *Dev Biol* 143:320-334.

569 Gey GO. 1933. An improved technic for massive tissue culture. *The American Journal of Cancer*
570 17:752-756.

571 Godfray H, Springmann M, Sexton A, Lynch J, Hepburn C, Jebb S. Meat: The future series
572 alternative proteins. *World Economic Forum*. p^pp.

573 Goodwin JN, Shoulders CW. 2013. The future of meat: A qualitative analysis of cultured meat
574 media coverage. *Meat Science* 95:445-450.

575 Gourmey. *Gourmey savoir-faire* [Online]. Available: <https://gourmey.com/en/>.

576 Greene E, Raub R. 1992. Procedures for harvesting satellite cells from equine skeletal muscle.

577 *Journal of Equine Veterinary Science* 12:33-35.

578 Greenlee A, Dodson M, Yablonka-Reuveni Z, Kersten C, Cloud J. 1995. In vitro differentiation of

579 myoblasts from skeletal muscle of rainbow trout. *Journal of fish biology* 46:731-747.

580 Grinnell F. 1978. Cellular adhesiveness and extracellular substrata. In *International review of*

581 *cytology*. Elsevier.

582 Han B, Tong J, Zhu MJ, Ma C, Du M. 2008. Insulin-like growth factor-1 (igf-1) and leucine activate

583 pig myogenic satellite cells through mammalian target of rapamycin (mtor) pathway. *Mol*

584 *Reprod Dev* 75:810-817.

585 Han X, Han J, Ding F, Cao S, Lim SS, Dai Y, Zhang R, Zhang Y, Lim B, Li N. 2011. Generation of

586 induced pluripotent stem cells from bovine embryonic fibroblast cells. *Cell Res* 21:1509-

587 1512.

588 Hoek AC, Luning PA, Stafleu A, De Graaf C. 2004. Food-related lifestyle and health attitudes of

589 dutch vegetarians, non-vegetarian consumers of meat substitutes, and meat consumers.

590 *Appetite* 42:265-272.

591 Hoekstra AY, Chapagain AK. 2006. Water footprints of nations: Water use by people as a function

592 of their consumption pattern. In *Integrated assessment of water resources and global*

593 *change*. Springer.

594 Hopkins PD, Dacey A. 2008. Vegetarian meat: Could technology save animals and satisfy meat

595 eaters? *Journal of Agricultural and Environmental Ethics* 21:579-596.

596 Hwang CM, Sant S, Masaeli M, Kachouie NN, Zamanian B, Lee SH, Khademhosseini A. 2010.

597 Fabrication of three-dimensional porous cell-laden hydrogel for tissue engineering.

598 *Biofabrication* 2:035003.

599 Just. 2018. Cultured meat: A vision of the future [Online]. Available:

600 <https://www.youtube.com/watch?v=f8li3DB6ejE>.

601 Kadim IT, Mahgoub O, Baqir S, Faye B, Purchas R. 2015. Cultured meat from muscle stem cells: A

602 review of challenges and prospects. *Journal of Integrative Agriculture* 14:222-233.

603 Karunaratne JF, Ashton CJ, Stickland NC. 2005. Fetal programming of fat and collagen in porcine

604 skeletal muscles. *J Anat* 207:763-768.

605 Kateman B. 2020. *Will cultured meat soon be a common sight in supermarkets across the globe?*

606 [Online]. Forbes. Available: [https://www.forbes.com/sites/briankateman/2020/02/17/will-](https://www.forbes.com/sites/briankateman/2020/02/17/will-cultured-meat-soon-be-a-common-sight-in-supermarkets-across-the-globe/#307f4af7c66a)

607 [cultured-meat-soon-be-a-common-sight-in-supermarkets-across-the-](https://www.forbes.com/sites/briankateman/2020/02/17/will-cultured-meat-soon-be-a-common-sight-in-supermarkets-across-the-globe/#307f4af7c66a)

608 [globe/#307f4af7c66a](https://www.forbes.com/sites/briankateman/2020/02/17/will-cultured-meat-soon-be-a-common-sight-in-supermarkets-across-the-globe/#307f4af7c66a).

609 Kim W, Lee H, Lee J, Atala A, Yoo JJ, Lee SJ, Kim GH. 2020. Efficient myotube formation in 3d

610 bioprinted tissue construct by biochemical and topographical cues. *Biomaterials*

611 230:119632.

612 Krieger J, Park BW, Lambert CR, Malcuit C. 2018. 3d skeletal muscle fascicle engineering is
613 improved with tgfbeta1 treatment of myogenic cells and their co-culture with
614 myofibroblasts. *PeerJ* 6:e4939.

615 Kunitake R, Suzuki A, Ichihashi H, Matsuda S, Hirai O, Morimoto K. 1997. Fully-automated roller
616 bottle handling system for large scale culture of mammalian cells. *J Biotechnol* 52:289-
617 294.

618 Laestadius LI. 2015. Public perceptions of the ethics of in-vitro meat: Determining an appropriate
619 course of action. *Journal of Agricultural and Environmental Ethics* 28:991-1009.

620 Laestadius LI, Caldwell MA. 2015. Is the future of meat palatable? Perceptions of in vitro meat as
621 evidenced by online news comments. *Public Health Nutrition* 18:2457-2467.

622 Lam AT-L, Li J, Toh JP-W, Sim EJ-H, Chen AK-L, Chan JK-Y, Choolani M, Reuveny S, Birch WR, Oh
623 SK-W. 2017. Biodegradable poly-ε-caprolactone microcarriers for efficient production of
624 human mesenchymal stromal cells and secreted cytokines in batch and fed-batch
625 bioreactors. *Cytotherapy* 19:419-432.

626 Laumonier T, Menetrey J. 2016. Muscle injuries and strategies for improving their repair. *J Exp*
627 *Orthop* 3:15.

628 Li C, Xiong T, Zhou M, Wan L, Xi S, Liu Q, Chen Y, Mao H, Liu S, Chen B. 2020. Characterization of
629 micrnas during embryonic skeletal muscle development in the shan ma duck. *Animals*
630 (Basel) 10.

631 Li H, Zhu C, Tao Z, Xu W, Song W, Hu Y, Zhu W, Song C. 2014. Myod and myf6 gene expression
632 patterns in skeletal muscle during embryonic and posthatch development in the domestic
633 duck (*anas platyrhynchos domestica*). *J Anim Breed Genet* 131:194-201.

634 Li L, Liu HH, Xu F, Si JM, Jia J, Wang JW. 2010. Myod expression profile and developmental
635 differences of leg and breast muscle in peking duck (*anas platyrhynchos domestica*)
636 during embryonic to neonatal stages. *Micron* 41:847-852.

637 Liu HH, Wang JW, Zhang RP, Chen X, Yu HY, Jin HB, Li L, Han CC, Xu F, Kang B, He H, Xu HY.
638 2012. In ovo feeding of igf-1 to ducks influences neonatal skeletal muscle hypertrophy
639 and muscle mass growth upon satellite cell activation. *J Cell Physiol* 227:1465-1475.

640 Liu J, Li L, Suo H, Yan M, Yin J, Fu JJM, Design. 2019. 3d printing of biomimetic multi-layered
641 gelma/nha scaffold for osteochondral defect repair. 171:107708.

642 Lucas A. 2019. *Lab-grown meat start-up raises \$14 million to build production plant*. [Online].
643 CNBC. Available: <https://www.cnbc.com/2019/10/10/future-meat-technologies-a-lab-grown-meat-start-up-raises-14-million-dollars.html>.

644

645 Macleod M, Gerber P, Mottet A, Tempio G, Falcucci A, Opio C, Vellinga T, Henderson B, Steinfeld
646 H. 2013. Greenhouse gas emissions from pig and chicken supply chains—a global life cycle
647 assessment. Food and Agriculture Organization of the United Nations.

648 Mancini MC, Antonioli F. 2020. To what extent are consumers' perception and acceptance of

649 alternative meat production systems affected by information? The case of cultured meat.
650 Animals 10:656.

651 Marga FS. 2012. Engineered comestible meat. 1601 S. Providence Road Columbia,MO 65211:
652 National Institute of Food and Agriculture.

653 Marga FS. 2016. Dried food products formed from cultured muscle cells. Google Patents.

654 Matsuda R, Spector DH, Strohman RC. 1983. Regenerating adult chicken skeletal muscle and
655 satellite cell cultures express embryonic patterns of myosin and tropomyosin isoforms.
656 Dev Biol 100:478-488.

657 Mau M, Oksbjerg N, Rehfeldt C. 2008. Establishment and conditions for growth and differentiation
658 of a myoblast cell line derived from the semimembranosus of newborn piglets. In
659 Vitro Cellular Developmental Biology-Animal 44:1-5.

660 Mccarthy JJ. 2008. Microrna-206: The skeletal muscle-specific myomir. Biochim Biophys Acta
661 1779:682-691.

662 Meats M. 2016a. *Meat culture* [Online]. Available:
663 <https://www.youtube.com/watch?v=DqaHWyimbQ8>.

664 Meats M. 2016b. The world's first cell-based meatball [Online]. Available:
665 <https://www.youtube.com/watch?v=Y027yLT2QY0>.

666 Meats M. 2017. *Histroic first: Cell-based poultry tasting* [Online]. Available:
667 <https://www.youtube.com/watch?v=b5ezRx23EMg>.

668 Melero-Martin J, Al-Rubeai M. 2007. In vitro expansion of chondrocytes. Topics in tissue
669 engineering 3:37.

670 Melvin RG. 1995. Non point sources of pollution on rangeland. Fact Sheet No. 3. Rangeland
671 Watershed Program; U.C. Cooperative Extension and USDA Natural Resources
672 Conservation Service; California Rangelands Research and Information Center Agronomy
673 and Range Science – UC Davis.

674 Merten OW. 2015. Advances in cell culture: anchorage dependence. Philosophical Transactions of
675 the Royal Society B: Biological Sciences 370: 1661.

676 Metzger K, Tuchscherer A, Palin MF, Ponsuksili S, Kalbe C. 2020. Establishment and validation of
677 cell pools using primary muscle cells derived from satellite cells of pig skeletal muscle. In
678 Vitro Cell Dev Biol Anim 56:193-199.

679 Moritz MS, Verbruggen SE, Post MJ. 2015. Alternatives for large-scale production of cultured beef:
680 A review. Journal of Integrative Agriculture 14:208-216.

681 Mouat MJ, Prince R. 2018. Cultured meat and cowless milk: On making markets for animal-free
682 food. Journal of Cultural Economy 11:315-329.

683 Musaro A, Mccullagh K, Paul A, Houghton L, Dobrowolny G, Molinaro M, Barton ER, Sweeney HL,
684 Rosenthal N. 2001. Localized igf-1 transgene expression sustains hypertrophy and
685 regeneration in senescent skeletal muscle. Nat Genet 27:195-200.

686 Nienow AW, Rafiq QA, Coopman K, Hewitt CJ. 2014. A potentially scalable method for the
687 harvesting of hmscs from microcarriers. *Biochemical Engineering Journal* 85:79-88.

688 Perruchot MH, Lefaucheur L, Barreau C, Casteilla L, Louveau I. 2013. Age-related changes in the
689 features of porcine adult stem cells isolated from adipose tissue and skeletal muscle. *Am J*
690 *Physiol Cell Physiol* 305:C728-738.

691 Petetin L. 2014. Frankenburger, risks and approval. *Eur. J. Risk Reg.* 5:168.

692 Post MJ. 2012. Cultured meat from stem cells: Challenges and prospects. *Meat Sci* 92:297-301.

693 Post MJ, Levenberg S, Kaplan DL, Genovese N, Fu J, Bryant CJ, Negowetti N, Verzijden K,
694 Moutsatsou P. 2020. Scientific, sustainability and regulatory challenges of cultured meat.
695 *Nature Food.* 1:403-415.

696 Purdy C. 2020. *I'll have the lab-grown duck pâté, please* [Online]. Newamerica. Available:
697 [https://www.newamerica.org/weekly/ill-have-the-lab-grown-duck-p%C3%A2t%C3%A9-
698 please/](https://www.newamerica.org/weekly/ill-have-the-lab-grown-duck-p%C3%A2t%C3%A9-please/).

699 Qiu H, Xu X, Fan B, Rothschild MF, Martin Y, Liu B. 2010. Investigation of *Idha* and *copb1* as
700 candidate genes for muscle development in the *myod1* region of pig chromosome 2. *Mol*
701 *Biol Rep* 37:629-636.

702 Rafiq QA, Coopman K, Hewitt CJ. 2013. Scale-up of human mesenchymal stem cell culture:
703 Current technologies and future challenges. *Current Opinion in Chemical Engineering* 2:8-
704 16.

705 Ridgeway AG, Skerjanc IS. 2001. *Pax3* is essential for skeletal myogenesis and the expression of
706 *six1* and *eya2*. *J Biol Chem* 276:19033-19039.

707 Rubin LL, And Feodor D. Price. 2019. *Satellite cells and compositions and methods for producing*
708 *the same*. United States patent application US16/297,548.

709 Sadler MJ. 2004. Meat alternatives—market developments and health benefits. *Trends in Food*
710 *Science & Technology* 15:250-260.

711 Savvides L. 2020. *Lab-grown meat: Taste-testing chicken of the future* [Online]. Cnet. Available:
712 [https://www.cnet.com/news/lab-grown-cultured-clean-meat-tasting-the-chicken-of-the-
713 future-just-inc/](https://www.cnet.com/news/lab-grown-cultured-clean-meat-tasting-the-chicken-of-the-future-just-inc/).

714 Schneider Z. 2012. In vitro meat: Space travel, cannibalism, and federal regulation. *Hous. L. Rev.*
715 50:991.

716 Shaffer E. 2020. Memphis meats sets sights on building a pilot plant [Online]. MEAT+POULTRY:
717 SOSLAND PUBLISHING COMPANY. Available: [https://www.meatpoultry.com/articles/22509-
718 memphis-meats-sets-sights-on-building-a-pilot-plant](https://www.meatpoultry.com/articles/22509-memphis-meats-sets-sights-on-building-a-pilot-plant).

719 Shi X, Garry DJ. 2006. Muscle stem cells in development, regeneration, and disease. *Genes Dev*
720 20:1692-1708.

721 Siegrist M, Hartmann C. 2020. Consumer acceptance of novel food technologies. *Nature Food*
722 1:343-350.

723 Siegrist M, Sütterlin B, Hartmann C. 2018. Perceived naturalness and evoked disgust influence
724 acceptance of cultured meat. *Meat science* 139:213-219.

725 Southey F. 2020. *'Ethical foie-gras': Cultured meat start-up taps duck egg cells to recreate french*
726 *delicacy* [Online]. Food Navigator. Available:
727 [https://www.foodnavigator.com/Article/2020/05/14/Ethical-foie-gras-Cultured-meat-start-](https://www.foodnavigator.com/Article/2020/05/14/Ethical-foie-gras-Cultured-meat-start-up-taps-duck-egg-cells-to-recreate-French-delicacy)
728 [up-taps-duck-egg-cells-to-recreate-French-delicacy.](https://www.foodnavigator.com/Article/2020/05/14/Ethical-foie-gras-Cultured-meat-start-up-taps-duck-egg-cells-to-recreate-French-delicacy)

729 Stanbury PF, Whitaker A, Hall SJ. 2013. Principles of fermentation technology. Elsevier.

730 Steinfeld H, Gerber P, Wassenaar T, Castel V, Rosales M, Rosales M, De Haan C. 2006. Livestock's
731 long shadow: Environmental issues and options. Food & Agriculture Org.

732 Stephens N, Di Silvio L, Dunsford I, Ellis M, Glencross A, Sexton A. 2018. Bringing cultured meat to
733 market: Technical, socio-political, and regulatory challenges in cellular agriculture. *Trends*
734 *in Food Science & Technology* 78:155-166.

735 Stickland NC. 1978. A quantitative study of muscle development in the bovine foetus (*bos*
736 *indicus*). *Anat Histol Embryol* 7:193-205.

737 Tavassoli H, Alhosseini SN, Tay A, Chan PPY, Weng Oh SK, Warkiani ME. 2018. Large-scale
738 production of stem cells utilizing microcarriers: A biomaterials engineering perspective
739 from academic research to commercialized products. *Biomaterials* 181:333-346.

740 Tomiyama AJ, Kawecki NS, Rosenfeld DL, Jay JA, Rajagopal D, Rowat AC. 2020. Bridging the gap
741 between the science of cultured meat and public perceptions. *Trends in Food Science and*
742 *Technology*.

743 Tuomisto HL, Teixeira De Mattos MJ. 2011. Environmental impacts of cultured meat production.
744 *Environmental science & technology* 45:6117-6123.

745 Un. 2019. World population prospects 2019. United nations. Department of economic and social
746 affairs.

747 Van Der Weele C, Tramper J. 2014. Cultured meat: Every village its own factory? *Trends in*
748 *Biotechnology* 32:294-296.

749 Van Wezel A. 1967. Growth of cell-strains and primary cells on micro-carriers in homogeneous
750 culture. *Nature* 216:64-65.

751 Verbeke W, Marcu A, Rutsaert P, Gaspar R, Seibt B, Fletcher D, Barnett J. 2015. 'Would you eat
752 cultured meat?': Consumers' reactions and attitude formation in Belgium, Portugal and the
753 United Kingdom. *Meat science* 102:49-58.

754 Verbruggen S, Luining D, Van Essen A, Post MJ. 2018. Bovine myoblast cell production in a
755 microcarriers-based system. *Cytotechnology* 70:503-512.

756 Welin S, Van Der Weele C. 2012. Cultured meat: Will it separate us from nature? In *Climate*
757 *change and sustainable development*. Springer.

758 Wesson LG, Jr., Cohn WE, Brues AM. 1949. The effect of temperature on potassium equilibria in
759 chick embryo muscle. *J Gen Physiol* 32:511-523.

760 Wojtczak J. 1979. Contractures and increase in internal longitudinal resistance of cow ventricular
761 muscle induced by hypoxia. *Circ Res* 44:88-95.

762 Wu N, Gu T, Lu L, Cao Z, Song Q, Wang Z, Zhang Y, Chang G, Xu Q, Chen G. 2019. Roles of
763 miRNA-1 and miRNA-133 in the proliferation and differentiation of myoblasts in duck
764 skeletal muscle. *J Cell Physiol* 234:3490-3499.

765 Wu Z, Chen J, Ren J, Bao L, Liao J, Cui C, Rao L, Li H, Gu Y, Dai H, Zhu H, Teng X, Cheng L, Xiao L.
766 2009. Generation of pig induced pluripotent stem cells with a drug-inducible system. *J*
767 *Mol Cell Biol* 1:46-54.

768 Yablonka-Reuveni Z, Quinn LS, Nameroff M. 1987. Isolation and clonal analysis of satellite cells
769 from chicken pectoralis muscle. *Dev Biol* 119:252-259.


770 Yamashita T, Sohal GS. 1986. Development of smooth and skeletal muscle cells in the iris of the
771 domestic duck, chick and quail. *Cell Tissue Res* 244:121-131.

772 Zeltinger J, Sherwood JK, Graham DA, Mueller R, Griffith LG. 2001. Effect of pore size and void
773 fraction on cellular adhesion, proliferation, and matrix deposition. *Tissue Eng* 7:557-572.

774 Zhu MJ, Ford SP, Nathanielsz PW, Du M. 2004. Effect of maternal nutrient restriction in sheep on
775 the development of fetal skeletal muscle. *Biol Reprod* 71:1968-1973.

776

777 **Figure legends**




778

779 **Figure 1. Muscle satellite cell myogenic differentiation pathway with expressing markers.** Muscle satellite
780 cells potential for differentiating into muscle fibre. Quiescent satellite cells express paired box 7 (Pax7) while
781 myogenic factor 5 (Myf5) is downregulated. In the process of developing into myoblast and muscle fibre,
782 satellite cells are proliferated as well as differentiated. Myogenic determination (MyoD) and myogenin (Myog)
783 marks the production of more complex filaments while differentiation undergoes.

784

ACCEPTED



785

786

Figure 2. Technical approach for producing cultured meat. Adult stem cells and induced pluripotent stem

787

(iPS) cells could both be considered cultured meat sources. Myogenic satellite cells and adipose stem cells are

788

proliferated through *in vitro* culturing and manufactured to resemble meat structure. iPS cells could differentiate

789

into several different cell types comprising muscle tissue that could be used, along with multiple other cell

790

types, to manufacture three-dimensional (3D) structures using tissue engineering or bioprinting technologies.

791

AC

Table 1. Comparison of traditional and cultured meat

| Attributes | Traditional meat | Cultured meat | References |
|--------------------------|-------------------------|----------------------|---|
| Production System | | | |
| Production method | Animal farming | Cell cultivation | (Bhat et al., 2019) |
| Land requirement | High | Low | (Alexander et al., 2017) |
| Location of production | Mostly rural | Rural and urban | (Bhat et al., 2019) |
| Production cost | High | (So far) Very High | (Van der Weele and Tramper, 2014) |
| Production time | Long | Short | (Bhat and Fayaz, 2011) |
| Production yield | Low | High | (Alexander et al., 2017) |
| Greenhouse gas emission | Very high | Low | (Bhat and Fayaz, 2011) |
| Energy requirement | High | High | (Tuomisto and Teixeira de Mattos, 2011) |
| Water and soil pollution | High | Low | (Welin and Van der Weele, 2012) |
| Sustainability | Low | High | (Siegrist and Hartmann, 2020) |
| Characteristics | | | |
| Manipulating composition | Impossible | Possible | (Bhat and Fayaz, 2011) |
| Human health | Low | High | (Joshi et al., 2020) |
| Food safety | Low | High | (Joshi et al., 2020) |
| Animal welfare | Low | High | (Mouat and Prince, 2018) |
| Ethical | Low | High | (Mancini and Antonioli, 2020) |

advantage

Consumer
acceptance

High

Low

(Siegrist et al., 2018)

793

ACCEPTED

Table 2. Diverse cultured meat products currently being developed

| Species | Company | Product | Manufacture year | Country |
|---------------------|--------------------------|----------------------|-------------------------|----------------|
| Chicken meat | JUST | chicken nugget | 2019 | USA |
| | Memphis Meats | chicken tender | 2017 | USA |
| | Peace of Meat | chicken nugget | 2020 | Belgium |
| | Future Meat Technologies | shawarma | 2019 | Israel |
| Duck meat | JUST | duck pâté & chorizo | 2020 | USA |
| | Memphis Meats | nugget | 2019 | USA |
| | Gourmey | foie gras | 2020 | France |
| Beef | Mosa Meat | burger | 2013 | Netherlands |
| | Memphis Meats | meat ball | 2016 | USA |
| Pork | Higher Steaks | pork belly and bacon | 2020 | UK |
| | New Age Meats | pork sausage | 2019 | USA |

795

796

797