TITLE PAGE - Food Science of Animal Resources -Upload this completed form to website with submission

ARTICLE INFORMATION	Fill in information in each box below
Article Type	Research article,
Article Title	Hovenia monofloral honey can attenuate Enterococcus faecalis mediated biofilm formation and inflammation
Running Title (within 10 words)	Hovenia monofloral honey has anti-biofilm and anti-inflammatory activities
Author	Ri You1, Oh Yun Kwon1, Hyun Joo Woo, Seung Ho Lee1,*
Affiliation	1Department of Nano-Bioengineering, Incheon National University, 119 Academy-ro, Incheon 22012, Korea
Special remarks – if authors have additional information to inform the editorial office	These author(Ri You and Oh Yun Kwon) contributed equally in this work
ORCID (All authors must have ORCID) https://orcid.org	Ri You: 0000-0002-0695-8709 Oh Yun Kwon: 0000-0002-3783-3220 Hyun Joo Woo: 0000-0001-8129-6760 Seung Ho Lee: 0000-0002-9941-4195
Conflicts of interest List any present or potential conflict s of interest for all authors. (This field may be published.)	The authors declare no potential conflict of interest.
Acknowledgements State funding sources (grants, funding sources, equipment, and supplies). Include name and number of grant if available. (This field may be published.)	This research was supported by an Incheon National University research grant in 2019. Hovenia monofloral honys (HMHs) which produced by using a net house system were provided from National Institute of Forest Science in South Korea.
Author contributions (This field may be published.)	Conceptualization: Lee SH. Data curation: You R, Kwon OY, Woo HJ. Formal analysis: You R, Kwon OY, Woo HJ. Methodology: You R, Kwon OY, Woo HJ Software: You R, Kwon OY, Woo HJ. Validation: You R, Kwon OY, Woo HJ HJ Investigation: You R, Kwon OY, Woo HJ Writing - original draft: Lee SH. Writing - review & editing: You R, Kwon OY, Woo HJ and Lee SH. (This field must list all authors)
(This field may be published.)	and animal participants.

CORRESPONDING AUTHOR CONTACT INFORMATION

For the <u>corresponding</u> author	Fill in information in each box below	
(responsible for correspondence,		
proofreading, and reprints)		
First name, middle initial, last name	Seun Ho Lee	
Email address – this is where your proofs	lsh1602@gmail.com	
will be sent		
Secondary Email address	seungho@inu.ac.kr	
Postal address	22012	

Cell phone number	+82-10-8014-1614
Office phone number	+82-32-835-8269
Fax number	+82-32-8350798

Hovenia monofloral honey can attenuate *Enterococcus faecalis* mediated biofilm formation and inflammation

Abstract

we evaluated the anti-biofilm formation and anti-inflammatory activity of Hovenia monofloral honey (HMH) against Enterococcus faecalis. Co-culture of HMH with E. faecalis attenuated the biofilm formation of *E. faecalis* on a polystyrene surface. In addition, HMH effectively eradicated the established E. faecalis biofilm. HMH significantly attenuated E. faecalis growth but did not affect the production of extracellular polymeric substances (EPSs) on E. faecalis, indicating that reduction of E. faecalis biofilm is a result of HMH-mediated killing of E. faecalis. Furthermore, we found that HMH can effectively attenuate E. faecalis-induced expression of a proinflammatory interleukin-8 (IL-8) in HT-29 cells. Interestingly, treatment of HMH significantly attenuated the E. faecalis-mediated expression of Toll-like receptor-2 (TLR-2) and its adaptor molecules, myeloid differentiation primary response 88 (MyD88), in HT-29 cells. In addition, E. faecalis-induced mitogen-activated protein kinases (MAPKs) phosphorylation was significantly attenuated by HMH administration. Furthermore, HMHmediated anti-inflammatory efficacy (0.2 mg/mL of HMHs) had an equal extent of inhibitory efficacy as 5 µM of MyD88 inhibitor to attenuate E. faecalis-mediated IL-8 expression in HT-29 cells. These results suggest that HMH could effectively inhibit E. faecalis-mediated gastrointestinal inflammation through regulating the TLR-2/MyD88/MAPKs signaling pathways. Collectively, our data suggest that HMH could be developed as a potential natural agent to control E. faecalis-mediated biofilm formation and inflammation.

Key words Hovenia monofloral honey, *Enterococcus faecalis*, biofilm, inflammation, Toll like receptor-2, MAPKs

Introduction

Enterococcus faecalis is a gram-positive bacteria that can be isolated from normal intestinal flora and optionally used as probiotics with *E. faecium* (Baccouri et al., 2019; Zommiti et al., 2018). However, *E. faecalis* is also known as an opportunistic pathogen that can induce peritonitis, urinary tract infections, and bacteremia (Lee et al., 2017). *E. faecalis* is reportedly able to make biofilms on food-processing plants, which could be a major microbial contaminant of foods (Liu et al., 2013; Pesavento et al., 2014). Therefore, *E. faecalis* is considered a foodborne pathogen, which can induce various infectious diseases accompanied by inflammatory responses (Giraffa, 2002).

A bacterial biofilm is a complex structure embedded in a matrix of exopolysaccharides, membrane vesicles, and bacterial DNA. This polymeric structure protects the bacterial cells from environmental stresses such as nutritional limitation, pH alteration, and osmolarity (Fux et al., 2005). In addition, this polymeric structure prevents the action of antibiotics, which results in the production of antibiotic-resistant bacterial populations. This causes severe problems for curing inflammatory responses in chronic wounds (Engemann et al., 2003; Hoyle and Costerton, 1991; Ngo et al., 2012; Wolcott et al., 2008). Therefore, demand for new agents that can more effectively control biofilm formation and biofilm-derived inflammatory responses continues to grow.

Honey is an edible natural agent that is produced by bees from the nectar of flowering plants. Honey mainly consists of primarily sugars, including oligosaccharides, and also contains minerals, vitamins, and plant-derived phenolic compounds (Bogdanov et al., 2008; da Silva et al., 2016; Qiu et al., 1999). Honey has been recognized as a medicinal food because of its antiinflammatory, anti-oxidative, and anti-bacterial activities (Ranneh et al., 2021; Sherlock et al., 2010; Stagos et al., 2018). Since the components of honey vary with weather condition, geographic location, and floral source, monofloral honeys, which are produced at different locations, show a range of biological activities (Allen et al., 1991; N et al., 2013).

The Hovenia (*Hovenia dulcis*) tree is a potential candidate for a honey plant in South Korea because Hovenia flowers produce higher amounts of nectar than Acacia flowers, and one of their characteristics is that they prefer to grow in a sunny position. Therefore, in South Korea, areas where Hovenia trees are being cultivated are gradually increasing. Physiochemical analysis, antioxidant, and anti-bacterial activities of Hovenia monofloral honeys (HMHs) were estimated in our previous study (Park et al., 2020). However, although HMH showed strong anti-bacterial activity against gram-positive (*Listeria monocytogenes* and *Staphylococcus aureus*) and gram-negative (*Salmonella* Typhimurium and *E. coli* O157:H7) strains, the effects of HMHs on *Enterococcus faecalis*-mediated biofilm formation and inflammation has not yet been elucidated. Therefore, in this study, the anti-biofilm and anti-inflammatory effects of HMH were evaluated and their underlying molecular mechanisms were also investigated.

Materials and Methods

Materials

HMHs were produced two times on different days (HMH-1, HMH-2) using a net house system. The purity of the HMHs was estimated by pollen analysis according to Park et al.'s method (Park et al., 2020), and the HMHs were determined to have more than 95% Hovenia

pollen. The HMHs were stored at 4°C in a refrigerator until analysis.

Rabbit anti-TLR-2 antibody was obtained from Invitrogen Co. (Invitrogen, Waltham, MA, USA). Rabbit anti-Myeloid differentiation primary response 88 (MyD88) (4283S), Rabbit anti-phospho-ERK (9102S), rabbit anti-ERK (9101S), rabbit anti-phospho-p38 (9212S), rabbit anti-p38 (9211S), rabbit anti-phospho-JNK (9251S), rabbit anti-JNK (9252S), and rabbit anti-GAPDH (5174S) were purchased from Cell Signaling Co. (Cell Signaling, Danver, MA, USA).

Biofilm formation and biofilm elimination assay

Enterococcus faecalis (KCTC 3511), which was obtained from the Korean Collection for Type Culture (KCTC), was used in this study. *E. faecalis* was incubated in a brain heart infusion (BHI; BD Biosciences, Franklin Lakes, NJ, USA) medium at 37°C. To assess the effects of HMH on the biofilm formation of *E. faecalis*, various concentrations of HMH (0~32%, w/v) were co-cultured with *E. faecalis* (1×10^7 /well) in a black polystyrene 96-well microtiter plate with a clear bottom (Greiner Bio-one, Kremsmunster, Austria) at 37°C in an incubator for 24 h. After washing with phosphate buffered saline (PBS) three times, the HMHs were stained with SYTO9 (25 µM) solution at room temperature for 30 min. After washing with PBS once, biofilm formation was examined visually using a confocal laser scanning microscope (Nickon, Tokyo, Japan).

To estimate the effects of HMH on the established *E. faecalis* biofilms, *E. faecalis* (1×10^{7} /well) was first added to a 96-well microtiter plate and incubated at 37°C in an incubator for 24 h to form biofilms. After washing with PBS three times, HMH (0~32%) was added and further incubated at 37°C in an incubator for 24 h. The biofilm was stained with a SYTO9 solution, as described as above, and examined visually using a confocal laser scanning microscope (Nickon, Tokyo, Japan). Biofilm mass were quantified by using the ImageJ

program.

Enterococcus faecalis viability assay

E. faecalis (2.5×10^4) with and without HMH was cultured in BHI broth at 37°C in an incubator for 2, 4, and 6 h. The growth of *E. faecalis* and cell viability were calculated by estimating the optical density at 595 nm and 450 nm respectively by using a microplate reader (iMark Microplate Reader, Bio-Rad laboratories, Inc., Contra Costa County, CA, USA). The effects of HMH on *E. faecalis* growth was further estimated using a WST-1(2-[4-Iodophenyl]-3-[4-nitrophenyl]-5-[2, 4-disulfophenyl]-2H-tetrazolium) cell viability assay kit according to the manufacturer's protocol. Briefly, *E. faecalis* (2.5×10^4) was cultured in 96-well microtiter plate with and without HMH in a 37°C incubator for 24 h. Then WST-1 solution was added, and the samples were incubated at room temperature for an additional 30 min.

Measuring extracellular polymeric substances

Estimating the amount of extracellular polymeric substances (EPSs) was conducted according to Yoon and Kang's method (Yoon and Kang, 2020) with minor modifications. Briefly, HMH (0, 16, and 32%) was co-cultured with *E. faecalis* for 16 h and then the bacterial pellet was diluted in a PBS to reach a turbidity of 1.0 at OD_{600} . Then, the *E. faecalis* suspension (1 mL) was mixed with 1 mL of Congo red solution (80 µg/mL) and further incubated at 37°C in an incubator for 2 h. After centrifuging at 4,000 relative centrifugal force (RCF) for 30 min, the optical density of the supernatant was estimated at OD_{492} using a UV/VIS spectrophotometer (Optizen POP, Mecasys, Daejeon, Korea) and the amount of Congo red was calculated by comparing the optical density of the standard curve of Congo red.

Cell culture and cytotoxicity assay

The HT-29 colorectal adenocarcinoma cell line, which was obtained from the American Type Culture Collection (ATCC; Manassa, VA, USA) and cultured in Dulbecco's Modified Eagle's Medium (DMEM; Welgen, Seoul, Korea), was mixed with 10% fetal bovine serum (Corning Inc., Corning, NY, USA), 100 unit/mL of penicillin, and 100 μ g/mL of streptomycin. To estimate the cytotoxicity of the HMH, HT-29 cells (1 × 10⁴) were cultured on a 96-well cell culture plate at 37 °C in a CO₂ incubator for 24 h. Then, the culture media was changed to serum-free DMEM containing HMH (0~2 mg/mL) and further incubated at 37 °C in a CO₂ incubator for 24 h. Subsequently, 10 μ L of WST-1 solution was added to each well and incubated for an additional 30 min in a CO₂ incubator. The number of viable cells was calculated by estimating the absorbance at 450 nm using a microplate reader (iMark Microplate Reader, Bio-Rad laboratories, Inc., Contra Costa County, CA, USA).

Western blotting

HT-29 cells (5 × 10⁵/wells) were seeded in six-well plates and cultured for 24 h. After incubation with serum-free DMEM for 16 h, *E. faecalis* (2 × 10⁶/well) was administered to HT-29 cells with and without HMH. After further incubation in a CO₂ incubator for 16 h, it was lysed with protein extraction buffer (Lee et al., 2010) on ice for 1 h. After centrifuging (12,000 × g) at 4°C for 15 min, the supernatants were separated and used for estimating the expression of each protein. Each protein sample was separated in a sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and the gels were separated and transferred to nitrocellulose membranes. After being incubated with a blocking buffer (5% nonfat milk in Tris-buffered saline with 1% Tween 20), each antibody was first reacted with the membranes (1:1,500) for 12 h at 4°C. The membranes were washed with a Tris-buffered saline with Tween 20 buffer (TBST), and reacted with a horseradish peroxidase conjugated rabbit secondary antibody (1:3,000) (Abcam, Cambridge, UK) at room temperature for 2 h. Each band was examined visually using an enhanced chemiluminescence (ECL) detection reagent (Bio-Rad, Hercules, CA, USA).

Quantitative real-time PCR (qRT-PCR)

HT-29 cells (5 \times 10⁵/wells) were seeded in six-well plates and cultured for 24 h. After incubation with serum-free DMEM for 16 h, E. faecalis (2×10^{6} /well), 10 µM of a control inhibitor (NBP2-29334, Novus, Centennial, CO, USA), and MyD88 inhibitor (NBP2-31226, Novus, Centennial, CO, USA) were administered to HT-29 cells with and without HMH. After further incubation in a CO₂ incubator for 16 h, the total RNA of the HT-29 cells, which were treated as above, was extracted using a TRIzol® Reagent (Invitrogen, Waltham, MA, USA). Complementary DNA (cDNA), which was synthesized with 1 µg of the total RNA and 10 pM of oligo dT primer, was mixed with SYBR® Green Realtime PCR Master Mix (Toyobo Co., Tokyo, Japan) and qRT-PCR was performed to compare the transcriptional expression of the IL-8 gene in a RT-PCR detection system (CFX ConnectTM, Bio-Rad Co., Hercules, CA, USA). A comparative ct method was used for evaluating the relative expression of the IL-8 gene and normalized to the expression of glyceraldehyde 3-phosphate dehydrogenase (GAPDH). The sequences of the IL-8 primers were 5'-TGTCCCATGCCACTCAGAGA-3' (forward) and 5'-5'-AGCAGGTGCTCCGGTTGTAT-3' (reverse); and those of the glyceraldehyde 3-phosphate dehydrogenase primers were 5'-ACGGGAAGCTCACTGGCA-3' (forward) and 5'-TCCAGGCGGCATGTCAGA -3' (reverse).

Statistical analysis

All results were expressed as mean \pm standard deviation (SD). Two-tailed, unpaired student's *t*-test and an analysis of variance (ANOVA) in Prism 5 software (Graph-Pad Software, Inc., San Diego, CA, USA) were used for statistical analysis. *p* < 0.05 considered to be statistically significant.

Results

Hovenia monofloral honey attenuated the formation of *E. faecalis*-mediated biofilms and stimulated the destruction of pre-formed *E. faecalis* biofilms

To investigate whether or not HMH can inhibit the formation of *E. faecalis* biofilms, HMHs (HMH-1 and HMH-2) were first incubated with *E. faecalis* for 24 h. Then, the *E. faecalis* biofilms were examined visually by SYTO9 staining (Invitrogen, Middlesex county, MA, USA) and quantified under a confocal laser scanning microscope. As shown in Fig. 1, the SYTO9 positive area, which is considered as the *E. faecalis* biofilm, was significantly (p < 0.05) decreased when 16% and 32% (w/v) of HMHs was co-incubated with *E. faecalis*. In addition, to examine whether HMH could destruct the *E. faecalis* biofilms, various concentrations (0~32%) of HMHs were administered to pre-formed *E. faecalis* and were examined visually by SYTO9 staining. As shown in Fig. 2, administration of 16% and 32% of HMHs to pre-formed *E. faecalis* biofilm significantly (p < 0.05) decreased the SYTO9 positive area when compared with that of the control (non-treated). These results suggest that HMH could attenuate the formation of *E. faecalis* biofilm and could effectively eradicate the pre-formed biofilm of *E. faecalis*.

Hovenia monofloral honey can inhibit the cell viability of E. faecalis

Since the HMH showed inhibitory activity on *E. faecalis* biofilm formation and also exhibited stimulatory properties on elimination of the *E. faecalis* biofilm (Fig. 2), we then examined the effects of HMH on the viability and EPSs production of *E. faecalis*. As shown in Fig. 3, administration of HMH on the planktonic growth of *E. faecalis* (which was estimated by measuring the optical density at 2, 4, and 6 h after inoculation) dose-dependently attenuated the growth of *E. faecalis*. The inhibitory activity of HMH was further examined using a WST-1 cell viability assay kit at 24 h after inoculation. About 60% and 80% of the *E. faecalis*, respectively. However, the production of EPSs, which is considered an important factor in the formation of *E. faecalis*-mediated biofilm, was not affected by HMH treatment. These data indicate that the HMHs' attenuation of the biofilm formation of *E. faecalis* was a result of the growth-inhibitory effects of the HMH but not that of EPS production suppression effect of HMH.

Hovenia monofloral honey inhibited the *E. faecalis*-mediated expression of inflammatory cytokines IL-8 in HT-29 cells

To investigate the effects of the HMHs on the *E. faecalis*-mediated inflammation in gastrointestinal epithelial cells, the inhibitory properties of the HMH on the *E. faecalis*-mediated induction of a proinflammatory cytokine, IL-8, was estimated by using HT-29 cells, which have been used as an *in-vitro* model of intestinal epithelial cells (Lee et al., 2010). At first, the cytotoxicity of the HMH in the HT-29 cells was evaluated. As shown in Fig. 4A, treatment of HMHs did not show cytotoxicity in HT-29 cells up to 2 mg/mL concentrations.

Based on these data, non-toxicological levels of HMHs ($0.1\sim0.2 \text{ mg/mL}$) were used for estimating the anti-inflammatory properties against *E. faecalis* in HT-29 cells. We then investigated whether or not HMHs can ameliorate the *E. faecalis*-mediated expression of the proinflammatory cytokine IL-8 in HT-29 cells. As shown in Fig. 4B and 4C, administration of *E. faecalis* to HT-29 cells stimulated IL-8 expression when compared with that of the control (non-treated). However, pretreatment of HMHs (HMH-1 and HMH-2) abolished the *E. faecalis*-induced IL-8 expression in HT-29 cells, suggesting that HMHs can prevent *E. faecalis*-mediated inflammation through attenuating the *E. faecalis*-induced expression of the proinflammatory cytokine IL-8.

Hovenia monofloral honey reduced the *E. faecalis*-mediated TLR-2 related inflammatory signaling in HT-29 cells

To further elucidate the action mechanisms of the HMH-mediated anti-inflammatory responses on the gastrointestinal track, the expression of Toll-like receptor-2 (TLR-2), which plays an important role in transmitting the *E. faecalis*-mediated inflammatory signal, and a TLR-2 adaptor protein, MyD88, were investigated. As shown in Fig. 5, administration of *E. faecalis* to HT-29 cells significantly (p < 0.05) increased the expression of TLR-2 as well as MyD88 when compared with that of the control (non-treated). Interestingly, those up-regulations disappeared in HT-29 cells treated with HMHs, suggesting that HMH could be used for attenuating the *E. faecalis*-mediated activation of TLR-2 related inflammatory signal transduction in HT-29 cells. Moreover, we found that administration of *E. faecalis* to HT-29 cells stimulated the phosphorylation of mitogen-activated protein kinases (MAPKs; ERK, p38, and JNK), which is a downstream signaling pathway of TLR-2, and it was significantly (p < 0.05) attenuated by HMH administration (0.2 mg/mL; Fig. 5A and 5D). These results indicate

that HMH could inhibit the *E. faecalis*-induced intracellular over-activation/phosphorylation of MAPKs, which is a critical signaling pathway that transduces the TLR-2 mediated inflammatory signals. Taken together, our data suggest that HMH could effectively ameliorate *E. faecalis*-mediated inflammatory responses through regulating the TLR-2/MyD88/MAPKs signaling pathways in HT-29 cells.

MyD88 inhibitor as well as Hovenia monofloral honey reduce the *E. faecalis*-mediated IL-8 expression in HT-29 cells

Since TLR-2 is known as a key regulatory receptor protein that regulates pathogen-mediated inflammatory responses, TLR-2 binding molecules have been targeted for developing antiinflammatory agents. MyD88 is an adaptor molecule that binds toll-like receptors (TLRs), thereby regulating the inflammatory responses. Therefore, finally, we compared the antiinflammatory efficacy of HMH with that of the MyD88 inhibitor in HT-29 cells. As shown in Fig. 6, although the expression of IL-8 did not increase when HMH (0.2 mg/mL), a control inhibitor (5 μ M), or the MyD88 inhibitor (5 μ M) were administered to the HT-29 cell, it was dramatically increased if *E. faecalis* was administered to HT-29 cells. Interestingly, *E. faecalis*mediated IL-8 expression was significantly (p < 0.05) attenuated by pretreatment of the MyD88 inhibitor and 0.2 mg/mL of HMH has an equal level of efficacy in suppressing *E. faecalis*-induced IL-8 expression in HT-29 cells. These results suggest that MyD88 plays a critical role in *E. faecalis*-induced IL-8 inflammatory responses in HT-29 cells, and support the anti-inflammatory effects of HMH whereby *E. faecalis*-mediated inflammatory responses could be effectively attenuated by HMH through controlling the expression of MyD88 in HT-29 cells.

Discussion

Foodborne disease is often induced by bacterial contamination from bacterial biofilms that are located in food processing plants (Liu et al., 2013; Pesavento et al., 2014). In addition, chronic wounds, which induce severe inflammatory responses, contain bacterial populations as a form of biofilms (Fux et al., 2005). The reason why chronic inflammation has problems in treating with antibiotics is because bacterial biofilms are more resistant to antibiotics that planktonic bacteria (Ngo et al., 2012). Thus, developing effective ways to eliminate existing biofilms and suppress newly formed biofilms may be an important means of attenuating foodborne diseases and chronic inflammatory responses. Therefore, in this study, we investigated whether or not HMH could effectively attenuate *E. faecalis*-mediated biofilm formation. In addition, the regulatory roles of HMH on *E. faecalis*-induced inflammatory responses in HT-29 cells were firstly evaluated.

Honey is a potential nutraceutical food and has also been recognized as a traditional medicine for curing chronic wounds and burns (Majtan, 2014). Since antimicrobial activity is a representative biological property of honey, anti-bacterial and anti-biofilm activities of several monofloral honeys have been investigated (Kim and Kang, 2020; Lu et al., 2019; Lu et al., 2014). Each monofloral honey is made by honeybees using distinct floral sources; therefore, the content and types of phenolic acids are different, and specific ingredients such as methylglyoxalate (MGO), which has a strong anti-bacterial effect, could be included in specific monofloral honeys, thereby showing distinct anti-bacterial activities among monofloral honeys (Lu et al., 2019).

Since honeybees can access honey plants that are far away to produce honey, many types of pollen are often detected even in monofloral honey. Therefore, in general, honey can be

recognized as monofloral honey if the content of major pollen is more than 45% and, if the content of major pollen is increased, the characteristics of the monofloral honey could be clearer (Alvarez-Suarez et al., 2010). The high-purity HMHs that were used in this study were produced by using a net house system; they contained more than 95% single pollen derived from the Hovenia tree (Park et al., 2020). We therefore think the HMHs used in this study are a good source for evaluating the honey potential of HMH. The antioxidant and anti-bacterial properties of HMH were first reported in our previous study (Park et al., 2020) and this study is the second paper to report the functionality of HMH. We found that HMHs can attenuate *E. faecalis*-mediated biofilm formation and stimulate the destruction of *E. faecalis* biofilms, indicating that HMH could be used for inhibiting *E. faecalis*-induced foodborne disease and chronic inflammation. We also found that the planktonic growth of *E. faecalis* was dramatically attenuated by HMH administration, but the EPS formation of *E. faecalis* was not affected by HMH. Thus, these results suggest that HMH-mediated attenuation or stimulation of *E. faecalis*.

Although *E. faecalis* is a common bacteria in the gastrointestinal track, it has been known to induce pathological disorders such as endocarditis and root canal infection (Ch'ng et al., 2019). In addition, food-contaminated *E. faecalis* may enter the gastrointestinal track and cause an inflammatory reaction. It has also been reported that *E. faecalis* is able to stimulate reactive oxygen species (ROS) production in gastrointestinal epithelial cells as well as colonic epithelia cells, which can induce epithelial cell damage (Rokutan et al., 2006; Strickertsson et al., 2013; Wang et al., 2008). In addition, *E. faecalis* infection was reported to be associated with Crohn's disease, a type of severe inflammatory disease (Zhou et al., 2016). Moreover, administration of *E. faecalis* to MKN74 gastric adenocarcinoma cells up-regulated proinflammatory cytokines such as tumor necrosis factor- α and IL-8. These previous reports suggest that *E. faecalis* could

directly induce inflammatory responses in gastrointestinal epithelial cells. In addition, the detailed molecular mechanisms of how *E. faecalis* transduces inflammatory signals to produce proinflammatory cytokines in gastrointestinal epithelial cells still need to be elucidated.

In our study, we demonstrated that administration of *E. faecalis* to HT-29 cells stimulated the expression of a proinflammtory cytokine, IL-8, and it was controlled by TLR-2 mediated signaling. The expression of TLR-2 and its adaptor protein MyD88 was increased when *E. faecalis* was administered to HT-29 cells. In addition, activation of MAPKs, a downstream signaling pathway of TLR-2, was shown when *E. faecalis* was administered to HT-29 cells. Furthermore, we found that *E. faecalis*-mediated expression of IL-8 was abolished by administration of a MyD88 specific inhibitor. These results strongly suggest that *E. faecalis* could induce the inflammatory response in HT-29 cells through activation of a TLR-2 mediated TLR-2 mediated inflammatory signaling could be attenuated by HMH treatments.

Various studies have suggested the anti-inflammatory activity of honey. Indeed, it was reported that clinical use of honey on wounds helps to inhibit infection (Armon, 1980; Phuapradit and Saropala, 1992) and rapid rates of healing were seen on wounds dressed with honey (Bergman et al., 1983; Hejase et al., 1996). In addition, many studies have reported that honey can inhibit inflammation through inhibition of ROS production and cyclooxygenase-2 (COX-2) expression (Hussein et al., 2012; van den Berg et al., 2008). Although the effects of HMHs on the expression of ROS and COX-2 in HT-29 cells was not examined in this study, anti-inflammatory properties of HMHs against *E. faecalis* in HT-29 cells was clearly demonstrated. Administration of HMHs to HT-29 cells effectively suppressed the adhesion of *E. faecalis* to HT-29 cells and thereby attenuated the *E. faecalis*-mediated TLR-2 inflammatory

signaling. Furthermore, our comparative study showed that 0.2 mg/mL of HMH and 5 μ M of the MyD88 inhibitor had equal efficacy in inhibiting IL-8 production by *E. faecalis* treatment. MyD88 is well-known as an intracellular adaptor protein of TLRs that plays a critical role in connecting TLRs and intracellular proteins to transmit signals from TLRs. Since TLRs-mediated signaling is closely involved in the progression of various diseases, the MyD88 protein has been targeted as a therapeutic target and, actually, several MyD88 inhibitors have been developed to treat inflammation-related diseases such as atherosclerosis, cancer, and leukaemia (Chen et al., 2019; Liu et al., 2020; Shiratori et al., 2017). Therefore, the results of our comparative study strongly support that HMS could effectively inhibit *E. faecalis*-mediated inflammatory responses in HT-29 cells.

Conclusion

Collectively, the effects of HMH on *E. faecalis*-mediated biofilm formation and the inflammatory response in HT-29 cells were firstly evaluated in this study. We found that HMH effectively attenuated *E. faecalis*-mediated biofilm formation by attenuating the growth of *E. faecalis*. In addition, we found that HMH could effectively inhibit the *E. faecalis*-mediated inflammatory responses in HT-29 cells through controlling the TLR-2/MyD88/MAPKs pathway. Taken together, these data suggest that HMH could be used to develop an antimicrobial agent for preventing *E. faecalis*-mediated inflammatory diseases.

Conflicts of Interest

All authors declare no potential conflict of interest.

Acknowledgments

This research was supported by an Incheon National University research grant in 2019. Hovenia monofloral honeys (HMHs) which produced by using a net house system were provided from National Institute of Forest Science in South Korea.

References

- Allen KL, Molan PC, Reid GM. 1991. A survey of the antibacterial activity of some new zealand honeys. J Pharm Pharmacol 43:817-822.
- Alvarez-Suarez JM, Tulipani S, Diaz D, Estevez Y, Romandini S, Giampieri F, Damiani E, Astolfi P, Bompadre S, Battino M. 2010. Antioxidant and antimicrobial capacity of several monofloral cuban honeys and their correlation with color, polyphenol content and other chemical compounds. Food Chem Toxicol 48:2490-2499.
- Armon PJ. 1980. The use of honey in the treatment of infected wounds. Trop Doct 10:91.
- Baccouri O, Boukerb AM, Farhat LB, Zebre A, Zimmermann K, Domann E, Cambronel M, Barreau M, Maillot
 O, Rince I, Muller C, Marzouki MN, Feuilloley M, Abidi F, Connil N. 2019. Probiotic potential and safety evaluation of enterococcus faecalis ob14 and ob15, isolated from traditional tunisian testouri cheese and rigouta, using physiological and genomic analysis. Front Microbiol 10:881.
- Bergman A, Yanai J, Weiss J, Bell D, David MP. 1983. Acceleration of wound healing by topical application of honey. An animal model. Am J Surg 145:374-376.
- Bogdanov S, Jurendic T, Sieber R, Gallmann P. 2008. Honey for nutrition and health: A review. J Am Coll Nutr 27:677-689.
- Ch'ng JH, Chong KKL, Lam LN, Wong JJ, Kline KA. 2019. Biofilm-associated infection by enterococci. Nat Rev Microbiol 17:82-94.
- Chen T, Luo W, Wu G, Wu L, Huang S, Li J, Wang J, Hu X, Huang W, Liang G. 2019. A novel myd88 inhibitor lm9 prevents atherosclerosis by regulating inflammatory responses and oxidative stress in macrophages. Toxicol Appl Pharmacol 370:44-55.
- Da Silva PM, Gauche C, Gonzaga LV, Costa AC, Fett R. 2016. Honey: Chemical composition, stability and authenticity. Food Chem 196:309-323.
- Engemann JJ, Carmeli Y, Cosgrove SE, Fowler VG, Bronstein MZ, Trivette SL, Briggs JP, Sexton DJ, Kaye KS. 2003. Adverse clinical and economic outcomes attributable to methicillin resistance among patients with staphylococcus aureus surgical site infection. Clin Infect Dis 36:592-598.
- Fux CA, Costerton JW, Stewart PS, Stoodley P. 2005. Survival strategies of infectious biofilms. Trends Microbiol 13:34-40.

Giraffa G. 2002. Enterococci from foods. FEMS Microbiol Rev 26:163-171.

- Hejase MJ, Simonin JE, Bihrle R, Coogan CL. 1996. Genital fournier's gangrene: Experience with 38 patients. Urology 47:734-739.
- Hoyle BD, Costerton JW. 1991. Bacterial resistance to antibiotics: The role of biofilms. Prog Drug Res 37:91-105.
- Hussein SZ, Mohd Yusoff K, Makpol S, Mohd Yusof YA. 2012. Gelam honey inhibits the production of proinflammatory, mediators no, pge(2), tnf-alpha, and il-6 in carrageenan-induced acute paw edema in rats. Evid Based Complement Alternat Med 2012:109636.
- Kim SY, Kang SS. 2020. Anti-biofilm activities of manuka honey against escherichia coli o157:H7. Food Sci Anim Resour 40:668-674.
- Lee K, Lee KM, Kim D, Yoon SS. 2017. Molecular determinants of the thickened matrix in a dual-species pseudomonas aeruginosa and enterococcus faecalis biofilm. Appl Environ Microbiol 83.
- Lee SH, Yu SY, Nakayama J, Khoo KH, Stone EL, Fukuda MN, Marth JD, Fukuda M. 2010. Core2 o-glycan structure is essential for the cell surface expression of sucrase isomaltase and dipeptidyl peptidase-iv during intestinal cell differentiation. J Biol Chem 285:37683-37692.
- Liu F, Du L, Xu W, Wang D, Zhang M, Zhu Y, Xu W. 2013. Production of tyramine by enterococcus faecalis strains in water-boiled salted duck. J Food Prot 76:854-859.
- Liu JH, Chen C, Li ZY, Zou ZM, Gao DC, Zhang X, Kuang XW, Sun ZH, Zheng WJ, Zhou P, Sun SR. 2020. The myd88 inhibitor tj-m2010-2 suppresses proliferation, migration and invasion of breast cancer cells by regulating myd88/gsk-3beta and myd88/nf-kappab signalling pathways. Exp Cell Res 394:112157.
- Lu J, Cokcetin NN, Burke CM, Turnbull L, Liu M, Carter DA, Whitchurch CB, Harry EJ. 2019. Honey can inhibit and eliminate biofilms produced by pseudomonas aeruginosa. Sci Rep 9:18160.
- Lu J, Turnbull L, Burke CM, Liu M, Carter DA, Schlothauer RC, Whitchurch CB, Harry EJ. 2014. Manuka-type honeys can eradicate biofilms produced by staphylococcus aureus strains with different biofilm-forming abilities. PeerJ 2:e326.
- Majtan J. 2014. Honey: An immunomodulator in wound healing. Wound Repair Regen 22:187-192.
- N AL-W, Al Ghamdi A, Ansari MJ, Al-Attal Y, Al-Mubarak A, Salom K. 2013. Differences in composition of honey samples and their impact on the antimicrobial activities against drug multiresistant bacteria and pathogenic fungi. Arch Med Res 44:307-316.

- Ngo QD, Vickery K, Deva AK. 2012. The effect of topical negative pressure on wound biofilms using an in vitro wound model. Wound Repair Regen 20:83-90.
- Park SH, Kim YK, Kim MS, Lee SH. 2020. Antioxidant and antibacterial properties of hovenia (hovenia dulcis) monofloral honey produced in south korea. Food Sci Anim Resour 40:221-230.
- Pesavento G, Calonico C, Ducci B, Magnanini A, Lo Nostro A. 2014. Prevalence and antibiotic resistance of enterococcus spp. Isolated from retail cheese, ready-to-eat salads, ham, and raw meat. Food Microbiol 41:1-7.
- Phuapradit W, Saropala N. 1992. Topical application of honey in treatment of abdominal wound disruption. Aust N Z J Obstet Gynaecol 32:381-384.
- Qiu PY, Ding HB, Tang YK, Xu RJ. 1999. Determination of chemical composition of commercial honey by nearinfrared spectroscopy. J Agric Food Chem 47:2760-2765.
- Ranneh Y, Akim AM, Hamid HA, Khazaai H, Fadel A, Zakaria ZA, Albujja M, Bakar MFA. 2021. Honey and its nutritional and anti-inflammatory value. BMC Complement Med Ther 21:30.
- Rokutan K, Kawahara T, Kuwano Y, Tominaga K, Sekiyama A, Teshima-Kondo S. 2006. Nadph oxidases in the gastrointestinal tract: A potential role of nox1 in innate immune response and carcinogenesis. Antioxid Redox Signal 8:1573-1582.
- Sherlock O, Dolan A, Athman R, Power A, Gethin G, Cowman S, Humphreys H. 2010. Comparison of the antimicrobial activity of ulmo honey from chile and manuka honey against methicillin-resistant staphylococcus aureus, escherichia coli and pseudomonas aeruginosa. BMC Complement Altern Med 10:47.
- Shiratori E, Itoh M, Tohda S. 2017. Myd88 inhibitor st2825 suppresses the growth of lymphoma and leukaemia cells. Anticancer Res 37:6203-6209.
- Stagos D, Soulitsiotis N, Tsadila C, Papaeconomou S, Arvanitis C, Ntontos A, Karkanta F, Adamou-Androulaki S, Petrotos K, Spandidos DA, Kouretas D, Mossialos D. 2018. Antibacterial and antioxidant activity of different types of honey derived from mount olympus in greece. Int J Mol Med 42:726-734.
- Strickertsson JA, Desler C, Martin-Bertelsen T, Machado AM, Wadstrom T, Winther O, Rasmussen LJ, Friis-Hansen L. 2013. Enterococcus faecalis infection causes inflammation, intracellular oxphos-independent ros production, and DNA damage in human gastric cancer cells. PLoS One 8:e63147.

- Van Den Berg AJ, Van Den Worm E, Van Ufford HC, Halkes SB, Hoekstra MJ, Beukelman CJ. 2008. An in vitro examination of the antioxidant and anti-inflammatory properties of buckwheat honey. J Wound Care 17:172-174, 176-178.
- Wang X, Allen TD, May RJ, Lightfoot S, Houchen CW, Huycke MM. 2008. Enterococcus faecalis induces aneuploidy and tetraploidy in colonic epithelial cells through a bystander effect. Cancer Res 68:9909-9917.
- Wolcott RD, Rhoads DD, Dowd SE. 2008. Biofilms and chronic wound inflammation. J Wound Care 17:333-341.
- Yoon JW, Kang SS. 2020. In vitro antibiofilm and anti-inflammatory properties of bacteriocins produced by pediococcus acidilactici against enterococcus faecalis. Foodborne Pathog Dis 17:764-771.
- Zhou Y, Chen H, He H, Du Y, Hu J, Li Y, Li Y, Zhou Y, Wang H, Chen Y, Nie Y. 2016. Increased enterococcus faecalis infection is associated with clinically active crohn disease. Medicine (Baltimore) 95:e5019.
- Zommiti M, Cambronel M, Maillot O, Barreau M, Sebei K, Feuilloley M, Ferchichi M, Connil N. 2018. Evaluation of probiotic properties and safety of enterococcus faecium isolated from artisanal tunisian meat "dried ossban". Front Microbiol 9:1685.

Figure legends



Fig. 1. Hovenia monofloral honeys (HMHs) attenuated the formation of *E. faecalis* biofilms. *E. faecalis* was cultured on a black polystyrene 96-well microtiter plate with a clear bottom with and without HMHs (0~32%). The *E. faecalis* biofilm was stained with SYTO9 (A) and the intensity of the SYTO9 positive region (green) was quantified using the ImagJ program (B). The data shown are representative of at least three independent experiments. Scale bar = 100 μ m. Data are presented as mean ± SD. *: significant differences (*p* < 0.05) when compared with the control (non-treated).



Fig. 2. Hovenia monofloral honeys (HMHs) eradicated the *E. faecalis* biofilms. Various concentration of HMHs (0~32%) were added to *E. faecalis* biofilms and incubated for an additional 24 h. The remaining *E. faecalis* biofilms were stained with SYTO9 solution (A) and the positive area (green) was quantified using the ImageJ program (B). The data shown are representative of at least three independent experiments. Scale bar = 100 μ m. Data are presented as mean ± SD. *: significant differences (*p* < 0.05) when compared with the control (non-treated).

Figure 3.



Fig. 3. Hovenia monofloral honeys (HMHs) attenuated the growth of *E. faecalis*. HMHs (0, 16, and 32%, w/v) were co-incubated with *E. faecalis* for 2, 4, and 6 h (A) or 24 h (B). The effect of the HMHs (0, 16, and 32%, w/v) on the planktonic growth of *E. faecalis* was evaluated by measuring the optical density (A) and WST-1 cell viability kit (B). The effect of the HMHs (0, 16, and 32%, w/v) on the production of extracellular polymeric substances (ESPs) was estimated by Congo red binding assay according to the protocol described in the materials and methods section (C). The data shown are representative of at least three independent experiments. Data are presented as mean \pm SD. *: significant differences (p < 0.05) when compared with the control (non-treated).

Figure 4.



Fig. 4. Hovenia monofloral honeys (HMHs) attenuated the *E. faecalis* (E.F) induced expression of IL-8 in HT-29 gastrointestinal epithelial cells. The cytotoxicity of HMHs (HMH-1 and HMH-2) against HT-29 cells was estimated using a WST-1 assay kit (A). *E. faecalis* (1×10^6 CFU/mL) was administered to HT-29 cells with and without HMHs for 24 h. The expression of the IL-8 gene was determined by qRT-PCR. The data shown are representative of at least three independent experiments. Data are presented as mean ± SD. Different letters indicate the significant (p < 0.05) differences between groups.



Fig. 5. Hovenia monofloral honeys (HMHs) attenuated the *E. faecalis*-mediated TLR-2 linked inflammatory signaling in HT-29 cells. *E. faecalis* (E.F) (1×10^6 CFU/mL) was administered to HT-29 cells with and without HMHs for 24 h. The expression of TLR-2, MyD88, and MAPKs (ERK, p38, and JNK) was estimated by western blotting (A) and the intensity of each band was measured using the imageJ program and tabulated (B). The data shown are representative of at least three independent experiments. Data are presented as mean ± SD. Different letters indicate the significant (p < 0.05) differences between groups.

Figure 6.



Fig. 6. *E. faecalis*-induced expression of the IL-8 gene was abolished by administering the MyD88 inhibitor as well as Hovenia monofloral honeys (HMHs). HT-29 cells were administered with *E. faecalis* (E.F) (1×10^6 CFU/mL) in the presence or absence of either the MyD88 inhibitor (5 µM) or HMHs (0.2 mg/mL). The expression of the IL-8 gene was estimated by qRT-PCR. Administration of the HMHs (0.2 mg/mL), the control inhibitor (5 µM), or the MyD88 inhibitor (5 µM) without administration of *E. faecalis* to the HT-29 cells did not stimulate the expression of IL-8. The data shown are representative of at least three independent experiments. Data are presented as mean ± SD. Different letters indicate the significant (p < 0.05) differences between groups.