

**TITLE PAGE**  
**- Food Science of Animal Resources -**  
**Upload this completed form to website with submission**

ARTICLE INFORMATION	Fill in information in each box below
<b>Article Type</b>	Review article
<b>Article Title</b>	Animal Models of Cognitive Deficits for Probiotic Treatment
<b>Running Title (within 10 words)</b>	Probiotics has anti-cognitive activities
<b>Author</b>	Oh Yun Kwon <sup>1</sup> , Seung Ho Lee <sup>1</sup>
<b>Affiliation</b>	1 Department of Nano-Bioengineering, Incheon National University, Incheon, Korea
<b>Special remarks</b> – if authors have additional information to inform the editorial office	
<b>ORCID (All authors must have ORCID) <a href="https://orcid.org">https://orcid.org</a></b>	Oh Yun Kwon ( <a href="https://orcid.org/0000-0002-3783-3220">https://orcid.org/0000-0002-3783-3220</a> ) Seung Ho Lee ( <a href="https://orcid.org/0000-0002-9941-4195">https://orcid.org/0000-0002-9941-4195</a> )
<b>Conflicts of interest</b> List any present or potential conflict s of interest for all authors. (This field may be published.)	The authors declare no potential conflict of interest.
<b>Acknowledgements</b> State funding sources (grants, funding sources, equipment, and supplies). Include name and number of grant if available. (This field may be published.)	This research was supported by an Incheon National University research grant (2018).
<b>Author contributions</b> (This field may be published.)	Conceptualization: Lee SH, Investigation: Kwon OY, Writing–original draft: Kwon OY, and Writing, review, and editing: Lee SH.
<b>Ethics approval (IRB/IACUC)</b> (This field may be published.)	This article did not require approval from the IACUC/IRB since there were no animal and human participants.

**CORRESPONDING AUTHOR CONTACT INFORMATION**

For the <u>corresponding</u> author (responsible for correspondence, proofreading, and reprints)	Fill in information in each box below
First name, middle initial, last name	Seung Ho, Lee
Email address – this is where your proofs will be sent	Lsh1602@gmail.com
Secondary Email address	seungho@inu.ac.kr
Postal address	22012
Cell phone number	010-8014-1614
Office phone number	
Fax number	

## Animal Models of Cognitive Deficits for Probiotic Treatment

### Abstract

Cognitive dysfunction is a common symptom of neurodegenerative diseases, such as Alzheimer's disease, Parkinson's disease, and Huntington's disease, and is known to be caused by the structural and functional loss of neurons. Many natural agents that can improve cognitive function have been developed and assessed for efficacy using various cognitive deficit animal models. As the gut environment is known to be closely connected to brain function, probiotics are attracting attention as an effective treatment target that can prevent and mitigate cognitive deficits as a result of neurodegenerative diseases. Thus, the objective of this review is to provide useful information about the types and characteristics of cognitive deficit animal models, which can be used to evaluate the anti-cognitive effects of probiotics. In addition, this work reviewed recent studies describing the effects and treatment conditions of probiotics on cognitive deficit animal models. Collectively, this review shows the potential of probiotics as edible natural agents that can mitigate cognitive impairment. It also provides useful information for the design of probiotic treatments for cognitive deficit patients in future clinical studies.

**Keywords:** Cognitive deficits, Probiotics, Neurodegenerative disease, Gut–Brain Axis

## Introduction

With rapid social development and the continued introduction of new medical technology, human life expectancy is gradually increasing, causing many countries around the world to become aging societies. Although aging is closely associated with cognitive decline, a broad range of different cognitive abilities was found even in people of the same age, and severe cognitive impairment could be detected in patients suffering from neurodegenerative diseases, such as Alzheimer's disease (AD) and Parkinson's disease (PD) (Chiu et al., 2006; Christensen et al., 1999). This indicates that age-dependent cognitive decline may be controlled. It has been reported that the prevalence of dementia is doubling every 20 years and is expected to reach 131.5 million cases by 2050 (Alladi et al., 2011; Bowler et al., 1998). Therefore, huge social and economic costs will be incurred as a result. As such, there is increasing demand for the development of efficient drugs that can treat dementia.

Many agents have been developed to mitigate cognitive deficits, a major effect of dementia. Since AD is the most common neurodegenerative disease, which manifests as severe cognitive impairments, researchers have focused on developing agents that can inhibit the amyloid beta ( $A\beta$ ) peptide-mediated cascade events, such as oxidative stress, mitochondria dysfunction, and neuronal death in the brain. Many natural agents, which can attenuate  $A\beta$ -induced neurological disorder with cognitive declines, have been developed from foods (Ravi et al., 2019), plants (Akter et al., 2021; Deng et al., 2020), and seaweed (Kwon and Lee, 2020; Kwon and Lee, 2021). Recently, various studies have reported that the gut environment can interact closely with the brain via the nervous system (Collins et al., 2012) and chemicals (Briguglio et al., 2018). In the connection between the brain and gut, the so-called gut-brain axis, gut microbiota have been proven to be a major regulator since they can exchange information between the gut and brain and also produce neuromodulators, such as short-chain fatty acids, glutamate, and serotonin. These neuromodulators can reach the central

neurons via the circulatory system, finally affecting neuronal activity and causing behavioral changes (Logsdon et al., 2018).

Probiotics, which are defined as “viable microbial food supplements that have beneficial effects on human health” (Salminen et al., 1998), have been studied for their effects on various diseases, such as inflammation (Vitetta et al., 2012) and immunological disorders (Vitetta et al., 2018). Since it is known that controlling the gut microbiota is important in regulating neurological disorders (Morais et al., 2021), many efforts have attempted to use probiotics to treat neurodegenerative diseases. Since a huge number of probiotic strains have been developed, the demand for an accurate estimating system to test each probiotic strain is gradually increasing. Therefore, the use of proper cognitive deficit animal models is important to evaluate the anti-cognitive deficit function of each probiotic strain. In this paper, we summarize various animal models and probiotic strains that have been used to test the effects of anti-cognitive deficits. These results provide useful information about ways to further the development of probiotic-based drugs for human treatments

## **Chemical-induced cognitive deficit animal models for probiotic treatment**

Various chemical-induced cognitive deficit animal models have been developed and used to test the anti-cognitive deficit activities of probiotics. The characteristics of each animal model are summarized in Table 1, and the treatment conditions and probiotic strains used in these mice are listed in Table 3.

### **D-galactose-induced accelerated aging mouse model**

Since D-galactose (D-gal), a normal carbohydrate, can be found in various foods, such as cheese, butter, and honey, D-gal is absorbed into the body through the intake

of these foods and used in the biochemical pathways to produce metabolic energy. However, high levels of D-gal in the body are considered an oxidative stressor to cells because D-gal can be oxidized by galactose oxidase, resulting in aldehyde and hydrogen peroxide production. Thus, the chronic administration of D-gal to rodents could lead to oxidative damage to cells, including those in the brain, resulting in a progressive loss of memory. Many studies have shown that mitochondrial dysfunction, oxidative stress, neuronal death, and cognitive deficits occur in mouse brains when excess D-gal is supplied (Kumar et al., 2009; Prakash and Kumar, 2013; Rehman et al., 2017). Therefore, a D-gal-induced rodent model could be used to test the anti-cognitive deficit effects of new agents, including probiotics.

To make D-gal-induced cognitive deficit rodent models, 100–200 mg/Kg body weight/day of D-gal should be injected for 6–10 weeks, which seems to be enough doses to cause progressive loss of memory (Lu et al., 2010; Shwe et al., 2018). In most reports, the candidate materials were continuously fed from the time D-gal was injected to the end of the D-gal injection period. A few reports have demonstrated the anti-cognitive deficit function of probiotics by using the D-gal-induced cognitive deficits model. For example, *Lactobacillus paracasei* PS23, which is isolated from human feces, has been reported to have an anti-cognitive deficit function. *L. paracasei* PS23 ( $10^9$  CFU/d/mice) was administrated to D-gal (100 mg/kg bw/d)-injected mice for 10 weeks to attenuate anxiety-like behavior and prevent the loss of long-term memory induced by D-gal (Cheng et al., 2022). The oral administration of *L. pentosus var. plantarum* C29 ( $10^{10}$  CFU/d/mice) isolated from kimchi, a traditional Korean food, to D-gal-injected mice (100 mg/kg bw/d of D-gal for 10 weeks) ameliorated D-gal-induced memory impairment (Woo et al., 2014). (Song et al., 2022) reported that the oral administration of *Bacillus coagulans* JA845 ( $10^9$  CFU /d/mice) to D-gal-induced cognitive deficit mice (injected with 120 mg/kg bw/d of D-gal for 10 weeks) inhibited oxidative stress in the brain, loss of hippocampal neurons, and loss of long-term memory. Collectively, these

data suggest that probiotics other than the above strains have great potential for attenuating D-gal-induced cognitive impairment.

### **Amyloid beta (A $\beta$ )-induced cognitive deficits model**

A $\beta$  is a 4 kDa peptide of 36–43 amino acids generated from the amyloid precursor protein (APP) by the proteolytic reaction of  $\beta$ -secretase and  $\gamma$ -secretase (Wang et al., 2021). A $\beta$  aggregates to form amyloid plaques, which can induce toxicity to neuronal cells, and amyloid plaques are easily detected in the brains of AD patients. Therefore, A $\beta$ -mediated neurotoxicity is considered a major target in developing anti-AD drugs. To make a cognitive deficit animal model that mimics AD, intracerebroventricular (ICV) injection of A $\beta$  has been used, and once A $\beta$  is injected into the mice brain directly, neuronal death followed by loss of learning and memory function could be detected from 3–6 d after injection (Ali et al., 2015; Kobayashi et al., 2017; Kwon and Lee, 2020). To construct the A $\beta$ -induced cognitive impairment animal model, stereotaxic apparatuses are often used to accurately inject A $\beta$  into the bregma region without brain damage. However, the direct injection method without stereotaxic apparatuses is also used to introduce cognitive deficits in mice by the direct injection of A $\beta$  peptide (Kim et al., 2016). Most reports injected 100  $\mu$ mol of A $\beta$  peptide, and cognitive impairment could be detected with a single injection. Therefore, A $\beta$ -injected mouse can be used to estimate the anti-AD activity of drug candidates.

Probiotics have been reported to be a potential candidate for preventing A $\beta$ -mediated neurodegeneration (Kobayashi et al., 2017). The oral administration of *Bifidobacterium breve* A1 ( $1 \times 10^9$  /d/mice) to A $\beta$ -injected mice for 10 d inhibited A $\beta$ -mediated cognitive dysfunction (Kobayashi et al., 2017). (Zhu et al., 2021) reported that oral treatment with *B. breve* CCFM1025 and WX ( $0.6 \times 10^9$ /d/mice) for 6 weeks recovered A $\beta$ -mediated synaptic plasticity decrease through the regulation of the gut

microbiome. Thus, these results suggest that a cognitive deficit animal model induced by A $\beta$  injection could be used to evaluate the regulatory function of probiotics in cognitive impairments.

### **Scopolamine-induced cognitive deficits in mice**

Acetylcholine is a neurotransmitter that plays an important role in cortical development, sleep–wake cycles, and memory and cognitive function (Bruehl-Jungerman et al., 2011; Lopez-Sobaler et al., 2021; Van Erum et al., 2019). Reduced cholinergic activity with a loss of cortical cholinergic neurons in the hippocampus is often detected in neurodegeneration, and it has been considered a cause of memory impairment in AD (Whitehouse et al., 1981; Whitehouse et al., 1982). Therefore, inhibiting the central cholinergic system has been proposed as a way to build an animal model with limited cognitive function. Scopolamine is an anti-cholinergic drug that blocks the binding between acetylcholine and muscarinic receptors, resulting in the excess release of acetylcholine (Lazareno et al., 2000). Thus, the administration of scopolamine to mice results in the loss of hippocampus neurons and learning and memory impairments. A single intraperitoneal injection of scopolamine (1–1.5 mg/kg bw) to mice could induce cognitive impairment within 1 h, but repetitive administration of scopolamine (once/d) is recommended during behavioral tests (Choi et al., 2021; Kim et al., 2021b; Yadang et al., 2020).

Several studies have shown that scopolamine-induced cognitive impairment can be attenuated by probiotic treatment. For example, the administration of *L. johnsonii* CJLJ103 ( $1 \times 10^9$  CFU/d/mice) for 5 d restored the scopolamine-induced decrease of spontaneous alteration (%) estimated by the Y-maze test (Lee et al., 2018). (Patel et al., 2020) reported that the oral administration of *L. rhamnosus* UBLR-58 ( $1 \times 10^9$  CFU/d/mice) for 10 d enhanced curcumin's effects against scopolamine-induced cognitive deficits. Collectively, these studies suggest that probiotics could be active

agents that control for cognitive deficits induced by the impairment of the cholinergic system.

## **LPS-induced cognitive deficits in mice**

Systemic inflammation induced by infectious agents has been recognized as a cause of cognitive dysfunctions (Banks et al., 2002; Konsman et al., 2002), and elevated inflammatory responses increase the amount of circulating proinflammatory cytokines that can change the central nerve system (Perry, 2004). In addition, it has been suggested that inflammatory cytokines can stimulate A $\beta$  production and attenuate the secretion of APPs (Blasko et al., 1999; Buxbaum et al., 1992). Thus, the association between chronic infection and AD has been intensively studied to prevent cognitive deficits (Panza et al., 2019). Since neuroinflammation can occur via infiltration from Gram-negative bacteria, such as *Chlamydia pneumoniae* and *Porphyromonas gingivalis* (MacIntyre et al., 2003; Shoemark and Allen, 2015), lipopolysaccharide (LPS), which is a cell wall component of Gram-negative bacteria, has been widely used to construct cognitive deficit animal models. Injection of LPS (0.25–1 mg/kg bw/d) in the abdominal cavity of mice for 4–7 consecutive d could produce neuroinflammatory responses with cognitive impairments. LPS can also be administered directly to brain tissue via ICV injection (Kamdi et al., 2021; Shoemark and Allen, 2015; Yang et al., 2020a). A single ICV injection of LPS (2–12  $\mu$ g) solution using a micro syringe or stereotaxic coordinates could induce neuroinflammatory responses with cognitive deficits from 1–7 d after injection (Zhao et al., 2019; Zhou et al., 2006). Therefore, an LPS-induced animal model could be used to evaluate the anti-cognitive deficit effects of probiotics. In fact, the *L. plantarum* NK51 and *B. longum* KN173 strains were reported to have preventive activity against LPS-induced neuroinflammation (Lee et al., 2021), and probiotic mixtures (*L. helveticus* R0052 and



*B. longum* R0175) were also proven to have anti-cognitive deficit effects on LPS-induced rat models (Mohammadi et al., 2019). In addition, it was reported that an oral gavage of *Lactococcus lactis* subsp. *cremois* LL95 ( $1 \times 10^9$  CFU/d/mice) for 7 d ameliorated mood disorders in an LPS-induced depression-like mice model (Ramalho et al., 2022), and the administration of a probiotic mixture (*B. animalis* subsp. *lactis* BL03, *B. animalis* subsp. *lactis* BI04, *B. breve* BB02, *L. acidophilus* BA05, *L. helveticus* BD08, *L. paracasei* BP07, *L. plantarum* BP06, and *Streptococcus thermophilus* BT01) ( $1 \times 10^9$  CFU/d/mice) for 15 d attenuated LPS-induced pro-inflammatory responses and sickness behavior (Petrella et al., 2021). These reports suggest that probiotics could be developed as effective agents that can ameliorate LPS-mediated neuroinflammation and cognitive deficits.

### Transgenic animal models that have cognitive deficit phenotypes

Various transgenic animal models that show cognitive impairment have been developed. Each mouse line showed subtly different characteristics in neuronal development and behavior disorders. The characteristics of the cognitive deficit transgenic animal models used for evaluating the anti-cognitive deficit effects of probiotics are summarized in Table 2, and the probiotic strains, treatment conditions, and behavior changes are listed in Table 3.

#### Senescence-accelerated mouse (SAM)

The senescence-accelerated mouse prone 8 (SAMP8) mouse line is derived from mice that have a naturally accelerated aging phenotype (Takeda et al., 1981). The SAMP8 mouse line shows an age-dependent increase in A $\beta$  in the hippocampal area from 4 to 12 months after birth. SAMP8 mice exhibit age-associated cognitive impairment (from 8 to 10 months), reduced anxiety-like behavior, and reduced

lifespan (Miyamoto et al., 1992; Miyamoto et al., 1986). Although the detailed molecular mechanisms behind early senescence in SAMP8 mice have not been fully elucidated, excessive oxidative stress has been suggested as a cause of cognitive dysfunction in the SAMP8 mouse line (Morley et al., 2012). Therefore, the SAMP8 mouse line could be an excellent animal model for studying age-dependent cognitive deficits.

Intriguingly, reports have shown that short- and long-term memory loss in SAMP8 mice was attenuated through the oral administration of a probiotic mixture (Yang et al., 2020b). SAMP8 mice (9-month-old males) were orally administered ProBiotic-4, a mixture of *B. lactis* (50%), *B. bifidum* (12.5%), *L. acidophilus* (12.5%), and *L. casei* (25%), for 12 weeks, and showed that memory deficits and neuronal injury were improved, and aging-related disruption of the intestinal barrier was attenuated. These results suggest that SAMP8 mice could be used as an animal model to evaluate probiotic effects on age-related cognitive impairment.

### ***App*<sup>NL-G-F</sup> mice**

The *App*<sup>NL-G-F</sup> mice are transgenic mice constructed by the knock-in of the *APP* gene containing four amino acids. In *App*<sup>NL-G-F</sup> mice, the lysine (K) and methionine (M) amino acids at the 670/671 positions of the *APP* gene are substituted with asparagine (N) and leucine (L), respectively (KM670/671NL). In addition, isoleucine (I) at the 716 position and glutamic acid (E) at the 693 position of the *APP* gene are mutated to phenylalanine (F) (I716F) and glycine (G) (E693G), respectively. These four mutation sites of the *APP* gene were found in human families that struggled with cognitive deficits (Guerreiro et al., 2010; Mullan et al., 1992; Nilsberth et al., 2001). The *App*<sup>NL-G-F</sup> mouse line shows aggressive A $\beta$  amyloidosis in an age-dependent manner. *App*<sup>NL-G-F</sup> mice show A $\beta$  deposition from 2 months after birth, and maximum memory impairment is shown at 12 months (Mehla et al., 2019; Saito et al., 2014). Therefore,

*App*<sup>NL-G-F</sup> mice are considered to have cognitive deficits in animal models that represent the typical A $\beta$  pathology.

It was reported that the oral administration of VSL#3, a probiotic mixture ( $1.28 \times 10^9$  CFU/d/mice) that consists of *L. plantarum*, *L. delbrueckii* subsp. *Bulgaricus*, *L. paracasei*, *L. acidophilus*, *B. breve*, *B. longum*, *B. infantis*, and *S. salivarius* subsp. *thermophiles*, for 8 weeks reduced intestinal inflammation and anxiety-like behavior in *App*<sup>NL-G-F</sup> mice (Kaur et al., 2020a; Kaur et al., 2020b). Although not many probiotic trials on *App*<sup>NL-G-F</sup> mice have been reported, *App*<sup>NL-G-F</sup> mice are thought to be an attractive animal model for estimating the probiotic effects on A $\beta$  amyloidosis.

#### ***APP/PS1* transgenic mice**

*APP/PS1* transgenic mice are created by introducing a human *APP* gene containing KM670/671NL mutations and presenilin-1 (PSEN1) possessing an L166P mutation. *APP/PS1* transgenic mice exhibit increased A $\beta$  production in an age-dependent manner, and A $\beta$  deposition is detected in the neocortex from approximately 6 weeks after birth. A $\beta$  deposition on the hippocampus can be detected from 3 to 4 months of age, and *APP/PS1* transgenic mice are known to show spatial learning and memory impairment from 7 months of age (Lok et al., 2013; Radde et al., 2006).

*APP/PS1* transgenic mice have also been used to estimate probiotics' effects on cognitive deficits. Oral administration of the *L. plantarum* ATCC 8014 strain ( $1 \times 10^9$  CFU/d/mice) to *APP/PS1* mice for 12 weeks ameliorated A $\beta$  deposition in the hippocampus and cognitive impairment (Wang et al., 2020). The administration of *B. lactis* Probio-M8 ( $2 \times 10^{10}$  CFU/d/mice) to *APP/PS1* transgenic mice for 45 d attenuated A $\beta$  plaque formation and improved spatial working memory, which was estimated by using the Y-maze test (Cao et al., 2021). (Sun et al., 2020) reported that the daily administration of *Clostridium butyricum* WZMC1016 ( $0.2 \times 10^9$  CFU/d/mice) for 4 weeks

effectively ameliorated A $\beta$  deposition, microglia activation, and cognitive impairments in *APP/PS1* transgenic mice. The evidence strongly suggested that *APP/PS1* transgenic mice could be used to evaluate probiotics' effects on age-dependent amyloid deposition accompanied by cognitive dysfunction.

### **3xTg-AD mice**

3xTg-AD mice are developed by integrating the three mutated AD-related genes, specifically *APP*, *PSEN1*, and microtubule-associated protein Tau (*MAPT*). 3xTg-AD mice harboring *APP* (KM670/671NL), *PSEN1* (M146V), and *MAPT* (P301L) genes showed progressive A $\beta$  deposition from 3 to 4 months of age, and hyperphosphorylated tau protein could be detected at 12–15 months of age (Oddo et al., 2003). 3xTg-AD mice were reported to have mildly impaired spatial learning and memory function, and the Barnes maze test was determined to be the most sensitive way to estimate cognitive deficits in 3xTg-AD mice (Stover et al., 2015). Thus, 3xTg-AD mice have been recognized as a valuable animal model for estimating anti-AD therapeutics.

3xTg-AD mice have also been used to evaluate the anti-AD effects of probiotics. (Bonfili et al., 2017) reported that oral treatment with SLAB51, a probiotic mixture that consists of *S. thermophilus*, *B. longum*, *B. breve*, *B. infantis*, *L. acidophilus*, *L. plantarum*, *L. paracasei*, *L. delbrueckii* subsp. *bulgaricus*, and *L. brevis*, to 3xTg-AD mice (8-week-old males) for 4 months attenuated A $\beta$  aggregation, brain damage, and cognitive decline. Another probiotic mixture, Lab4b, which consists of *L. salivarius* CUL61 (NCIMB 30211), *L. paracasei* CUL08 (NCIMB 30154), *B. bifidum* CUL20 (NCIMB 30153), and *B. animalis* subsp. *lactis* CUL34 (NCIMB 30172), was proven to have anti-AD effects on the 3xTg-AD mouse model. Oral treatment with Lab4b ( $5 \times 10^8$  CFU/d/mice) in 12-week-old 3xTg-AD mice for 12 weeks improved recognition memory sensitivity,

which was estimated using the novel objection recognition (NOR) test (Webberley et al., 2022). Collectively, these reports support the possibility of using 3xTg-AD mice to evaluate the anti-cognitive impairment effects of probiotics.

## **5xFAD mice**

It has been reported that mutations in AD-related genes, such as *APP* and presenilins (PS1 and PS2), are closely associated with familial AD (FAD), which express high levels of A $\beta$ . 5xFAD mice is an *APP/PS1* transgenic mouse model that expresses five FDA mutations (*APP*<sup>KM670/671NL</sup>, *APP*<sup>I716V</sup>, *APP*<sup>V717I</sup>, *PSEN1*<sup>M146L</sup>, and *PSEN1*<sup>L286V</sup>) that result in the rapid accumulation of cerebral A $\beta$ . The advantage of 5xFAD mice is that intraneuronal A $\beta$  accumulation was detected at 1.5 months of age, which is relatively early, and neuronal loss started from 6 months of age with amyloidosis. In addition, it was reported that the impairment of spatial working memory was detected at 4–5 months (Devi and Ohno, 2010; Oakley et al., 2006). Therefore, 5xFAD mice could be used as an animal model for estimating the effects of anti-cognitive deficit agents.

Several reports have assessed functional probiotics against AD-like phenotypes by using 5xFAD mice. (Lee et al., 2019) reported that *B. longum* NK46, which is isolated from human fecal matter, has anti-AD effects. Oral administration of *B. longum* NK46 (1 $\times$ 10<sup>9</sup>CFU/mouse/d) to 5xFAD mice (6 months old) for 1 and 2 months attenuated the hippocampal accumulation of A $\beta$  and cognitive decline. *B. bifidum* BGN4 and *B. longum* BORI were also reported to have anti-AD activities in 5xFAD mice. The administration of a probiotic mixture (*B. bifidum* BGN4 + *B. longum* BORI) (1 $\times$ 10<sup>9</sup>CFU/mouse/d) to 5xFAD mice (3 months old) for 30 d inhibited the loss of hippocampal neurons and cognitive decline estimated by the Y-maze test (Kim et al., 2021a). These reports suggest that 5xFAD, which exhibits AD-like physiology at a relatively early

stage, could be an excellent animal model for evaluating the anti-cognitive deficit activity of probiotics.

## **Conclusion**

Several animal models for cognitive impairment have been developed. If these animal models are properly used to estimate the anti-cognitive deficit activity of probiotics, more efficient and powerful probiotic-based anti-neurodegenerative disease agents could be developed. Additionally, these approaches will shed light on the detailed molecular mechanisms behind how probiotics modulate the gut–brain axis to attenuate the cognitive impairments that occur as a result of neurodegenerative diseases.

## **Conflicts of Interest**

The authors declare no potential conflicts of interest.

## **Acknowledgements**

This research was supported by an Incheon National University research grant (2018).

## **Author Contributions**

Conceptualization: Lee SH, Investigation: Kwon OY, Writing–original draft: Kwon OY, and Writing, review, and editing: Lee SH.

## **Ethics approval**

This article did not require approval from the IACUC/IRB since there were no animal and human participants.

**Table 1. Chemical-induced cognitive deficit animal models**

	Name	Doses/Methods	Major phenotypes	Refs
<b>Chemical-induced cognitive deficit animal models</b>	D-galactose-induced mice	IP injection of D-galactose (100–120 mg/kg bw/d) for 6–12 weeks.	Cognitive impairment Mitochondrial dysfunction Neuronal degeneration Apoptosis Depression and anxiety	(Liu et al., 2021; Parameshwaran et al., 2010; Woo et al., 2014)
	A $\beta$ -induced mice	ICV injection of A $\beta$ peptide solution (500–3000 $\mu$ g). A $\beta$ peptide aggregation occurs within 3–28 d.	Cognitive impairment A $\beta$ accumulation Neuronal loss	(Ali et al., 2015; Kobayashi et al., 2017; Kwon and Lee, 2020)
	Scopolamine-induced mice	IP injection of scopolamine (1–1.5 mg/kg bw/d) to mice could induce cognitive impairment within 1 h, but repetitive administration of scopolamine (once/d) is recommended during behavioral tests.	Cognitive impairment Neuronal degeneration Apoptosis	(Choi et al., 2021; Kim et al., 2021b; Yadang et al., 2020)
	LPS-induced mice	IP injection of LPS (0.25–1 mg/kg bw/d) in the abdominal cavity for 5–7 consecutive d.	Cognitive impairment Mitochondrial dysfunction Neuroinflammation	(Kamdi et al., 2021; Shoemark and Allen, 2015; Yang et al., 2020a)
		ICV injections of LPS (2–12 $\mu$ g). Cognitive deficiency due to neuroinflammation occurs 1–7 d after LPS injection.	Cognitive impairment Mitochondrial dysfunction Neuroinflammation	(Zhao et al., 2019; Zhou et al., 2006)

IP: Intraperitoneal; ICV: Intracerebroventricular; LPS: Lipopolysaccharides; A $\beta$ : beta-amyloid.

**Table 2. Transgenic mouse models that have cognitive deficit phenotypes**

	Name	Promoter	Mutation(s)	Symptoms	Refs
Senescence-accelerated mouse prone (SAMP)	SAMP8 mice	–	–	Age-associated increase in A $\beta$ deposition in hippocampal from 4 to 12 months ⇒ Astrogliosis, microgliosis, and cognitive deficits	(Takeda et al., 1981)
Single APP knock-in	<i>App</i> <sup>NL-G-F</sup> mice	Endogenous APP	<i>APP</i> <sup>KM670/671NL, E693G, I716F</sup>	A $\beta$ plaque deposition observed starting at 2 months and is nearly saturated by 7 months ⇒ Gliosis and cognitive deficits	(Guerreiro et al., 2010; Mullan et al., 1992; Nilsberth et al., 2001)
Double transgenic APP-Tg × PSEN1-Tg	APP/PS1 transgenic mice	Mouse Thy1 (APP, PS1)	<i>APP</i> <sup>KM670/671NL</sup> <i>PS1</i> <sup>H166P</sup>	A $\beta$ plaque deposition starting at approximately 6 weeks of age in the neocortex ⇒ Astrogliosis, microgliosis, formation of dystrophic neuritis, neuronal loss, and cognitive deficits	(Maia et al., 2013; Radde et al., 2006)
	5xFAD transgenic mice	Mouse Thy1.2 (APP, PS1)	<i>APP</i> <sup>KM670/671NL, I716V, V717I</sup> <i>PS1</i> <sup>M146L, L286V</sup>	Neuron loss occurs in multiple brain regions beginning at 6 months of age ⇒ Astrogliosis, microgliosis, neuronal loss, and cognitive deficits	(Devi and Ohno, 2010; Oakley et al., 2006)
Triple transgenic	3xTg-AD mice	Mouse Thy1.2 (APP) Endogenous (PS1)	<i>APP</i> <sup>K670N, M671L</sup> <i>PS1</i> <sup>M146V</sup> <i>MAPT</i> <sup>P301L</sup>	A $\beta$ deposition with intracellular immunoreactivity detected in some brain regions as early as 3–4 months of age ⇒ Gliosis and cognitive deficits	(Oddo et al., 2003)

Tg: Transgenic, Thy: Promotor of Thy-1 cell surface antigen gene, APP: Amyloid precursor protein, PS1: Presenilin-1, MAPT: Microtubule-associated protein tau, NFT: Neurofibrillary tangles.



**Table 3. Effects of probiotics on cognitive deficits in animal models**

Animal models	Strains (age, sex)	Proposed disease models	Probiotics (sources)	Doses	Effects	Behavior experiments	Refs
Chemical- induced cognitive deficit animal models	D- galactose induced mice	C57BL/6J (20-week- old males)	Alzheimer's disease	<i>Lactobacillus. pentosus</i> var. <i>plantarum</i> C29 (Kimchi)	$1 \times 10^{10}$ CFU/d/mouse for 5 weeks	Improvement of spatial memory and non-declarative connecting memory functions	Y-maze Morris water maze Passive avoidance (Woo et al., 2014)
		C57BL/6J (8-week-old males)	Alzheimer's disease	<i>L. paracasei</i> PS23 (Fermented milk)	$1 \times 10^{10}$ CFU/d/mouse for 9 weeks	Improvement of spatial memory	Morris water maze Open field (Cheng et al., 2022)
		ICR (males)	Alzheimer's disease	<i>Bacillus coagulans</i> JA845 (Fermented pickle)	$1 \times 10^9$ CFU/d/mouse for 10 weeks	Improvement of spatial memory	Morris water maze (Song et al., 2022)
	A $\beta$ induced mice	ddY mice (10-week- old males)	Alzheimer's disease	<i>Bifidobacterium breve</i> A1 (Yogurt)	$1 \times 10^9$ CFU/d/mouse for 10 d	Improvement of spatial memory and non-declarative connecting memory functions	Y-maze test Passive avoidance (Kobay ashi et al., 2017)
		C57BL/6J (8-week-old males)	Alzheimer's disease	<i>B. breve</i> NMG <i>B. breve</i> MY <i>B. breve</i> CCFM1025 <i>B. breve</i> XY <i>B. breve</i> WX (Human feces)	$0.6 \times 10^9$ CFU/d/mouse for 6 weeks	Improvement of spatial memory and non-declarative connecting memory functions	Y-maze Morris water maze Passive avoidance (Zhu et al., 2021)
	Scopola mine induced	Swiss albino (Females)	Alzheimer's disease	<i>L. rhamnosus</i> UBLR-58 (Kimchi)	$1 \times 10^9$ CFU/d/mouse for 10 d	Improvement of spatial memory	Morris water maze (Patel et al., 2020)

	mice		ICR (6-week-old males)	Alzheimer's disease	<i>L. johnsonii</i> CJLJ103 (Human feces)	$1 \times 10^9$ CFU/d/mouse for 5 d	Improvement of spatial memory and non-declarative connecting memory functions	Y-maze Passive avoidance	(Lee et al., 2018)
	LPS-induced mice	C57BL/6 (8-week-old males)	Alzheimer's disease	<i>Lactococcus lactis</i> subsp. <i>cremoris</i> LL95 (Mozzarella cheese)	$1 \times 10^9$ CFU/d/mouse for 7 d	Improvement of spatial memory	Sucrose preference Open field Forced swim	(Ramalho et al., 2022)	
		C57BL/6 (6-week-old males)	Alzheimer's disease	<i>B. longum</i> NK173 <i>L. plantarum</i> NK151 (Human feces)	$1 \times 10^9$ CFU/d/mouse for 5 d	Improvement of spatial memory	Y-maze, Novel object recognition	(Lee et al., 2021)	
		C57BL/6J, (8-week-old males)	Alzheimer's disease	<i>B. animalis</i> subsp. <i>lactis</i> BL03 <i>B. animalis</i> subsp. <i>lactis</i> BI04 <i>B. breve</i> BB02 <i>L. acidophilus</i> BA05 <i>L. helveticus</i> BD08 <i>L. paracasei</i> BP07 <i>L. plantarum</i> BP06 <i>Streptococcus thermophilus</i> BT01 (Human feces)	$1 \times 10^9$ CFU/d/mouse for 15 d	Improvement of spatial memory Finding space well has the effect of reducing anxiety symptoms	Open field Plus maze	(Petrella et al., 2021)	
Transgenic mouse models which have cognitive deficits phenotypes	SAMP8 mice	9-month-old (Male)	Alzheimer's disease	ProBiotic-4 ( <i>B. lactis</i> , <i>L. casei</i> , <i>B. bifidum</i> , <i>L. acidophilus</i> ) (Fermented milk)	$2 \times 10^9$ CFU/d/mouse for 12 weeks	Improvement of spatial memory and non-declarative connecting memory functions	Y-maze Passive avoidance	(Yang et al., 2020b)	
	<i>App<sup>NL-G-F</sup></i> mice	6–8 months old	Alzheimer's disease	VSL#3 ( <i>L. plantarum</i> , <i>L. delbrueckii</i> subsp.	$1.28 \times 10^9$ CFU/d/mouse	Finding space well has the effect of	Cross-maze rodent behavior	(Kaur et al.,	

(Female)		<i>bulgaricus</i> , <i>L. paracasei</i> , <i>L. acidophilus</i> , <i>B. breve</i> , <i>B. longum</i> , <i>B. infantis</i> , and <i>S. salivarius</i> sub) (Yogurt, fermented milk)	for 8 weeks	reducing anxiety symptoms		2020b)
6–8-month- old (Female)	Alzheimer's disease	VSL#3 ( <i>L. plantarum</i> , <i>L. delbrueckii</i> subsp. <i>bulgaricus</i> , <i>L. paracasei</i> , <i>L. acidophilus</i> , <i>B. breve</i> , <i>B. longum</i> , <i>B. infantis</i> , and <i>S. salivarius</i> sub) (Yogurt, fermented milk)	$1.28 \times 10^9$ CFU/d/mouse for 8 weeks	Improvement of non-declarative connecting memory functions	Passive avoidance	(Kaur et al., 2020a)
6-month- old (ND)	Alzheimer's disease	<i>Clostridium butyricum</i> WZMC1016 (Milk and cheeses)	$0.2 \times 10^9$ CFU/d/mouse for 4 weeks	Improvement of spatial memory	Morris water maze Object recognition	(Sun et al., 2020)
6-month- old (Male)	Alzheimer's disease	<i>L. plantarum</i> ATCC 8014 (Fermented milk)	$1 \times 10^9$ CFU/d/mouse for 12 weeks	Improvement of spatial memory and working memory	Spontaneous locomotor activity test Nest building, Novel object recognition Morris water maze Shuttlebox	(Wang et al., 2020)
4-month- old (ND)	Alzheimer's disease	<i>B. Lactis</i> Probio-M8 (Yogurt)	$2 \times 10^{10}$ CFU/d/mouse for 45 d	Improvement of spatial memory and working memory	Novel object recognition Y-maze	(Cao et al., 2021)

	8-week-old (Male)	Alzheimer's disease	SLAB51 ( <i>S. thermophilus</i> , <i>B. longum</i> , <i>B. breve</i> , <i>B. infantis</i> , <i>L. acidophilus</i> , <i>L. plantarum</i> , <i>L. paracasei</i> , <i>L. delbrueckii</i> subsp. <i>bulgaricus</i> , <i>L. brevis</i> [Yogurt])	$4 \times 10^9$ CFU/d/mouse for 4 months	Improvement of spatial memory and working memory and non-declarative connecting memory functions	Open field Novel-object recognition Passive avoidance	(Bonfili et al., 2017)
3xTg-AD mice	12-week-old (Male)	Alzheimer's disease	<i>L. salivarius</i> CUL61, <i>L. paracasei</i> CUL08, <i>B. bifidum</i> CUL20, <i>B. animalis</i> subsp. <i>lactis</i> CUL34 (Fermented milk)	$5 \times 10^8$ CFU/mouse/d for 12 weeks	Improvement of spatial memory and working memory	Novel object recognition Open field	(Webb et al., 2022)
	6-month-old (Male)	Alzheimer's disease	<i>B. longum</i> NK46 (Human feces)	$1 \times 10^9$ CFU/ d/mouse for 8 weeks	Improvement of spatial memory and non-declarative connecting memory functions	Y-maze Morris water maze Passive avoidance	(Lee et al., 2019)
5xFAD-Tg mice	3-month-old (ND)	Alzheimer's disease	<i>B. bifidum</i> BGN4 and <i>B. longum</i> BORI (Human feces)	$1 \times 10^9$ CFU/d/mouse for 30 d	Improvement of spatial memory	A contextual fear conditioning test Y-maze Morris water maze	(Kim et al., 2021a)

ND: Not determined.

## References

- Akter R, Chowdhury MaR, Rahman MH. 2021. Flavonoids and polyphenolic compounds as potential talented agents for the treatment of alzheimer's disease and their antioxidant activities. *Curr Pharm Des* 27:345-356.
- Ali T, Yoon GH, Shah SA, Lee HY, Kim MO. 2015. Osmotin attenuates amyloid beta-induced memory impairment, tau phosphorylation and neurodegeneration in the mouse hippocampus. *Sci Rep* 5:11708.
- Alladi S, Mekala S, Chadalawada SK, Jala S, Mridula R, Kaul S. 2011. Subtypes of dementia: A study from a memory clinic in india. *Dement Geriatr Cogn Disord* 32:32-38.
- Banks WA, Farr SA, Morley JE. 2002. Entry of blood-borne cytokines into the central nervous system: Effects on cognitive processes. *Neuroimmunomodulation* 10:319-327.
- Blasko I, Marx F, Steiner E, Hartmann T, Grubeck-Loebenstien B. 1999. Tnfalpha plus ifngamma induce the production of alzheimer beta-amyloid peptides and decrease the secretion of apps. *FASEB J* 13:63-68.
- Bonfili L, Cecarini V, Berardi S, Scarpona S, Suchodolski JS, Nasuti C, Fiorini D, Boarelli MC, Rossi G, Eleuteri AM. 2017. Microbiota modulation counteracts alzheimer's disease progression influencing neuronal proteolysis and gut hormones plasma levels. *Sci Rep* 7:2426.
- Bowler JV, Munoz DG, Merskey H, Hachinski V. 1998. Fallacies in the pathological confirmation of the diagnosis of alzheimer's disease. *J Neurol Neurosurg Psychiatry* 64:18-24.
- Briguglio M, Dell'osso B, Panzica G, Malgaroli A, Banfi G, Zanaboni Dina C, Galentino R, Porta M. 2018. Dietary neurotransmitters: A narrative review on current

- knowledge. *Nutrients* 10.
- Bruel-Jungerman E, Lucassen PJ, Francis F. 2011. Cholinergic influences on cortical development and adult neurogenesis. *Behav Brain Res* 221:379-388.
- Buxbaum JD, Oishi M, Chen HI, Pinkas-Kramarski R, Jaffe EA, Gandy SE, Greengard P. 1992. Cholinergic agonists and interleukin 1 regulate processing and secretion of the alzheimer beta/a4 amyloid protein precursor. *Proc Natl Acad Sci U S A* 89:10075-10078.
- Cao J, Amakye WK, Qi C, Liu X, Ma J, Ren J. 2021. *Bifidobacterium lactis* probio-m8 regulates gut microbiota to alleviate alzheimer's disease in the app/ps1 mouse model. *Eur J Nutr* 60:3757-3769.
- Cheng LH, Chou PY, Hou AT, Huang CL, Shiu WL, Wang S. 2022. *Lactobacillus paracasei* ps23 improves cognitive deficits via modulating the hippocampal gene expression and the gut microbiota in d-galactose-induced aging mice. *Food Funct* 13:5240-5251.
- Chiu MJ, Chen TF, Yip PK, Hua MS, Tang LY. 2006. Behavioral and psychologic symptoms in different types of dementia. *J Formos Med Assoc* 105:556-562.
- Choi JH, Lee EB, Jang HH, Cha YS, Park YS, Lee SH. 2021. *Allium hookeri* extracts improve scopolamine-induced cognitive impairment via activation of the cholinergic system and anti-neuroinflammation in mice. *Nutrients* 13.
- Christensen H, Mackinnon AJ, Korten AE, Jorm AF, Henderson AS, Jacomb P, Rodgers B. 1999. An analysis of diversity in the cognitive performance of elderly community dwellers: Individual differences in change scores as a function of age. *Psychol Aging* 14:365-379.
- Collins SM, Surette M, Bercik P. 2012. The interplay between the intestinal microbiota and the brain. *Nat Rev Microbiol* 10:735-742.
- Deng X, Zhao S, Liu X, Han L, Wang R, Hao H, Jiao Y, Han S, Bai C. 2020. *Polygala tenuifolia*: A source for anti-alzheimer's disease drugs. *Pharm Biol* 58:410-416.

- Devi L, Ohno M. 2010. Phospho-eif2alpha level is important for determining abilities of bace1 reduction to rescue cholinergic neurodegeneration and memory defects in 5xfad mice. *PLoS One* 5:e12974.
- Guerreiro RJ, Baquero M, Blesa R, Boada M, Bras JM, Bullido MJ, Calado A, Crook R, Ferreira C, Frank A, Gomez-Isla T, Hernandez I, Lleo A, Machado A, Martinez-Lage P, Masdeu J, Molina-Porcel L, Molinuevo JL, Pastor P, Perez-Tur J, Relvas R, Oliveira CR, Ribeiro MH, Rogaeva E, Sa A, Samaranch L, Sanchez-Valle R, Santana I, Tarraga L, Valdivieso F, Singleton A, Hardy J, Clarimon J. 2010. Genetic screening of alzheimer's disease genes in iberian and african samples yields novel mutations in presenilins and app. *Neurobiol Aging* 31:725-731.
- Kamdi SP, Raval A, Nakhate KT. 2021. Phloridzin attenuates lipopolysaccharide-induced cognitive impairment via antioxidant, anti-inflammatory and neuromodulatory activities. *Cytokine* 139:155408.
- Kaur H, Golovko S, Golovko MY, Singh S, Darland DC, Combs CK. 2020a. Effects of probiotic supplementation on short chain fatty acids in the appnl-g-f mouse model of alzheimer's disease. *J Alzheimers Dis* 76:1083-1102.
- Kaur H, Nagamoto-Combs K, Golovko S, Golovko MY, Klug MG, Combs CK. 2020b. Probiotics ameliorate intestinal pathophysiology in a mouse model of alzheimer's disease. *Neurobiol Aging* 92:114-134.
- Kim H, Kim S, Park SJ, Park G, Shin H, Park MS, Kim J. 2021a. Administration of bifidobacterium bifidum bgn4 and bifidobacterium longum bori improves cognitive and memory function in the mouse model of alzheimer's disease. *Front Aging Neurosci* 13:709091.
- Kim HY, Lee DK, Chung BR, Kim HV, Kim Y. 2016. Intracerebroventricular injection of amyloid-beta peptides in normal mice to acutely induce alzheimer-like cognitive deficits. *J Vis Exp*.
- Kim Y, Kim J, He M, Lee A, Cho E. 2021b. Apigenin ameliorates scopolamine-induced

- cognitive dysfunction and neuronal damage in mice. *Molecules* 26.
- Kobayashi Y, Sugahara H, Shimada K, Mitsuyama E, Kuhara T, Yasuoka A, Kondo T, Abe K, Xiao JZ. 2017. Therapeutic potential of bifidobacterium breve strain a1 for preventing cognitive impairment in alzheimer's disease. *Sci Rep* 7:13510.
- Konsman JP, Parnet P, Dantzer R. 2002. Cytokine-induced sickness behaviour: Mechanisms and implications. *Trends Neurosci* 25:154-159.
- Kumar A, Dogra S, Prakash A. 2009. Effect of carvedilol on behavioral, mitochondrial dysfunction, and oxidative damage against d-galactose induced senescence in mice. *Naunyn Schmiedebergs Arch Pharmacol* 380:431-441.
- Kwon OY, Lee SH. 2020. Ameliorating activity of ishige okamurae on the amyloid beta-induced cognitive deficits and neurotoxicity through regulating erk, p38 mapk, and jnk signaling in alzheimer's disease-like mice model. *Mol Nutr Food Res* 64:e1901220.
- Kwon OY, Lee SH. 2021. Ishige okamurae suppresses trimethyltin-induced neurodegeneration and glutamate-mediated excitotoxicity by regulating maps/nrf2/ho-1 antioxidant pathways. *Antioxidants (Basel)* 10.
- Lazareno S, Popham A, Birdsall NJ. 2000. Allosteric interactions of staurosporine and other indolocarbazoles with n-[methyl-(3)h]scopolamine and acetylcholine at muscarinic receptor subtypes: Identification of a second allosteric site. *Mol Pharmacol* 58:194-207.
- Lee DY, Shin YJ, Kim JK, Jang HM, Joo MK, Kim DH. 2021. Alleviation of cognitive impairment by gut microbiota lipopolysaccharide production-suppressing lactobacillus plantarum and bifidobacterium longum in mice. *Food Funct* 12:10750-10763.
- Lee HJ, Lee KE, Kim JK, Kim DH. 2019. Suppression of gut dysbiosis by bifidobacterium longum alleviates cognitive decline in 5xfad transgenic and aged mice. *Sci Rep* 9:11814.



- Lee HJ, Lim SM, Kim DH. 2018. *Lactobacillus johnsonii* cjlj103 attenuates scopolamine-induced memory impairment in mice by increasing bdnf expression and inhibiting nf-kappab activation. *J Microbiol Biotechnol* 28:1443-1446.
- Liu Y, Liu Y, Guo Y, Xu L, Wang H. 2021. Phlorizin exerts potent effects against aging induced by d-galactose in mice and pc12 cells. *Food Funct* 12:2148-2160.
- Logsdon AF, Erickson MA, Rhea EM, Salameh TS, Banks WA. 2018. Gut reactions: How the blood-brain barrier connects the microbiome and the brain. *Exp Biol Med (Maywood)* 243:159-165.
- Lok K, Zhao H, Shen H, Wang Z, Gao X, Zhao W, Yin M. 2013. Characterization of the app/ps1 mouse model of alzheimer's disease in senescence accelerated background. *Neurosci Lett* 557 Pt B:84-89.
- Lopez-Sobaler AM, Lorenzo Mora AM, Salas Gonzalez MF, Peral Suarez A, Aparicio A, Ortega RMF. 2021. [importance of choline in cognitive function]. *Nutr Hosp* 37:18-23.
- Lu J, Wu DM, Zheng YL, Hu B, Zhang ZF. 2010. Purple sweet potato color alleviates d-galactose-induced brain aging in old mice by promoting survival of neurons via pi3k pathway and inhibiting cytochrome c-mediated apoptosis. *Brain Pathol* 20:598-612.
- Macintyre A, Abramov R, Hammond CJ, Hudson AP, Arking EJ, Little CS, Appelt DM, Balin BJ. 2003. *Chlamydia pneumoniae* infection promotes the transmigration of monocytes through human brain endothelial cells. *J Neurosci Res* 71:740-750.
- Maia LF, Kaeser SA, Reichwald J, Hruscha M, Martus P, Staufenbiel M, Jucker M. 2013. Changes in amyloid-beta and tau in the cerebrospinal fluid of transgenic mice overexpressing amyloid precursor protein. *Sci Transl Med* 5:194re192.
- Mehla J, Lacoursiere SG, Lapointe V, Mcnaughton BL, Sutherland RJ, Mcdonald RJ, Mohajerani MH. 2019. Age-dependent behavioral and biochemical

- characterization of single app knock-in mouse (app(nl-g-f/nl-g-f)) model of alzheimer's disease. *Neurobiol Aging* 75:25-37.
- Miyamoto M, Kiyota Y, Nishiyama M, Nagaoka A. 1992. Senescence-accelerated mouse (sam): Age-related reduced anxiety-like behavior in the sam-p/8 strain. *Physiol Behav* 51:979-985.
- Miyamoto M, Kiyota Y, Yamazaki N, Nagaoka A, Matsuo T, Nagawa Y, Takeda T. 1986. Age-related changes in learning and memory in the senescence-accelerated mouse (sam). *Physiol Behav* 38:399-406.
- Mohammadi G, Dargahi L, Peymani A, Mirzanejad Y, Alizadeh SA, Naserpour T, Nassiri-Asl M. 2019. The effects of probiotic formulation pretreatment (lactobacillus helveticus r0052 and bifidobacterium longum r0175) on a lipopolysaccharide rat model. *J Am Coll Nutr* 38:209-217.
- Morais LH, Schreiber HLT, Mazmanian SK. 2021. The gut microbiota-brain axis in behaviour and brain disorders. *Nat Rev Microbiol* 19:241-255.
- Morley JE, Armbrrecht HJ, Farr SA, Kumar VB. 2012. The senescence accelerated mouse (somp8) as a model for oxidative stress and alzheimer's disease. *Biochim Biophys Acta* 1822:650-656.
- Mullan M, Crawford F, Axelman K, Houlden H, Lilius L, Winblad B, Lannfelt L. 1992. A pathogenic mutation for probable alzheimer's disease in the app gene at the n-terminus of beta-amyloid. *Nat Genet* 1:345-347.
- Nilsberth C, Westlind-Danielsson A, Eckman CB, Condron MM, Axelman K, Forsell C, Stenh C, Luthman J, Teplow DB, Younkin SG, Naslund J, Lannfelt L. 2001. The 'arctic' app mutation (e693g) causes alzheimer's disease by enhanced abeta protofibril formation. *Nat Neurosci* 4:887-893.
- Oakley H, Cole SL, Logan S, Maus E, Shao P, Craft J, Guillozet-Bongaarts A, Ohno M, Disterhoft J, Van Eldik L, Berry R, Vassar R. 2006. Intraneuronal beta-amyloid aggregates, neurodegeneration, and neuron loss in transgenic mice with five

- familial alzheimer's disease mutations: Potential factors in amyloid plaque formation. *J Neurosci* 26:10129-10140.
- Oddo S, Caccamo A, Shepherd JD, Murphy MP, Golde TE, Kaye R, Metherate R, Mattson MP, Akbari Y, Laferla FM. 2003. Triple-transgenic model of alzheimer's disease with plaques and tangles: Intracellular abeta and synaptic dysfunction. *Neuron* 39:409-421.
- Panza F, Lozupone M, Solfrizzi V, Watling M, Imbimbo BP. 2019. Time to test antibacterial therapy in alzheimer's disease. *Brain* 142:2905-2929.
- Parameshwaran K, Irwin MH, Steliou K, Pinkert CA. 2010. D-galactose effectiveness in modeling aging and therapeutic antioxidant treatment in mice. *Rejuvenation Res* 13:729-735.
- Patel C, Pande S, Acharya S. 2020. Potentiation of anti-alzheimer activity of curcumin by probiotic *Lactobacillus rhamnosus* ublr-58 against scopolamine-induced memory impairment in mice. *Naunyn Schmiedeberg's Arch Pharmacol* 393:1955-1962.
- Perry VH. 2004. The influence of systemic inflammation on inflammation in the brain: Implications for chronic neurodegenerative disease. *Brain Behav Immun* 18:407-413.
- Petrella C, Strimpakos G, Torcinaro A, Middei S, Ricci V, Gargari G, Mora D, De Santa F, Farioli-Vecchioli S. 2021. Proneurogenic and neuroprotective effect of a multi strain probiotic mixture in a mouse model of acute inflammation: Involvement of the gut-brain axis. *Pharmacol Res* 172:105795.
- Prakash A, Kumar A. 2013. Pioglitazone alleviates the mitochondrial apoptotic pathway and mito-oxidative damage in the d-galactose-induced mouse model. *Clin Exp Pharmacol Physiol* 40:644-651.
- Radde R, Bolmont T, Kaeser SA, Coomaraswamy J, Lindau D, Stoltze L, Calhoun ME, Jaggi F, Wolburg H, Gengler S, Haass C, Ghetti B, Czech C, Holscher C,

- Mathews PM, Jucker M. 2006. Abeta42-driven cerebral amyloidosis in transgenic mice reveals early and robust pathology. *EMBO Rep* 7:940-946.
- Ramalho JB, Spiazzi CC, Bicca DF, Rodrigues JF, Sehn CP, Da Silva WP, Cibir FWS. 2022. Beneficial effects of *Lactococcus lactis* subsp. *Cremoris* 1195 treatment in an lps-induced depression-like model in mice. *Behav Brain Res* 426:113847.
- Ravi SK, Narasingappa RB, Vincent B. 2019. Neuro-nutrients as anti-alzheimer's disease agents: A critical review. *Crit Rev Food Sci Nutr* 59:2999-3018.
- Rehman SU, Shah SA, Ali T, Chung JI, Kim MO. 2017. Anthocyanins reversed d-galactose-induced oxidative stress and neuroinflammation mediated cognitive impairment in adult rats. *Mol Neurobiol* 54:255-271.
- Saito T, Matsuba Y, Mihira N, Takano J, Nilsson P, Itohara S, Iwata N, Saido TC. 2014. Single app knock-in mouse models of alzheimer's disease. *Nat Neurosci* 17:661-663.
- Salminen S, Bouley C, Boutron-Ruault MC, Cummings JH, Franck A, Gibson GR, Isolauri E, Moreau MC, Roberfroid M, Rowland I. 1998. Functional food science and gastrointestinal physiology and function. *Br J Nutr* 80 Suppl 1:S147-171.
- Shoemark DK, Allen SJ. 2015. The microbiome and disease: Reviewing the links between the oral microbiome, aging, and alzheimer's disease. *J Alzheimers Dis* 43:725-738.
- Shwe T, Pratchayasakul W, Chattipakorn N, Chattipakorn SC. 2018. Role of d-galactose-induced brain aging and its potential used for therapeutic interventions. *Exp Gerontol* 101:13-36.
- Song X, Zhao Z, Zhao Y, Jin Q, Li S. 2022. Protective effects of *Bacillus coagulans* ja845 against d-galactose/alcl3-induced cognitive decline, oxidative stress and neuroinflammation. *J Microbiol Biotechnol* 32:212-219.
- Stover KR, Campbell MA, Van Winssen CM, Brown RE. 2015. Early detection of cognitive deficits in the 3xtg-ad mouse model of alzheimer's disease. *Behav*

Brain Res 289:29-38.

- Sun J, Xu J, Yang B, Chen K, Kong Y, Fang N, Gong T, Wang F, Ling Z, Liu J. 2020. Effect of clostridium butyricum against microglia-mediated neuroinflammation in alzheimer's disease via regulating gut microbiota and metabolites butyrate. *Mol Nutr Food Res* 64:e1900636.
- Takeda T, Hosokawa M, Takeshita S, Irino M, Higuchi K, Matsushita T, Tomita Y, Yasuhira K, Hamamoto H, Shimizu K, Ishii M, Yamamuro T. 1981. A new murine model of accelerated senescence. *Mech Ageing Dev* 17:183-194.
- Van Erum J, Van Dam D, De Deyn PP. 2019. Alzheimer's disease: Neurotransmitters of the sleep-wake cycle. *Neurosci Biobehav Rev* 105:72-80.
- Vitetta L, Briskey D, Hayes E, Shing C, Peake J. 2012. A review of the pharmacobiotic regulation of gastrointestinal inflammation by probiotics, commensal bacteria and prebiotics. *Inflammopharmacology* 20:251-266.
- Vitetta L, Vitetta G, Hall S. 2018. Immunological tolerance and function: Associations between intestinal bacteria, probiotics, prebiotics, and phages. *Front Immunol* 9:2240.
- Wang H, Kulas JA, Wang C, Holtzman DM, Ferris HA, Hansen SB. 2021. Regulation of beta-amyloid production in neurons by astrocyte-derived cholesterol. *Proc Natl Acad Sci U S A* 118.
- Wang QJ, Shen YE, Wang X, Fu S, Zhang X, Zhang YN, Wang RT. 2020. Concomitant memantine and lactobacillus plantarum treatment attenuates cognitive impairments in app/ps1 mice. *Aging (Albany NY)* 12:628-649.
- Webberley TS, Masetti G, Bevan RJ, Kerry-Smith J, Jack AA, Michael DR, Thomas S, Glymenaki M, Li J, McDonald JA, John D, Morgan JE, Marchesi JR, Good MA, Plummer SF, Hughes TR. 2022. The impact of probiotic supplementation on cognitive, pathological and metabolic markers in a transgenic mouse model of alzheimer's disease. *Front Neurosci* 16:843105.

- Whitehouse PJ, Price DL, Clark AW, Coyle JT, DeLong MR. 1981. Alzheimer disease: Evidence for selective loss of cholinergic neurons in the nucleus basalis. *Ann Neurol* 10:122-126.
- Whitehouse PJ, Price DL, Struble RG, Clark AW, Coyle JT, DeLong MR. 1982. Alzheimer's disease and senile dementia: Loss of neurons in the basal forebrain. *Science* 215:1237-1239.
- Woo JY, Gu W, Kim KA, Jang SE, Han MJ, Kim DH. 2014. *Lactobacillus pentosus* var. *Plantarum* c29 ameliorates memory impairment and inflammaging in a d-galactose-induced accelerated aging mouse model. *Anaerobe* 27:22-26.
- Yadang FSA, Nguézeye Y, Kom CW, Betote PHD, Mamat A, Tchokouaha LRY, Taiwe GS, Agbor GA, Bum EN. 2020. Scopolamine-induced memory impairment in mice: Neuroprotective effects of *Carissa edulis* (forssk.) valh (apocynaceae) aqueous extract. *Int J Alzheimers Dis* 2020:6372059.
- Yang L, Zhou R, Tong Y, Chen P, Shen Y, Miao S, Liu X. 2020a. Neuroprotection by dihydrotestosterone in lps-induced neuroinflammation. *Neurobiol Dis* 140:104814.
- Yang X, Yu D, Xue L, Li H, Du J. 2020b. Probiotics modulate the microbiota-gut-brain axis and improve memory deficits in aged samp8 mice. *Acta Pharm Sin B* 10:475-487.
- Zhao J, Bi W, Xiao S, Lan X, Cheng X, Zhang J, Lu D, Wei W, Wang Y, Li H, Fu Y, Zhu L. 2019. Neuroinflammation induced by lipopolysaccharide causes cognitive impairment in mice. *Sci Rep* 9:5790.
- Zhou H, Lapointe BM, Clark SR, Zbytnuik L, Kubes P. 2006. A requirement for microglial tlr4 in leukocyte recruitment into brain in response to lipopolysaccharide. *J Immunol* 177:8103-8110.
- Zhu G, Zhao J, Zhang H, Chen W, Wang G. 2021. Administration of *Bifidobacterium breve* improves the brain function of Aβ1-42-treated mice via the modulation

of the gut microbiome. Nutrients 13.

ACCEPTED