



ARTICLE

Oxidative Stability and Quality Characteristics of Duck, Chicken, Swine and Bovine Skin Fats Extracted by Pressurized Hot Water Extraction

Dong-Min Shin¹, Do Hyun Kim¹, Jong Hyeok Yune¹, Hyuk Cheol Kwon¹,
Hyo Juong Kim², Han Geuk Seo¹, and Sung Gu Han^{1,*}

¹Department of Food Science and Biotechnology of Animal Resources, Konkuk University, Seoul 05029, Korea

²Taekyung Food and Processing R&D Center, Seoul 07057, Korea



Received April 4, 2019

Revised April 24, 2019

Accepted April 30, 2019

*Corresponding author : Sung Gu Han
Department of Food Science and
Biotechnology of Animal Resources,
Konkuk University, Seoul 05029, Korea
Tel: +82-2-450-0526
Fax: +82-2-455-1044
E-mail: hansg@konkuk.ac.kr

*ORCID

Dong-Min Shin
<https://orcid.org/0000-0003-2755-433X>
Do Hyun Kim
<https://orcid.org/0000-0002-2500-8688>
Jong Hyuk Yune
<https://orcid.org/0000-0002-3015-7661>
Hyuk Cheol Kwon
<https://orcid.org/0000-0001-6234-2530>
Hyo Juong Kim
<https://orcid.org/0000-0003-2819-0822>
Han Geuk Seo
<https://orcid.org/0000-0002-9123-3816>
Sung Gu Han
<https://orcid.org/0000-0002-1485-861X>

Abstract The aim of this study was to investigate the oxidative status and quality characteristics of four animal skin-derived fats extracted using an identical extraction method. Pressurized hot water extraction, a green extraction method, was used to extract animal skin fats (duck, chicken, swine, and bovine skin). Multiple experiments were performed during accelerated storage at 60°C for 90 days. Quality characteristics, such as extraction yield, iodine value (IV), fatty acid composition, and fat viscosity were determined. In addition, indicators for oxidative status, including acid value (AV), peroxide value (PV), *p*-anisidine value (*p*-AV), thiobarbituric acid reactive substances (TBARS), conjugated dienes (CD), and total oxidation (totox) values were evaluated. The fat extraction yield was highest in bovine fat, followed by duck, swine, and chicken fats. The IV was higher in duck and chicken fats. Duck fats contained the most unsaturated fats and the least saturated fats. Fat oxidation indicators, such as PV, TBARS, and totox values, were relatively higher in duck fats during storage compared to the other fats. Other indicators, including AV, *p*-AV, and CD, were similar in duck, chicken, and swine fats. Viscosity was similar in all the tested fats but markedly increased after 70 days of storage in duck fats. Our data indicate that duck skin fat was more vulnerable to oxidative changes in accelerated storage conditions and this may be due to its higher unsaturated fatty acid content. Supplementation with antioxidants might be a reasonable way to solve the oxidation issue in duck skin fats.

Keywords duck skin, edible fat, oxidative stability, unsaturated fatty acid, fat oxidation

Introduction

Dietary fat is a major nutritional and biochemical component of the human body necessary for maintaining physiological, biochemical, and molecular mechanisms and the composition of cell components and for providing a source of energy. As a dietary component, fat plays a major role in taste, flavor, and aroma in most foods. In contrast, excessive fat consumption has been associated with the occurrence of chronic human

diseases, such as obesity, cardiovascular disease, diabetes, and cancer (Hariri and Thibault, 2010). Thus, efforts to reduce total fat consumption have continued for decades. However, according to recent reports from the National Health and Nutrition Examination Survey, the incidence of chronic diseases has remained high, despite reductions in fat intake (Austin et al., 2011). Therefore, recent studies have recommended that people should consider improving the quality of fat consumed, instead of reducing total fat intake, because the balance of health benefits to risks is determined by the amount of saturated fatty acids (SFA) and unsaturated fatty acids (USFA) consumed (Moro and Capel, 2019). Current recommendations limit SFA consumption to less than 10% of total calories due to the risk of cardiovascular disease (USDA and HHS, 2015). In contrast, USFA consumption showed positive health effects through decreased low-density lipoprotein cholesterol levels and reduced coronary heart disease (Li and Sun, 2019). Thus, the consumption of unsaturated fats (e.g., vegetable and fish oil) has been recommended, instead of saturated fats (e.g., lard and tallow), to reduce adverse health effects (Moghtadaei et al., 2018).

Duck meat is a type of poultry meat with a unique flavor. It also contains high levels of essential amino acids and polyunsaturated fatty acids (Jo et al., 2018). Particularly, a higher amount of long-chain fatty acids, such as oleic acid (C18:1) and linoleic acid (C18:2) in duck skin and tissue, have been reported (Chen et al., 2017; Heo et al., 2013). In fact, these fatty acids have been known to prevent cardiovascular disease, including arteriosclerosis and cholesterol and triglyceride accumulation in blood vessels. It was estimated that the worldwide duck meat consumption is approximately 4.53 million metric tons (MMT) per annum, while poultry meat accounts for 121 MMT per annum (FAOSTAT, 2018; Khan et al., 2019). The increased production and consumption of duck meat has resulted in large amounts of by-products, such as duck skin. Duck skin fat has the potential for use as a raw material in the production of healthy animal-derived fats (Huda et al., 2013).

Pressurized hot water extraction (PHWE), a process which combines pressure and high water temperature, is a green extraction method for foods and herbal plants (Chemat et al., 2012). The advantage of PHWE is that the degradation of minor components is minimized because of the short extraction time (dos Santos Freitas et al., 2008). More importantly, PHWE utilizes hot water, whereas traditional extraction methods, such as pressurized liquid extraction requires non-environmentally friendly organic solvents (Kronholm et al., 2007). Studies on the quality properties of vegetables and fruit oil prepared by PHWE have been published (Mustafa and Turner, 2011). PHWE-employed fat extraction from animal skin has not been frequently used but its advantages to protect minor components in the fats and protect the environment warrant further investigation.

Therefore, in the current study, the oxidative stability and quality characteristics of duck skin fat were evaluated in comparison with those of other animal fats (chicken, swine, and bovine skin fats). An identical fat extraction method (i.e., PHWE) was used for all animal skins. This study was also performed to provide information on the potential use of duck skin fats as a healthy edible animal fat in the food industry and for cooking.

Material and Methods

Chemicals

Potassium iodide, *p*-anisidine, potassium hydroxide, heptadecanoic acid methyl ester, starch, thiobarbituric acid, and phenolphthalein were purchased from Sigma Aldrich Chemical Co. (St. Louis, MO, USA). Chloroform, isooctane, glacial acetic acid, methanol, sodium thiosulfate (0.1 N and 0.01 N), ethanol, and ether were purchased from Daejung Co. (Siheung, Korea). Benzene and iodine monochloride were purchased from Samchun (Seoul, Korea). All chemicals were analytical grade and used without further purification.

Fat extraction from animal skin

Animal skins (duck, chicken, swine, and bovine skin) were obtained Farm Duck Co. Ltd., Jeongeup-si, Korea). The PHWE method has previously been reported for animal fat extraction (Plaza and Turner, 2015). In detail, animal skins (2 kg) were washed several times and the visible fat and connective tissue were removed from the skin. Fat extraction was performed using a pressure extractor (Yu-il ENT Co. Ltd, Seoul, Korea) at 115°C, 1.4 kgf/cm² for 3 h and dehydration for 30 min. The samples were filtered using a 40-mesh bucket filter, then the filtrate was centrifuged using a decanter (GEA Westfalia Separator, Oelde, Germany) at 7,500 rpm for one hour. The fat phase was collected from the supernatant and re-centrifuged at 15,000 rpm for one hour. The extract was packed in a plastic container and stored at -80°C until use.

Oven storage stability test

A modified Schaal oven test (accelerated storage test) was used to measure the oxidative stability of fat at 60°C (Warner et al., 1989). Briefly, the extracted fat samples (1 kg) were poured into glass beakers and placed in a drying oven (SW-90D, Sangwoo Scientific Co., Korea) which was set to 60±2°C for 90 days. Experimental analyses were performed within a 10-day interval.

Extract yield

The extract yield was determined by calculating the difference in the weights of the samples before and after processing:

$$\text{Extract yield (\%)} = (\text{Weight of sample after extraction} / \text{Weight of raw material}) \times 100$$

Fatty acid composition

The fatty acid composition was determined using the method reported previously by Muguerza et al. (2001). Briefly, the fatty acid composition was determined using gas chromatography (5890, Agilent Technologies, Santa Clara, USA) with a capillary column SP-2560 (100 m×0.25 mm×0.2 µm). Chromatographic conditions were as follows: the temperature of both the injector and detector was 225°C, the oven temperature was programmed to increase from 100°C to 240°C at a rate of 1°C/min. Hydrogen was the carrier gas at a flow rate of 1 mL/min. The quantification of individual fatty acids was determined by using heptadecanoic acid methyl ester as an internal standard.

Iodine value (IV)

The iodine value was measured as described previously (Kyriakidis and Katsiloulis, 2000). The samples (0.2–0.4 g) were dissolved in 20 mL of chloroform and 25 mL of iodine monochloride solution in a flask. The mixture was kept for 30 min in the dark at room temperature, then 20 mL of 1 N KI solution and 100 mL of distilled water was added. Then, the mixture was titrated against 0.1 N sodium thiosulphate solution with 1% starch solution as an indicator. IV was calculated with the following equation:

$$\text{IV (g/100 g)} = 1.269 \times (b - a) / W$$

Where *a* was the volume (mL) of 0.1 N sodium thiosulfate consumed in the titration, *b* was the normality of the 0.1 N

sodium thiosulfate, and W was the weight of the sample (g).

Acid value (AV)

Free fatty acid content was determined by a standard titrimetry method (Ghadge and Raheman, 2005). The samples (5 g) were mixed with 100 mL of ethanol/ether solution (1:2, v/v) and the mixture was titrated against 0.1 N potassium hydroxide with 1% phenolphthalein as an indicator. Acid value (AV) was calculated with the following equation:

$$AV \text{ (mg KOH/g)} = 5.611 \times (a - b)/W$$

Where a was the volume (mL) of potassium hydroxide consumed in the titration, b was the normality of the potassium hydroxide and W was the weight of the sample (g).

Peroxide value (PV)

The PV of the samples was determined by the AOAC (2007) method. The samples were dissolved with 10 mL of chloroform and 15 mL of glacial acetic acid in a flask. Then, 1 mL of saturated solution of KI was added to the flask and kept for 10 min in the dark. Samples added with 30 mL of distilled water were thoroughly mixed and the mixture was titrated against 0.01 N sodium thiosulphate solution with 1% starch solution as an indicator. The PV was calculated as follows:

$$PV \text{ (meq/kg)} = (a - b) \times f/W \times 10$$

Where a was the volume (mL) of sodium thiosulfate consumed in the titration, b was the normality of the sodium thiosulfate, and W was the weight of the sample (g).

p-Anisidine value (*p*-AV)

Determination of the *p*-AVs was carried out as previously described (Hashempour-Baltork et al., 2017). Samples (1 g) were dissolved in isooctane in a 25 mL volumetric flask and mixed with 1 mL *p*-anisidine solution (0.25% w/v in 99.5% glacial acetic acid). The mixture was reacted for 10 min in the dark. The absorbance was measured at 350 nm using a UV/VIS spectrophotometer (Optizen 2120 UV Plus, Mecasys Co., Ltd., Daejeon, Korea) and the *p*-AV was calculated as follows:

$$p\text{-AV} = 25 \times (1.2A_s - A_b)/W$$

Where, A_s was the absorbance of the sample with *p*-anisidine solution, A_b was the absorbance of the sample without *p*-anisidine solution, and W was the weight of the sample (g).

Total oxidation (TOTOX) value

The total oxidation state (TOTOX value) of the oils and fats was calculated according to the formula:

$$TOTOX = p\text{-AV} + 2PV$$

Thiobarbituric acid reactive substances (TBARS)

The thiobarbituric acid reactive substances (TBARS) value was obtained following a procedure described previously (Papastergiadis et al., 2012). A 1 g sample was weighed into a 50 mL conical tube and mixed with 5 mL distilled water for two minutes. The mixture was centrifuged at 5,000 g for five minutes. The aqueous layer was collected, and the procedure was repeated twice. The collected sample (2.5 mL) and 2.5 mL of TBA reagent (46 mM in 99.5% glacial acetic acid) were mixed in a test tube and heated in a water bath at 95°C for 35 minutes. After cooling, the absorbance was measured at 532 nm using a UV/VIS spectrophotometer. The results were expressed as mg MDA/kg.

Conjugated dienes (CD)

The conjugated dienes (CDs) were determined using a UV/VIS at 233 nm. Before analysis, 100 mg of sample was diluted with isooctane in a 25 mL of volumetric flask. The CD expressed as a percentage of conjugated dienoic acid was calculated as follows:

$$\text{CD (\%)} = 0.84 \times \text{Ab}_{233} / (bc - K_0)$$

Where Ab_{233} was the absorbance observed at 233 nm, b was the cell length (cm), c was the concentration of the sample (g/L), and K_0 was the absorptivity of the acid (value of 0.03).

Viscosity

The viscosity of the samples was analyzed using a DV-E viscometer (Brookfield, Toronto, Canada) during the storage period. Thirty mL of sample was added to a conical tube and the analysis was conducted at a constant rotational speed of 50 rpm using the No. 62 spindle. The measurement was repeated five times for 30 s and the results were expressed as centipoise (cP) units.

Statistical analyses

Statistical analysis was performed with two-way analysis of variance (ANOVA). The ANOVA was performed on all the variables using the General Linear Model (GLM) procedure of SPSS Ver. 24.0 (SPSS INC., USA). Duncan's multiple range test ($p < 0.05$) was used to determine significant differences between the sample groups.

Results and Discussion

Extraction yield and iodine value

The extraction yields of animal skin-derived fats using PHWE are shown in Table 1. The highest ($p < 0.05$) fat extraction (60.73%) was obtained from bovine skin, followed by duck skin (34.05%), swine skin (23.20%), and chicken skin (14.52%). Bovine skin products contained the highest amount of fats, whereas chicken skin contained the lowest amount of fats. The IV represents the degree of unsaturated fats and oils and also indicates the relative content of USFA (Naz et al., 2005). The IV determines the quality and grade of oil and is used as an authentication test in the oil industry (Yan et al., 2018). The IV of animal skin fat in the present study is shown in Table 1. The IV of the samples ranged from 55.71 to 79.57 g $\text{I}_2/100$ g. The IV

Table 1. Quality characteristics of the animal skin (duck, chicken, swine, and bovine) fats

Parameters	Animal skin fats			
	Duck fat	Chicken fat	Swine fat	Bovine fat
Extraction yield (%)	34.05±1.22 ^b	14.52±0.88 ^d	23.20±1.44 ^c	60.73±2.25 ^a
Iodine value (g/100 g)	77.57±1.26 ^a	77.32±1.69 ^a	71.59±0.36 ^b	55.71±2.04 ^c

All values are the mean±SD of three replicates.

^{a-d} means within a row with different letters are significantly different ($p < 0.05$).

of duck and chicken fats was 77.57 and 77.32 g I₂/100 g, respectively. Fats containing high IVs are chemically unstable because the double bonds in USFA are relatively reactive and vulnerable to oxidation (Encinar et al., 2019). Duck fat has been characterized by a high ratio of USFA. The major fatty acids identified were palmitic acid (C16:0), stearic acid (18:0), oleic acid (C18:1), linoleic acid (C18:2), and arachidonic acid (C20:4) (Cobos et al., 2000). The higher IV duck skin fat observed in this study reflected its higher USFA contents.

Fatty acid composition

The fatty acid composition of fat is a critical indicator of nutritional value. The fatty acid composition is associated with oxidative stability and thus, high levels of polyunsaturated fatty acids can contribute to the rapid deterioration of fats (Symoniuk et al., 2019). The fatty acid composition of the animal skin fats is summarized in Table 2. The fatty acid compositions were significantly different based on the origin of the animal skin fat ($p < 0.05$). For most skin fats, oleic acid (C18:1) was the major USFA, followed by palmitic acid (C16:0), linoleic acid (C18:2), and stearic acid (C18:0). Only bovine skin fat showed more stearic acid (C18:0) than linoleic acid (C18:2). The fatty acid composition measured in the skin of animals were similar to previously reported data (da Silva et al., 2019; Kang et al., 2014; Marion and Woodroof, 1963; Park

Table 2. Fatty acid composition of animal skin (duck, chicken, swine, and bovine) fats

FA (%)	Animal skin fats			
	Duck fat	Chicken fat	Swine fat	Bovine fat
Lauric 12:0	0.04±0.01 ^c	ND	0.11±0.01 ^b	0.17±0.01 ^a
Myristic 14:0	0.68±0.03 ^c	0.83±0.02 ^b	0.42±0.01 ^d	4.69±0.02 ^a
Palmitic 16:0	22.57±0.74 ^c	23.20±0.52 ^b	24.06±0.92 ^b	26.65±0.01 ^a
Stearic 18:0	5.24±0.33 ^c	5.67±0.06 ^c	11.81±0.61 ^a	7.24±0.01 ^b
Oleic 18:1	48.70±0.14 ^a	44.64±0.54 ^b	41.87±0.25 ^c	43.49±0.41 ^b
Linoleic 18:2	15.08±0.91 ^a	14.31±0.43 ^a	12.99±0.46 ^b	1.93±0.05 ^c
α -Linolenic 18:3	0.73±0.03 ^b	0.79±0.01 ^a	0.62±0.01 ^c	0.11±0.01 ^d
Σ SFA ¹⁾	28.53±1.06 ^d	29.70±0.57 ^c	36.40±0.31 ^b	38.75±0.03 ^a
Σ USFA	64.51±0.61 ^a	59.74±0.82 ^b	55.48±0.36 ^c	45.53±0.42 ^d
Σ USFA/SFA	2.26±0.08 ^a	2.01±0.02 ^b	1.52±0.01 ^c	1.17±0.01 ^d
Total FA	93.04±1.44 ^a	89.44±1.34 ^b	91.89±0.67 ^a	84.28±0.44 ^c

All values are the mean±SD of three replicates.

^{a-d} Means within a row with different letters are significantly different ($p < 0.05$).

¹⁾ Σ SFA: saturated fatty acid=C12:0+C14:0+C16:0+C18:0; Σ USFA: unsaturated fatty acid=C18:1+C18:2+C18:3; Σ USFA/SFA: ratio of unsaturated fatty acid to saturated fatty acid; FA, fatty acid; ND, not detected.

et al., 2014). In the current study, duck skin fat displayed a higher content of oleic acid ($p < 0.05$) than other skin fats. Oleic acid plays an important role in human nutrition (Rekas et al., 2015). Particularly, oleic acid is an USFA that provides biologically beneficial effects in the human body. A previous report demonstrated that the consumption of canola oils enriched with high oleic acid decreased atherogenic lipids and lipoproteins compared to commercial canola oils (Bowen et al., 2019). Thus, high levels of oleic acid in duck skin fat may confer extra value as a food ingredient. In addition, a few reports support the association between oleic acid and flavor. For example, high oleic acid composition provided flavor stability and improved flavor of peanuts (Mugendi et al., 1998; Talcott et al., 2005). The total SFA and USFA contents were significantly different between the groups ($p < 0.05$). Duck skin fat showed the highest total USFA content and the lowest total SFA content, while bovine skin fat exhibited the lowest total USFA and the highest SFA content. More importantly, the ratio of USFA/SFA of duck skin fat was approximately 2-fold higher than that of bovine skin fat ($p < 0.05$). The consumption of unsaturated fat has been linked to the prevention of cardiovascular diseases, such as atherosclerosis (da Silva et al., 2019). Our fatty acid composition analysis demonstrated that duck skin fat contained higher levels of oleic acid and a high USFA/SFA ratio and these might be useful indicators for the increased use of duck skin fat in cooking and food industries.

Comparisons of oxidation stability of animal skin fats

Quality characteristics of oil and fat are affected by multiple factors, including fatty acid composition, extraction methods, thermal treatment, and the presence of transition metals, pigments, and antioxidants (Choe and Min, 2006). In particular, high levels of polyunsaturated fatty acids may result in the rapid oxidation of fats (Resende et al., 2019). Thus, the oxidative stability of animal skin fats was evaluated during the 90-day storage period at 60°C (Figs. 1A–F). Results of the AV analysis are shown in Fig. 1A. The AV is generated by the hydrolysis and oxidation of triacylglycerol (Ghobadi et al., 2018). The initial AVs (day 0) ranged from 0.64 to 1.81 mg/g fat and the levels were markedly increased and ranged from 3.43 to 6.21 mg/g after 90 days of storage at 60°C. The initial AVs were relatively lower in duck and chicken skin fats than in swine and bovine skin fats. During 90 days of storage, the AVs were increased in all animal skin fats. Specifically, the values were rapidly increased after day 60, probably due to increased hydrolysis and oxidation of triacylglycerol. Increases in AVs also occur in other foods, such as peanuts, during storage. AV levels were increased in raw peanuts stored at room temperature for 720 days (Martin et al., 2018). At the end of the storage period in this study, the AV levels were higher in duck and chicken skin fats than in swine and bovine skin fats. This may have been due to higher concentrations of USFA in the duck and chicken skin fats. USFAs are more susceptible to oxidation and hydrolysis than SFAs (Zhou et al., 2019).

The PV represents the peroxides (primary oxidation products) produced by the oxidation process in fats. Primary oxidation products are broken down into minor substances, such as low molecular weight aldehydes, ketones, alcohols and short-chain hydrocarbons (Kurtys et al., 2016; Strieder et al., 2019). The *p*-AV indicates the amounts of aldehydes and ketones (secondary oxidation products) and the TBARS value measures the formation of secondary oxidation products, such as malondialdehyde, alkenals, and alkadienals (Shahidi et al., 2003). These measurements have been used to determine the oxidation status of fats and oils.

The data showed a rapid increase in the PVs in duck, chicken, and swine skin fats up to 40 to 50 days of storage (Fig. 1B). After that time, the PVs declined until the end of the storage period. However, in bovine skin fats, the PV increased more gradually and the value was similar to duck skin fats at 90 days (Fig. 1B). This suggests that peroxides were rapidly produced at the beginning of storage in warm conditions (60°C) because of the oxidation of fats. The peroxide product may be further broken down to secondary products, such as aldehydes and ketones (Afonso et al., 2016). This chemical reaction explains the




Fig. 1. Changes in the oxidation stability of animal skin (duck, chicken, swine, and bovine) fats during storage at 60°C for 90 days. (A) The acid value (AV), (B) peroxide value (PV), (C) *p*-anisidine value (*p*-AV), (D) thiobarbituric acid reactive substances (TBARS), (E) conjugated dienes (CD), and (F) total oxidation (totox) value in animal skin fats were determined. The error bars indicate standard deviations.

decline in PVs after 40 to 50 days of storage.

The *p*-AV showed a rapid increase up to 30 to 40 days of storage in duck, chicken, and swine skin fats (Fig. 1C). These *p*-AVs remained constant until the end of the storage period and there was no significant difference between the three groups at the end of storage ($P > 0.05$). In contrast, the *p*-AV in bovine skin fat gradually increased during 60 days of storage (Fig. 1C). The *p*-AVs tended to depend on the USFA content ($p < 0.05$). High USFA contents have been shown to be the major factor influencing *p*-AV in samples (Zuo et al., 2017). Our *p*-AV data also demonstrated a relationship between the amount of USFA and the production of aldehydes and ketones during storage.

Analyzing TBARS is a common technique for measuring lipid peroxidation, particularly the levels of malondialdehyde. In our data, the TBARS value was significantly higher at the beginning of the storage period (at 10 to 40 days) in duck skin fat

(Fig. 1D). Chicken skin fat was relatively higher than swine and bovine skin fats (at 10 to 20 days) (Fig. 1D). These results imply that the higher lipid peroxidation in duck and chicken skin fats was due to higher levels of USFA which were more susceptible to oxidation. SFA is less vulnerable to lipid oxidation and that resulted in the lower TBARS value observed in bovine skin fats. Also, higher levels of PV (primary oxidation products) in duck, chicken, and swine fat might have caused the higher TBARS values. In fact, the unstable peroxides can be the result of the decomposition of polyunsaturated fatty acids (Gheisari, 2011). Our data suggest that the addition of antioxidants to duck fats may be useful for improving oxidative stability during storage.

CD is a measure of the oxidative stability of fats or oils where CD is a lipid hydroperoxide formed by triplet oxygen or singlet oxygen (Akhtar et al., 2018). When the oxidation of lipids progresses, the unconjugated double bonds are converted to conjugated structures that absorb ultraviolet radiations of 233 nm (Peri and Saguy, 2015). Our data showed that the CD values gradually increased from the beginning of the storage period in duck, chicken, and swine skin fats (Fig. 1E). The CD value of these skin fats reached the highest point at the end of the storage period. Bovine skin fat, however, remained at low levels during 90 days of storage (Fig. 1E). In fact, a high level of CD values is associated with the presence of higher USFA (Iqbal and Bhanger, 2007). Thus, the observed high CD values for duck skin fat were due to the higher USFA content.

The totox value is known as the sum of 2PV and *p*-AV. The totox value provides the overall oxidative status of oils and fats (Wai et al., 2009) and lower totox values are expected in better quality cooking oils (Halim et al., 2016). Similar trends of totox values were observed in duck, chicken, and swine skin fats, whereas duck fats showed higher totox values (Fig. 1F). Bovine skin fat showed significantly lower totox values compared to other animal skin fats ($p < 0.05$). In duck, chicken, and swine fats, the totox values continuously increased around 40 to 50 days, then decreased until the end of the storage period. This result has been attributed to the formation of secondary oxidation products which replaced the primary oxidation products (Kasimoglu et al., 2018). These data might also be explained by the ratio of USFA/SFA in the samples. In this regard, a study found that soybean oil containing high amounts of USFA had higher totox values than palm oil with a higher percentage of SFA, such as palmitic acid (Abdulkarim et al., 2007).

Viscosity

Changes in the viscosity of the animal skin fats during the 90-day storage period are presented in Fig. 2. Viscosities increased as the storage time increased in duck, chicken, and swine skin fats ($p < 0.05$). Duck skin fats showed the highest viscosity values, while bovine skin fats showed relatively lower values during the storage period. The formation of high molecular compounds by polymerization during the thermal oxidation process can change the rheological properties (Shin and Kim, 1982). This suggests that polymerized compounds were higher in duck skin fats. Specifically, the increase of viscosity has been explained by the conjugation and isomerization of double bonds in fatty acids and the formation of polymers during thermal oxidation (Ahn et al., 2008). Interestingly, the viscosity of swine skin fat rapidly increased at 90 days. In fact, swine fat contains higher amounts of USFA, particularly linoleic acid (C18:2), than bovine fat. Thus, as shown for the CD (Fig. 1E), the increased viscosity in swine skin fat may be due to the conjugation of double bonds, resulting in polymer formation.

Conclusion

This study evaluated four different animal skin fats (duck, chicken, swine, and bovine fats) extracted with an identical fat extraction method, PHWE. The oxidative stability and quality characteristics of the fats were evaluated under accelerated




Fig. 2. Changes in the viscosity of animal skin (duck, chicken, swine, and bovine) fats were analyzed using a DV-E viscometer during storage at 60°C for 90 days. The measurements were repeated five times for 30 s and the results are expressed as centipoise (cP) units. The error bars indicate SD.

storage conditions (at 60°C, for 90 days). The data demonstrated that duck skin contained more USFA than other animal skin fats, indicating that duck fat is a healthier edible oil for human consumption. However, duck skin fat was more susceptible to oxidation than other animal skin fats in hot storage conditions. This was probably due to the higher level of USFA in duck skin fats. Supplementation of duck fat with antioxidants might be a reasonable way to prevent oxidation during long-term storage. Further studies on the effects of antioxidant supplementation in duck fat and specific health effects in the human body are needed.

Conflict of Interest

The authors declare no potential conflict of interest.

Acknowledgments

This research was supported by Korea Institute of Planning and Evaluation for Technology in Food, Agriculture, Forestry and Fisheries (IPET) through High Value-added Food Technology Development Program, funded by Ministry of Agriculture, Food and Rural Affairs (MAFRA) (2018-118011).

Author Contributions

Conceptualization: Shin DM. Data curation: Shin DM. Formal analysis: Shin DM, Kim DH, Yune JH, Kwon HC, Kim HJ. Methodology: Shin DM. Software: Shin DM. Validation: Seo HG, Han SG. Investigation: Shin DM, Kim DH, Yune JH, Kwon HC, Kim HJ. Writing - original draft: Shin DM, Han SG. Writing - review & editing: Shin DM, Kim DH, Yune JH, Kwon HC, Kim HJ, Seo HG, Han SG.

Ethics Approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

References

- Abdulkarim SM, Long K, Lai OM, Muhammad SKS, Ghazali HM. 2007. Frying quality and stability of high-oleic moringa oleifera seed oil in comparison with other vegetable oils. *Food Chem* 105:1382-1389.
- Afonso C, Bandarra NM, Nunes L, Cardoso C. 2016. Tocopherols in seafood and aquaculture products. *Crit Rev Food Sci Nutr* 56:128-140.
- Ahn MS, Suh MS, Kim HJ. 2008. Measurement of trans fatty acid formation and degree of rancidity in fat and oils according to heating conditions. *J Korean Soc Food Cult* 23:469-478.
- Akhtar S, Tanveer M, Ismail A, Ismail T, Hussain M. 2018. Safety evaluation of oil samples collected from different food points of multan city of Pakistan. *Int J Food Allied Sci* 3:43-48.
- AOAC Internatioanl. 2007. Official methods of analysis of AOAC International. 2nd ed. AOAC International, Gaithersburg, MD, USA.
- Austin GL, Ogden LG, Hill JO. 2011. Trends in carbohydrate, fat, and protein intakes and association with energy intake in normal-weight, overweight, and obese individuals: 1971–2006. *Am J Clin Nutr* 93:836-843.
- Bowen KJ, Kris-Etherton PM, West SG, Fleming JA, Connelly PW, Lamarche B, Couture P, Jenkins DJ, Taylor CG, Zahrada P. 2019. Diets enriched with conventional or high-oleic acid canola oils lower atherogenic lipids and lipoproteins compared to a diet with a western fatty acid profile in adults with central adiposity. *J Nutr* 149:471-478.
- Chemat F, Vian MA, Cravotto G. 2012. Green extraction of natural products: Concept and principles. *Int J Mol Sci* 13:8615-8627.
- Chen X, Du X, Shen J, Lu L, Wang W. 2017. Effect of various dietary fats on fatty acid profile in duck liver: Efficient conversion of short-chain to long-chain omega-3 fatty acids. *Exp Biol Med* 242:80-87.
- Choe E, Min DB. 2006. Mechanisms and factors for edible oil oxidation. *Compr Rev Food Sci Food Saf* 5:169-186.
- Cobos A, Veiga A, Díaz O. 2000. Chemical and fatty acid composition of meat and liver of wild ducks (*Anas platyrhynchos*). *Food Chem* 68:77-79.
- Da Silva SL, Amaral JT, Ribeiro M, Sebastiao EE, Vargas C, De Lima Franzen F, Schneider G, Lorenzo JM, Fries LLM, Cichoski AJ, Campagnol PCB. 2019. Fat replacement by oleogel rich in oleic acid and its impact on the technological, nutritional, oxidative, and sensory properties of bologna-type sausages. *Meat Sci* 149:141-148.
- Dos Santos Freitas L, Jacques RA, Richter MF, Da Silva AL, Caramao EB. 2008. Pressurized liquid extraction of vitamin e from brazilian grape seed oil. *J Chromatogr A* 1200:80-83.
- Encinar JM, Gonzalez JF, Sanchez N, Nogales-Delgado S. 2019. Sunflower oil transesterification with methanol using immobilized lipase enzymes. *Bioprocess Biosyst Eng* 42:157-166.
- FAOSTAT. 2018. Meat, poultry. In Livestock primary. Available from: <http://www.fao.org/faostat/en/#search/duck>. Accessed at Jan 03, 2019.
- Ghadge SV, Raheman H. 2005. Biodiesel production from mahua (*Madhuca indica*) oil having high free fatty acids. *Biomass Bioenerg* 28:601-605.
- Gheisari HR. 2011. Correlation between acid, tba, peroxide and iodine values, catalase and glutathione peroxidase activities of chicken, cattle and camel meat during refrigerated storage. *Vet World* 4:153-157.
- Ghobadi S, Akhlaghi M, Shams S, Mazloomi SM. 2018. Acid and peroxide values and total polar compounds of frying oils in fast food restaurants of Shiraz, Southern Iran. *Int J Nutr Sci* 3:25-30.

- Halim Y, Natania, Halim JM, Soedirga LC, Yakhi LA. 2016. Physical and chemical characteristics of frying oil in Indonesia in a repeated frying model. *J Chem Pharm Res* 8:583-589.
- Hariri N, Thibault L. 2010. High-fat diet-induced obesity in animal models. *Nutr Res Rev* 23:270-299.
- Hashempour-Baltork F, Torbati M, Azadmard-Damirchi S, Savage GP. 2017. Quality properties of sesame and olive oils incorporated with flaxseed oil. *Adv Pharm Bull* 7:97-101.
- Heo KN, Choo HJ, Kim CD, Kim SH, Kim HK, Lee MJ, Son BR, Choi HC, Hong EC. 2013. Changes of fatty acids and amino acids contents of Korean native commercial ducks meats with different raising periods. *Korean J Poult Sci* 40:235-241.
- Huda N, Seow EK, Normawati MN, Aisyah NNM, Fazilah A, Easa AM. 2013. Effect of duck feet collagen addition on physicochemical properties of surimi. *Int Food Res J* 20:537-544.
- Iqbal S, Bhangar MI. 2007. Stabilization of sunflower oil by garlic extract during accelerated storage. *Food Chem* 100:246-254.
- Jo Y, An KA, Arshad MS, Kwon JH. 2018. Effects of e-beam irradiation on amino acids, fatty acids, and volatiles of smoked duck meat during storage. *Innov Food Sci Emerg Technol* 47:101-109.
- Kang G, Seong PN, Cho S, Moon S, Park K, Kang SM, Park BY. 2014. Effect of addition duck skin on quality characteristics of duck meat sausages. *Korean J Poult Sci* 41:45-52.
- Kasimoglu Z, Tontul I, Soyulu A, Gulen K, Topuz A. 2018. The oxidative stability of flavoured virgin olive oil: The effect of the water activity of rosemary. *J Food Meas Charact* 12:2080-2086.
- Khan MA, Ali S, Yang H, Kamboh AA, Ahmad Z, Tume RK, Zhou G. 2019. Improvement of color, texture and food safety of ready-to-eat high pressure-heat treated duck breast. *Food Chem* 277:646-654.
- Kronholm J, Hartonen K, Riekkola ML. 2007. Analytical extractions with water at elevated temperatures and pressures. *Trends Anal Chem* 26:396-412.
- Kurtys E, Eisel ULM, Verkuyl JM, Broersen LM, Dierckx RAJO, De Vries EFJ. 2016. The combination of vitamins and omega-3 fatty acids has an enhanced anti-inflammatory effect on microglia. *Neurochem Int* 99:206-214.
- Kyriakidis NB, Katsiloulis T. 2000. Calculation of iodine value from measurements of fatty acid methyl esters of some oils: Comparison with the relevant american oil chemists society method. *J Am Oil Chem Soc* 77:1235-1238.
- Li J, Sun Q. 2019. Consumption of saturated fatty acids and coronary heart disease risk. *Int J Cardiol* 279:27-28.
- Marion JE, Woodroof JG. 1963. The fatty acid composition of breast, thigh, and skin tissues of chicken broilers as influenced by dietary fats. *Poult Sci* 42:1202-1207.
- Martin MP, Asensio CM, Nepote V, Grosso NR. 2018. Improving quality preservation of raw peanuts stored under different conditions during a long-term storage. *Eur J Lipid Sci Technol* 120:1800150.
- Moghtadaei M, Soltanizadeh N, Goli SAH. 2018. Production of sesame oil oleogels based on beeswax and application as partial substitutes of animal fat in beef burger. *Food Res Int* 108:368-377.
- Moro C, Capel F. 2019. Regulation of skeletal muscle metabolism by saturated and monounsaturated fatty acids. In *Nutrition and skeletal muscle*. Walrand S (ed). Academic Press, London, UK. pp 367-378.
- Mugendi JB, Sims CA, Gorbet DW, O'keefe SF. 1998. Flavor stability of high-oleic peanuts stored at low humidity. *J Am Oil Chem Soc* 75:21-25.
- Muguerza E, Gimeno O, Ansorena D, Bloukas JG, Astiasaran I. 2001. Effect of replacing pork backfat with pre-emulsified olive oil on lipid fraction and sensory quality of chorizo de pamplona—a traditional spanish fermented sausage. *Meat Sci* 59:251-258.
- Mustafa A, Turner C. 2011. Pressurized liquid extraction as a green approach in food and herbal plants extraction: A review.

- Anal Chim Acta 703:8-18.
- Naz S, Siddiqi R, Sheikh H, Sayeed SA. 2005. Deterioration of olive, corn and soybean oils due to air, light, heat and deep-frying. *Food Res Int* 38:127-134.
- Papastergiadis A, Mubiru E, Van Langenhove H, De Meulenaer B. 2012. Malondialdehyde measurement in oxidized foods: Evaluation of the spectrophotometric thiobarbituric acid reactive substances (TBARS) test in various foods. *J Agric Food Chem* 60:9589-9594.
- Park KS, Park HS, Choi YJ, Lee JS, Park SS, Jung IC. 2014. Comparison of fatty acid and nutritional composition of Korean native black cattle and Hanwoo. *Korean J Food Cook Sci* 30:556-563.
- Peri I, Saguy IS. 2015. Continuous injection of water and antioxidants possible roles on oil quality during frying. *LWT-Food Sci Technol* 64:919-925.
- Plaza M, Turner C. 2015. Pressurized hot water extraction of bioactives. *Trends Anal Chem* 71:39-54.
- Rekas A, Wroniak M, Krygier K. 2015. Effects of different roasting conditions on the nutritional value and oxidative stability of high-oleic and yellow-seeded *Brassica napus* oils. *Grasas Aceites* 66:e092.
- Resende MT, Campisi-Pinto S, Linder C, Wiesman Z. 2019. Multidimensional proton nuclear magnetic resonance relaxation morphological and chemical spectrum graphics for monitoring and characterization of polyunsaturated fatty-acid oxidation. *J Am Oil Chem Soc* 96:125-135.
- Shahidi F, Desilva C, Amarowicz R. 2003. Antioxidant activity of extracts of defatted seeds of niger (*Guizotia abyssinica*). *J Am Oil Chem Soc* 80:443-450.
- Shin AJ, Kim DH. 1982. Studies on thermal oxidation of soybean oil-i. Changes in some chemical and physical properties of a soybean oil during thermal oxidation. *Korean J Food Sci Technol* 14:257-264.
- Strieder MM, Engelmann JI, Pohndorf RS, Rodrigues PA, Juliano RS, Dotto GL, Pinto LAA. 2019. The effect of temperature on rice oil bleaching to reduce oxidation and loss in bioactive compounds. *Grasas Aceites* 70:e287.
- Symoniuk E, Ratusz K, Krygier K. 2019. Evaluation of the oxidative stability of cold-pressed rapeseed oil by rancimat and pressure differential scanning calorimetry measurements. *Eur J Lipid Sci Technol* 121:1800017.
- Talcott ST, Passeretti S, Duncan CE, Gorbet DW. 2005. Polyphenolic content and sensory properties of normal and high oleic acid peanuts. *Food Chem* 90:379-388.
- US Department of Agriculture [USDA], US Department of Health and Human Services [HHS]. 2015. 2015–2020 dietary guidelines for americans. USDA/HHS, Washington, DC, USA.
- Wai WT, Saad B, Lim BP. 2009. Determination of totox value in palm oleins using a fi-potentiometric analyzer. *Food Chem* 113:285-290.
- Warner K, Frankel EN, Mounts TL. 1989. Flavor and oxidative stability of soybean, sunflower and low erucic acid rapeseed oils. *J Am Oil Chem Soc* 66:558-564.
- Yan H, Zhang J, Gao J, Huang Y, Xiong Y, Min S. 2018. Towards improvement in prediction of iodine value in edible oil system based on chemometric analysis of portable vibrational spectroscopic data. *Sci Rep* 8:14729.
- Zhou D, Zhou F, Ma J, Ge F. 2019. Microcapsulation of *Ganoderma lucidum* spores oil: Evaluation of its fatty acids composition and enhancement of oxidative stability. *Ind Crop Prod* 131:1-7.
- Zuo W, Hu X, Yang Y, Jiang L, Ren L, Huang H. 2017. Development of an improved method to determine saturated aliphatic aldehydes in docosahexaenoic acid-rich oil: A supplement to p-anisidine value. *Eur J Lipid Sci Technol* 119:1700243.