



ARTICLE

Isolation of the Cholesterol-Assimilating Strain *Pediococcus acidilactici* LRCC5307 and Production of Low-Cholesterol Butter

Yunsik Kim^{1,†}, Seokmin Yoon^{2,†}, Hyejung Shin³, Miyoun Jo², Sunmin Lee², and Sae-hun Kim^{3,*}

¹Department of Biosystems and Biotechnology, College of Life Science and Biotechnology, Korea University, Seoul 02841, Korea

²Food-Biotech Team, Division of Basic Research, Lotte R&D Center, Seoul 07594, Korea

³Department of Food Bioscience and Technology, College of Life Science and Biotechnology, Korea University, Seoul 02841, Korea

OPEN ACCESS

Received September 15, 2020

Revised December 8, 2020

Accepted December 26, 2020

*Corresponding author : Sae-hun Kim
Department of Food Bioscience and Technology, College of Life Science and Biotechnology, Korea University, Seoul 02841, Korea
Tel: +82-2-3290-3055
Fax: +82-2-3290-3506
E-mail: saehkim@korea.ac.kr

*ORCID
Yunsik Kim
<https://orcid.org/0000-0001-8608-4951>
Seokmin Yoon
<https://orcid.org/0000-0002-7004-5947>
Hyejung Shin
<https://orcid.org/0000-0003-2571-6294>
Miyoun Jo
<https://orcid.org/0000-0001-6248-6091>
Sunmin Lee
<https://orcid.org/0000-0002-8438-3365>
Sae-hun Kim
<https://orcid.org/0000-0002-0990-2268>

† These authors contributed equally to this work.

Abstract The objective of the present study was to evaluate the cholesterol-assimilation ability of lactic acid bacteria (LAB), which were isolated from *kimchi*, a Korean traditional fermented cabbage. The isolated strain, using modified MRS medium, showed 30.5% cholesterol assimilation activity and was named *Pediococcus acidilactici* LRCC5307. Types and concentrations of bile were investigated for their effects on increasing the cholesterol assimilation ability of the LRCC5307 strain, a 74.5% decrease in cholesterol was observed when 0.2% bile salts were added. In addition, the manufacture of low-cholesterol butter using LRCC5307 was examined. After fermentation, LRCC5307 with butter showed 8.74 Log CFU/g viable cells, pH 5.43, and a 11% decrease in cholesterol. These results suggest that LRCC5307 could help in the production of healthier butter by decreasing cholesterol and including living LAB.

Keywords cholesterol, *Pediococcus acidilactici*, assimilation, bile salts, butter

Introduction

Cholesterol, an essential substrate used in cell membranes and for the production of certain hormones, is obtained through food or produced in the liver. However, an excessive amount of cholesterol causes hypercholesterolemia, a known risk factor for coronary artery disease (Akalin et al., 1997; Anderson and Gilliland, 1999; Lin and Chen, 2000). For this reason, the World Health Organization (WHO) and the American Heart Association recommend that consumers limit their intake of saturated fatty acids and cholesterol to reduce the risk of coronary artery disease (Hansel et al., 2007). However, animal foods such as eggs, meat, and shrimp, in addition to milk and dairy

products, which are consumed on a daily basis, contain high levels of cholesterol.

In general, dairy products are considered healthy. However, products such as butter, cream, and certain types of cheese, which contain high amounts of fat, are not necessarily healthy. In particular, butter is used in various ways in the dairy industry. It serves as a basic material for other dairy products and is provided directly to consumers due to its unique flavor and sensuality. Nevertheless, cholesterol content in butter is approximately 210 mg/100 g in general, which is higher than that in cream cheese (110 mg/100 g) and condensed milk (35 mg/100 g). Therefore, it is essential to manage cholesterol intake from butter.

Various processing methods that remove cholesterol from butter have been studied. These include new processes, such as ultrasound, nano-filtration, accelerated solvent extraction, and solid-phase extraction, as well as methods involving β -cyclodextrin and lactic acid bacteria (LAB) (Allègre et al., 2006; Jia et al., 2020; Lye et al., 2012; Richter et al., 1996). In particular, there have been many studies attempting to reduce cholesterol in butter using the chemical mechanisms of β -cyclodextrin; however, this process is costly, resulting in relatively expensive butter products. As such, this process reduces the flavor components of butter, and these products often fail to be selected by consumers for both sensory and economic reasons (Aloğlu and Öner, 2006; Kim et al., 2006).

Meanwhile, there are currently several efforts underway to apply biological methods using microbes, which incur no extra costs and are thought to have beneficial effects on sensory properties, taste, and texture. In particular, studies have focused on LAB and yeast, which have demonstrated not only reductions in serum cholesterol after intake into the body, but also assimilation of cholesterol during the fermentation process. Aloğlu et al. (2016) found that cholesterol in butter could be assimilated by probiotic LAB and yeast (Aloğlu and Öner, 2006; Aloğlu et al., 2015). Additionally, Gilliland et al. (1985) and Pan et al. (2011) reported, respectively, that *Lactobacillus acidophilus* and *Lactobacillus fermentum* SM-7 assimilate cholesterol *in vitro*.

In this study, we examined the effects of cholesterol-assimilating LAB on butter production. To this end, we isolated LAB from a traditional fermented food of Korea, *kimchi*, and acquired strains that exhibited cholesterol-assimilating abilities when grown in a medium containing cholesterol as the only carbon source. Further, to investigate the effects of bile on cholesterol assimilation, various types and concentrations of bile were tested. The ultimate goal of this study was to produce butter with a reduced amount of cholesterol using bile and isolated LAB to provide butter, which otherwise is high in cholesterol despite its excellent flavor and nutritional value, as a healthy food.

Materials and Methods

Isolation of cholesterol-assimilating lactic acid bacteria

To isolate bacterial strains, approximately 200 *kimchi* samples were collected from a traditional market in Korea and diluted in sterile distilled water before being homogenized with a mill homogenizer. Depending on the degree of fermentation, there were 66, 95, and 45 kinds of *kimchi* less than 3-days old, less than 4-weeks old, and more than 1-month old, respectively. After serial dilution of the homogenized solution, the appropriate concentration of solution was spread on de Man, Rogosa, and Sharpe (MRS) agar containing 0.002% bromo-cresol purple and cultured for 48 h, following which, colonies showing a yellow hue were selected. To isolate the selected strains that showed cholesterol-assimilating ability, cholesterol-MRS agar was prepared based on the method of Gilliland et al. (1985), where the only carbon source was cholesterol. The specific composition of the agar medium was as follows: cholesterol 0.2 g/L, proteose peptone 10 g/L,

ammonium citrate 2 g/L, sodium acetate 5 g/L, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.1 g/L, MnSO_4 0.05 g/L, K_2HPO_4 2 g/L, and agar 15 g/L. After placing sterilized 8-mm paper discs (Paperdisc, Advantec, Ehime, Japan) onto the agar medium, 10 μL of each seed culture broth from the purified strains was added, and the agar was incubated for 48 h at 37°C. Subsequently, the areas surrounding the paper discs were examined for the growth of bacterial colonies, and strains that used cholesterol as a nutrient were selected. As a positive control for cholesterol assimilation, 10 μL of the cholesterol-oxidizing enzyme peroxidase solution (peroxidase from horseradish, 1,000 U/mL, Sigma-Aldrich, St. Louis, MO, USA) was used, and as a negative control, 10 μL of cholesterol-MRS broth without bacteria inoculation was used.

Measuring the cholesterol-assimilating ability of the isolated strains

The primarily selected strains were inoculated into 15-mL tubes containing 10 mL of MRS broth, and after 48 h of incubation at 37°C, the tubes were centrifuged (10,000 \times g, 10 min), the supernatant was removed, and the pellet was washed with 0.1 M phosphate buffer (pH 6.8). After repeating this process three times, 5 mL of 0.1 M phosphate buffer was added to the pellet to obtain a bacterial suspension and complete the production of the seed culture. After inoculating 1% of the seed culture in MRS broth with cholesterol as the only carbon source, the broth was incubated for 48 h at 37°C. Thereafter, 10 mL of isopropyl alcohol was added, the mixture was vortexed for 5 min and centrifuged at 5,600 \times g for 5 min, followed by collection of 4 mL of the supernatant. To the supernatant, 100 μL of 4 M KOH, 1.5 g of NaCl, and 4 mL of distilled water were added, the mixture was centrifuged (5,600 \times g, 5 min), the supernatant was collected, and the cholesterol concentration was measured using a gas chromatography-tandem mass spectrometer system (GC-MS/MS 5977A, Agilent Technologies, Santa Clara, CA, USA). Medium without bacterial inoculation was used as the negative control, and *L. acidophilus* ATCC43121, the most studied strain regarding serum cholesterol-lowering effects *in vivo*, was obtained from a U.S. bioresource center (ATCC, Manassas, VA, USA) and used as a comparison group.

Strain identification

The API 50 CHL kit (Biomerieux, Rue des Aqueducs, France) was used as a simple way of identifying the isolated bacterial strains by measuring sugar utilization. Colonies cultured in MRS broth were inoculated into the API 50 CHL kit and incubated at 37°C. After 24 h and 48 h, the change in color (yellow) was measured according to the type of sugar. The measurements were used for simple identification of the bacteria via the Biomerieux DB (<https://apiweb.biomerieux.com>).

In addition, 16S rDNA was analyzed for genetic identification. Specifically, after extracting genomic DNA using a genomic DNA preparation kit (Promega, Madison, WI, USA), a polymerase chain reaction (PCR) reaction was run using the universal primer pair 27F (5'-AGA GTT TGA TCC TGG CTC AG-3') and 1492R (5'-TAC GGY TAC CTT GTT ACG ACT T-3') to amplify the 16S rDNA (Buck and Gilliland, 1994). The PCR products were purified using a QIA quick PCR kit (QIAGEN, Hilden, Germany) and sequencing was outsourced to Macrogen (Seoul, Korea). The base sequences were then compared with the NCBI GenBank DB using BLAST analysis (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>).

Determination of cholesterol assimilation with different types and concentrations of bile

After activation and seed culturing of the isolated bacterial strains into MRS agar and broth, the seed culture was inoculated into cholesterol-MRS broth containing different types of bile. After adding 0.1% (w/v) of either bile extract, oxgall powder, bile acid, or bile salt (Sigma-Aldrich), 1 M NaOH was used to adjust the pH to 6.8 \pm 0.2. Following incubation, the bile that produced the greatest decrease in cholesterol concentration was selected as optimal. The selected bile was added to

cholesterol-MRS broth at concentrations of 0, 0.1, 0.2, 0.3, 0.4, and 0.5% (w/v), after which 1% (v/v) of the seed culture of the isolated strains was also inoculated, and the broth was cultured. As stated previously herein, the bile concentration that resulted in the greatest decrease in cholesterol was selected as the optimal concentration.

Fermentation of butter for the reduction of cholesterol

LRCC5307 LAB was cultured, centrifuged, and washed. The supernatant was removed, and cells were diluted in PBS (pH 6.8±0.2) to prepare seed cultures. In a 1-L beaker immersed in a 90°C water bath, 500 g of commercially available unsalted butter was completely dissolved. While stirring the butter in a paste form at 150 rpm using a magnetic mixer (digital stirring hot plates, Corning®, Corning, NY, USA) maintained at 40°C or higher, 1 g of bile salts and 2.5 g of seed culture were added for 1 h to increase dispersibility. The beaker was sealed, placed on a magnetic mixer, and put in an incubator for fermentation for 72 h at 42°C to prevent the butter from hardening. After incubation, viable cells, pH, and cholesterol concentrations were assessed and compared to those before incubation.

Statistical analysis

All data are presented as means (±SDs) of at least three independent experiments; each experiment had three replicates of each sample. Data were analyzed statistically using IBM SPSS Statistics software version 25.0 (IBM, Armonk, NY, USA). The statistical differences between the mean values of test groups were analyzed by one-way analysis of variance and a paired sample t-test. Statistical significance was defined as $p < 0.05$. Multiple comparisons between different groups were assessed using Duncan's test.

Results and Discussion

Screening of cholesterol-assimilating bacterial strains

Approximately 300 strains of bacteria were isolated from traditional *kimchi*. Fig. 1 shows the patterns of the colonies that formed after the incubation of each of these strains on cholesterol-MRS agar. As shown in Fig. 1, when peroxidase was applied as a positive control, there was a clearly visible brown halo around the paper disc; however, when only the cholesterol-MRS broth was applied as a negative control, there was no significant change on the paper disc. When the isolated strains were added to the medium, although there were differences in size, the results could be divided into two major patterns: those that formed a clear halo (strains A and D) and those that did not (strains B, C, E, and F).

Several bacteria and their enzymes have been reported to have the ability to degrade cholesterol and 7-ketocholesterol. The degradation of these compounds is initiated by mechanisms like cholesterol oxidation (Ghosh and Khare, 2016). These bacteria have been reported to be involved in the biodegradation of cholesterol via cholesterol oxidase. Cholesterol oxidase is a FAD-dependent (flavin adenine dinucleotide) enzyme that catalyzes the oxidation and isomerization of sterols to sterones, typically cholesterol (5-cholesten-3- β -ol) to 4-cholesten-3-one (cholestenone) (Lashgarian et al., 2016; Pan et al., 2011; Sakodinskaya and Ryabov, 2000). Studies have shown that cholestenone, which is produced as a metabolite, is safe and can be used to control obesity, treat liver disease, and prevent keratinization of the skin (Elia et al., 2019). Wali et al. (2019) isolated *Bacillus pumilus* W1 and *Serratia marcescens* W8 from soil contaminated with oil and reported that these strains degrade cholesterol and produce red colonies in M9 medium containing 0.1% cholesterol as the only carbon source.

Thus, of the 300 isolated bacterial strains, 54 formed red halos and halo sizes were measured to determine their cholesterol-




Fig. 1. Cholesterol-MRS agar culture of bacterial strains isolated from traditional *kimchi*. P-CNT, positive control: peroxidase 1,000 U 10 μ L; N-CNT, negative control: cholesterol-MRS broth 10 μ L; Strains A, *L. acidophilus* ATCC43121; Strain B, *E. faecalis* EF5326 (isolated strain from kimchi); Strain C, *Pediococcus acidilactici* PA5296; Strain D, *P. acidilactici* LRCC5307; Strain E, *L. plantarum* LP5272; Strain F, *Lactobacillus brevis* LB5238 (isolated strain from kimchi) strains isolated from *kimchi* that formed yellow colonies on MRS agar.

assimilation ability. There were 33, 12, and nine strains with a halo diameter of 10–12 mm, 12–14 mm, and 15 mm and above, respectively. We aimed to determine the cholesterol assimilating ability of these strains quantitatively.

Measurement of cholesterol-assimilating ability

The 54 bacterial strains isolated from traditional *kimchi* were inoculated into cholesterol-MRS broth, and the cholesterol concentration in the medium was measured after cultivation. The five strains that produced a decrease in the cholesterol concentration are shown in Table 1. The initial mean cholesterol concentration in the medium was 206.25 ± 1.68 mg/L, and after 48 h of incubation with the comparison strain *L. acidophilus* ATCC43121, there was a 23.0% decrease in cholesterol to 158.83 ± 3.39 mg/L. The strain that showed the highest decrease in cholesterol was *P. acidilactici* LRCC5307, which produced a 30.5% reduction to 143.38 ± 2.48 mg/L, and this was closely followed by *P. acidilactici* PA5296, which produced a 28.0% reduction to 148.43 ± 1.84 mg/L. Other strains showed a cholesterol concentration of ≥ 170 mg/L after cultivation, representing a decrease of $< 20\%$. Thus, the LRCC5307 strain was selected as the strain with the highest cholesterol-assimilation ability.

Pereira and Gibson (2002) assessed the *in vitro* cholesterol-assimilation effects of LAB and bifidobacteria isolated from the human gut, cholesterol were decreased by 47% in the medium containing 100 mg/L and 0.4% of oxgall.

Table 1. The decrease in cholesterol with the isolated bacterial strains

Strain	Cholesterol ¹⁾ (mg/L)	Cholesterol reduction rate ²⁾ (%)
Blank	206.25 ± 1.68	–
<i>L. acidophilus</i> ATCC43121	158.83 ± 3.39	23.0 ^b
<i>P. acidilactici</i> LRCC5307	143.38 ± 2.48	30.5 ^c
<i>P. acidilactici</i> PA5296	148.43 ± 1.84	28.0 ^c
<i>L. plantarum</i> LP5272	168.29 ± 3.67	18.4 ^a
<i>E. faecalis</i> EF5315	172.95 ± 1.79	16.1 ^a
<i>P. acidilactici</i> PA5265	175.70 ± 2.20	14.8 ^a

Blank, the initial media without cultivation. *L. acidophilus* ATCC43121 was used as reference strains.

¹⁾ Results are expressed as mean \pm SE (n=3).

²⁾ Results are calculated as (cholesterol concentration of each strain / cholesterol concentration of blank) \times 100.

^{a-c} Means in the same column with different lowercase superscript letters are significantly different at $p < 0.05$.

Considering that the initial amount of cholesterol administered in this study was approximately 200 mg/L and that LRCC5307 decreased cholesterol levels by 30%, an excellent cholesterol-assimilation activity of approximately 17% was observed. Moreover, Anila et al. (2016) demonstrated that culturing *L. casei* and *L. brevis* in medium containing 100 µg/mL of cholesterol resulted in assimilation of 18.18–47.70 µg/mL, which suggests that 17% more cholesterol was reduced compared to that with LRCC5307. As both studies showed excellent cholesterol-assimilation activity when oxgall or oxbile was added to the culture medium, we also decided to assess the effects of bile.

Bacterial strain identification

Table 2 shows the results from the analysis with API 50 CHL to investigate the sugar utilization of the isolated strain LRCC5307. LRCC5307 utilized galactose, glucose, fructose, mannose, cellobiose, lactose, trehalose, and esculin, but not mannitol, sorbitol, salicin, and inulin. When the sugar utilization results were compared with the API website (<https://apiweb.biomerieux.com>), they were similar to those of *P. acidilactici* standard strain (99.9% ID, T index 0.91), and the utilization of rhamnose and salicin were each 75% different. The next closest identified species was *Lactococcus lactis* ssp. *lactis* standard strain 1; however, this showed very low similarity (0.1% ID, T index 0.44) and various contrasting characteristics.

After amplifying the 16S rDNA of the LRCC5307 strain through PCR, sequencing of the 1,440 bp (base-pair) was outsourced to MacroGen, and a homology search was conducted using the NCBI BLASTN program (<https://blast.ncbi.nlm.nih.gov>). After

Table 2. Carbohydrate fermentation patterns analysis of PA5307

Sugar	Result ¹⁾	Sugar	Result ¹⁾	Sugar	Result ¹⁾
Control	–	α-Methyl-D-mannoside	–	Turanose	–
Glycerol	–	α-Methyl-D-glucoside	–	Lyxose	–
Erythritol	–	N-Acetyl-glucosamine	+	Tagatose	+
D-Arabinose	–	Amygdalin	–	D-Fucose	–
L-Arabinose	+	Arbutin	–	L-Fucose	–
Ribose	+	Esculin	+	D-Arabitol	–
D-Xylose	+	Salicin	–	L-Arabitol	–
L-Xylose	–	Cellobiose	+	Gluconate	–
Adonitol	–	Maltose	–	2-Ketone-gluconate	–
β-Methyl-D-xylose	–	Lactose	+	5-Keto-gluconate	–
Galactose	+	Melibiose	–		
Glucose	+	Sucrose	–		
Fructose	+	Trehalose	+		
Mannose	+	Inulin	–		
Sorbose	–	Melezitose	–		
Rhamnose	–	Raffinose	–		
Dulcitol	–	Starch	–		
Inositol	–	Glycogen	–		
Mannitol	–	Gentiobiose	–		
Sorbitol	–	Gentiobiose	+		

¹⁾ The results were compared against the database from bioMerieux at <https://apiweb.biomerieux.com>.

comparing this sequence with the GenBank database and performing a homology search with the BLASTN program, a phylogenetic tree was constructed using the neighbor joining method as shown in Fig. 2. Based on the results, the strain was identified as *P. acidilactici*, and the species closest in similarity was *P. acidilactici* strain N9. Moreover, the strain was similar to *P. acidilactici* strain CE73b and *P. acidilactici* strain 5541. Therefore, the isolated strain LRCC5307 was named *P. acidilactici* LRCC5307 (Lotte R&D Culture Collection).

Genetic identification, performed by analyzing 16S rDNA of approximately 300 types of LAB isolated from *kimchi*, showed 57, 35, 13, 71, 10, 78, and 31 species of *Leuconostoc mesenteroides*, *L. brevis*, *Lactobacillus pentosus*, *Lactobacillus plantarum*, *L. casei*, *Lactobacillus sakei*, and *P. acidilactici*, respectively. Moreover, a total of 308 species were observed, including four, three, and six species of *Leuconostoc citreum*, *Lactobacillus curvatus*, and *Weissella cibaria*, respectively. According to Jeong et al. (2009), 58.9%, 10.7%, and 7.1% of LAB species isolated from commercially available *kimchi* were *L. plantarum*, *L. casei*, and *Lactobacillus coryniformis*, respectively, whereas 5% each of *L. mesenteroides* and *L. sakei* were also observed. In a study by Song et al. (2015), various LAB species were isolated from *kimchi*, including *P. acidilactici*, in addition to various *Lactobacillus* sp. Moreover, Kim et al. (2019) reported that *P. acidilactici* K10, isolated from *kimchi*, inhibited the adhesion of *Salmonella typhimurium* and *Escherichia coli* O157:H7, which are enteropathogenic bacteria, to HT-29 cells. Therefore, although *P. acidilactici* is not a major species among many LAB that are present in *kimchi*, it is thought that it can be isolated and has various functions. Additionally, we deposited LRCC5307, isolated in this study, into the Korean Culture Center of Microorganisms (Daejeon, Korea) as KCCM11734P, and as of June 13, 2017, it was registered as a Korean patent strain.

Effects of bile type and concentration

Fig. 3 shows the cholesterol-assimilation ability measured when different types of bile were added to a *P. acidilactici* LRCC5307 cultivation. The cholesterol concentration in the medium in the absence of LRCC5307 was 201.73 ± 1.44 mg/L,




Fig. 2. Phylogenetic tree of *Pediococcus acidilactici* LRCC5307. The analyzed sequences were compared against the GenBank database on the NCBI website using BLAST analysis (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>).




Fig. 3. The cholesterol clearing ability of LRCC5307 with the addition of different types of bile. Blank, media with cholesterol, not fermented; Non-bile, LRCC5307 fermented without bile; BE, LRCC5307 fermented with bile extract; OX, LRCC5307 fermented with oxgall; BA, LRCC5307 fermented with bile acids; BS, LRCC5307 fermented with bile salts. Results are expressed as mean \pm SE (n=3). ^{a-d} Means in different marks superscript letters are significantly different at p<0.05.

which decreased to 146.06 \pm 2.01 mg/L in the medium inoculated with LRCC5307 in the absence of bile. When the culture was incubated under the same conditions but with the addition of different types of bile, the cholesterol concentration significantly decreased compared to that after cultivation without bile and produced a 68% decrease in cholesterol to 64.27 \pm 2.84 mg/L.

Fig. 4 shows the results of culturing *P. acidilactici* LRCC5307 treated with different concentrations of bile salts. The cholesterol concentration decreased to 65.64 \pm 1.76 mg/L when 0.1% bile salts were added and 51.54 \pm 1.65 mg/L when 0.2% bile salts were added. The addition of 0.3% and 0.5% bile salts resulted in cholesterol concentrations of 54.38 \pm 2.10 mg/L and 54.29 \pm 1.78 mg/L, respectively, which were no better than those obtained with the addition of 0.2% bile salts.

Microbial assimilation of cholesterol is related to the presence of bile, and cholesterol-assimilation ability has been




Fig. 4. The cholesterol clearing ability of LRCC5307 at different concentrations of bile salts. Blank, media with cholesterol, not fermented; 0–0.5, cholesterol concentration in media, fermented with LRCC5307. Results are expressed as mean \pm SE (n=3). ^{a-c} Means in different marks superscript letters are significantly different at p<0.05.

surmised to increase when there is an appropriate concentration of bile in the medium. Many *in vitro* studies have reported much higher rates of cholesterol assimilation in the presence of bile (Anila et al., 2016; Tok and Aslim, 2010). Gilliland et al. (1985) reported a 70% decrease in cholesterol when *L. acidophilus* was cultured in medium containing $\geq 0.4\%$ oxgall. Pereira and Gibson (2002) reported a significant increase in the cholesterol-reducing ability of *L. casei* Shirota in medium containing 0.4% oxgall. However, the effects are believed to differ depending on the species of microbe used for cholesterol assimilation, and the optimal bile type and concentration differ between microbes. Therefore, to achieve effective cholesterol assimilation, it is important to examine the optimal bile type and concentration for each microbe. In this study, a reduction in cholesterol levels by LRCC5307 occurred when the optimal bile type was bile salts and the optimal concentration was 0.2%.

Reduction of cholesterol levels in butter

Commercially available unsalted butter was fermented for 48 h with LRCC5307, and the viable cells, pH, and cholesterol were measured before and after fermentation (Table 3). The viable cells increased from 7.49 ± 0.10 Log CFU/g before fermentation to 8.74 ± 0.06 Log CFU/g after fermentation, and the pH decreased by approximately 1.2, from 6.62 ± 0.00 before fermentation to 5.43 ± 0.01 after fermentation. The cholesterol concentration decreased by approximately 230 mg/L, or 11% of the initial concentration, from $2,105.21 \pm 27.99$ mg/kg before fermentation to $1,873.16 \pm 15.20$ mg/kg after fermentation. Compared to that with cultivation in MRS medium, the pH was approximately 1.0 units higher and the viable cells were approximately 1 Log CFU/g lower; this was thought to be due to the lack of nutrients and a suitable environment for growth in butter (excessive fat content, etc.) compared to those in a medium that is optimized for LAB growth. Moreover, cell viability, pH, and cholesterol showed statistically significant differences compared to those with cultivation in butter without fermentation.

In butter fermented without bile salts, the number of viable cells and pH were similar to those of butter fermented with bile salt, and there was no statistically significant difference. However, cholesterol was 2017.37 ± 14.73 mg/kg, which was reduced by approximately 4.2% compared to that in butter before fermentation. Therefore, similar results to the increased cholesterol assimilation after the addition of bile salts to MRS medium containing cholesterol were also observed in butter.

To reduce the risk of heart disease, the WHO, the American Heart Association, and others recommend reducing one's fatty acid and cholesterol intake and suggest a maximum daily cholesterol intake of 300 mg/d. Although it varies depending on the region and diet, daily butter intake ranges 1–10 g/d, and as a result, cholesterol intake ranges from 2–20 mg/d. In this study, cholesterol in butter was reduced by 10% with LRCC5307. If this effect can be enhanced to produce butter without a risk of cholesterol, it would broaden options for consumers and greatly contribute to the growth of the butter industry.

Aloğlu and Öner (2006) reported research using 10 strains of LAB to degrade cholesterol in cream and butter. In that study,

Table 3. Butter before and after LRCC5307 fermentation

Contents	Pre-fermentation	Post-fermentation	
		Without bile salts	With bile salts
Viable cells (Log CFU/g)	7.49 ± 0.10^a	8.72 ± 0.10^b	8.74 ± 0.06^b
pH	6.62 ± 0.00^a	5.49 ± 0.06^b	5.43 ± 0.01^b
Cholesterol (mg/kg)	$2,105.21 \pm 27.99^a$	$2,017.37 \pm 14.73^b$	$1,873.16 \pm 15.20^c$

Results are expressed as mean \pm SE (n=3).

A difference of $p < 0.05$ was considered significant.

^{a-c} Means in the same row with different lowercase superscript letters are significantly different at $p < 0.05$.

Lactobacillus maltaromicus AC3-16 and *L. casei* ssp. *casei* AB16-65 were reported to degrade 0.1%–25% of the cholesterol in butter, although the cholesterol-assimilation ability was assumed to differ depending on the growth of the strain. Meanwhile, Albano et al. (2018) measured the cholesterol-assimilation ability when cheese was produced and matured with seven strains of LAB and reported cholesterol-assimilation rates of 21% for *L. plantarum* VS513 and 18% for *Lactobacillus paracasei* VC213. Using cholesterol-degrading LAB to ferment actual dairy products, such as cream, cheese, and butter, is known to greatly reduce cholesterol-assimilation ability compared to that with fermentation in broth. This is thought to be because, compared to that in broth, the dairy products present a suboptimal environment for LAB growth, and butter is believed to be especially poor for LRCC5307 growth due to its high fat content and low water content. Therefore, in future research it will be important to establish the optimal conditions for butter fermentation using LRCC5307 to improve the cholesterol assimilation effect without negatively impacting the sensory properties of butter.

Conclusion

As excessive amounts of cholesterol can lead to hypercholesterolemia, the WHO and The American Heart Association recommend restricted intake of food with high levels of saturated fatty acids and cholesterol. Although butter has high calorie and fat contents for its high nutritional value and excellent flavor, it contains more than 2,000 mg/kg of cholesterol.

In this study, methods using fermentation and LAB were studied to reduce cholesterol amounts in butter. The cholesterol-assimilating ability of *P. acidilactici* LRCC5307 isolated from *kimchi* was evaluated, and the optimal conditions for cholesterol assimilation by this strain in the presence of different types and concentrations of bile in the cultures were determined. When cholesterol was added to general MRS broth as the only carbon source, there was a 30% reduction in cholesterol, but when 0.2% bile salt was added, the cholesterol concentration decreased by 74.5%. When actual butter was fermented, the cholesterol concentration decreased by approximately 11%, showing potential for this strain to produce lower-cholesterol butter. Therefore, if the cholesterol assimilation rate could be further improved by optimizing the LRCC5307 fermentation conditions for butter, we anticipate that it would enable the production of healthier butter. However, due to properties of the LAB, as fermentation conditions improve, the unique sour taste imparted by LAB would also increase, thus making it important to ensure that this does not drastically alter the particular flavor of butter.

Therefore, in future studies, finding the optimal fermentation condition that does not affect the inherent flavor of butter while reducing cholesterol would be crucial, and this would allow for the manufacturing of healthy butter with consumer acceptance.

Conflicts of Interest

The authors declare no potential conflicts of interest.

Acknowledgements

This work was supported by Lotte Foods Co. Ltd. and Korea University.

Author Contributions

Conceptualization: Kim YS. Data curation: Yoon SM. Formal analysis: Jo MY. Methodology: Jo MY. Software: Lee SM.

Validation: Shin HJ. Investigation: Kim SH. Writing - original draft: Kim YS, Yoon SM. Writing - review & editing: Kim YS, Yoon SM, Shin HJ, Jo MY, Lee SM, Kim SH.

Ethics Approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

References

- Akalin AS, Göncü S, Düzel S. 1997. Influence of yogurt and acidophilus yogurt on serum cholesterol levels in mice. *J Dairy Sci* 80:2721-2725.
- Albano C, Morandi S, Silveti T, Casiraghi MC, Manini F, Brasca M. 2018. Lactic acid bacteria with cholesterol-lowering properties for dairy applications: *In vitro* and *in situ* activity. *J Dairy Sci* 101:10807-10818.
- Allègre C, Moulin P, Gleize B, Pieroni G, Charbit F. 2006. Cholesterol removal by nanofiltration: Applications in nutraceuticals and nutritional supplements. *J Membr Sci* 269:109-117.
- Aloğlu H, Öner Z. 2006. Assimilation of cholesterol in broth, cream, and butter by probiotic bacteria. *Eur J Lipid Sci Technol* 108:709-713.
- Aloğlu HŞ, Özer ED, Öner Z. 2016. Assimilation of cholesterol and probiotic characterization of yeast strains isolated from raw milk and fermented foods. *Int J Dairy Technol* 69:63-70.
- Anderson JW, Gilliland SE. 1999. Effect of fermented milk (yogurt) containing lactic acid bacteria acidophilus L1 on serum cholesterol in hypercholesterolemic humans. *J Am Coll Nutr* 18:43-50.
- Anila K, Kunzes A, Bhalla TC. 2016. *In vitro* cholesterol assimilation and functional enzymatic activities of putative probiotic *Lactobacillus* sp. isolated from fermented foods/beverages of North West India. *J Nutr Food Sci* 6:1-5.
- Buck LM, Gilliland SE. 1994. Comparison of freshly isolated strains of *Lactobacillus acidophilus* of human intestinal origin for ability to assimilate cholesterol during growth. *J Dairy Sci* 77:2925-2933.
- Elia J, Carbone D, Logé C, Ory L, Huvelin JM, Tannoury M, Diab-Assaf M, Petit K, Nazih H. 2019. 4-Cholesten-3-one decreases breast cancer cell viability and alters membrane raft-localized EGFR expression by reducing lipogenesis and enhancing LXR-dependent cholesterol transporters. *Lipids Health Dis* 18:168.
- Ghosh S, Khare SK. 2016. Biodegradation of cytotoxic 7-ketocholesterol by *Pseudomonas aeruginosa* PseA. *Bioresour Technol* 213:44-49.
- Gilliland SE, Nelson CR, Maxwell C. 1985. Assimilation of cholesterol by *Lactobacillus acidophilus*. *Appl Environ Microbiol* 49:377-381.
- Hansel B, Nicolle C, Lalanne F, Tondu F, Lassel T, Donazzolo Y, Ferrières J, Krempf M, Schlienger JL, Verges B, Chapman MJ, Bruckert E. 2007. Effect of low-fat, fermented milk enriched with plant sterols on serum lipid profile and oxidative stress in moderate hypercholesterolemia. *Am J Clin Nutr* 86:790-796.
- Jeong L, Oh CK, OH, MC, Kim SH. 2009. Isolation and identification of lactic acid bacteria from commercial Kimchi. *J Korean Soc Food Sci Nutr* 38:732-741.
- Jia X, Yang N, Qi X, Chen L, Zhao Y. 2020. Adsorptive removal of cholesterol by biodegradable zein-graft- β -cyclodextrin film. *Int J Biol Macromol* 155:293-304.
- Kim JJ, Jung TH, Ahn J, Kwak HS. 2006. Properties of cholesterol-reduced butter made with β -cyclodextrin and added

- evening primrose oil and phytosterols. *J Dairy Sci* 89:4503-4510.
- Kim SH, Kim WJ, Kang SS. 2019. Inhibitory effect of bacteriocin-producing *Lactobacillus brevis* DF01 and *Pediococcus acidilactici* K10 isolated from kimchi on enteropathogenic bacterial adhesion. *Food Biosci* 30:100425.
- Lashgarian HE, Jahanbakhsh S, Shahzamani K. 2016. Molecular identification of cholesterol oxidase enzyme producing *Streptomyces* bacteria in soil of Lorestan province, Iran. *Int J Med Res Health Sci* 5:54-62.
- Lin MY, Chen TW. 2000. Reduction of cholesterol by *Lactobacillus acidophilus* in culture broth. *J Food Drug Anal* 8:97-102.
- Lye HS, Alias KA, Rusul G, Liong MT. 2012. Ultrasound treatment enhances cholesterol removal ability of lactobacilli. *Ultrason Sonochem* 19:632-641.
- Pan DD, Zeng XQ, Yan YT. 2011. Characterization of *Lactobacillus fermentum* SM-7 isolated from koumiss, a potential probiotic bacterium with cholesterol-lowering effects. *J Sci Food Agric* 91:512-518.
- Pereira DIA, Gibson GR. 2002. Cholesterol assimilation by lactic acid bacteria and bifidobacteria isolated from the human gut. *Appl Environ Microbiol* 68:4689-4693.
- Richter BE, Jones BA, Ezzell JL, Porter NL, Avdalovic N, Pohl C. 1996. Accelerated solvent extraction: A technique for sample preparation. *Anal Chem* 68:1033-1039.
- Sakodinskaya IK, Ryabov AD. 2000. Crown ether activates cholesterol oxidase in low water media. *Biotechnol Lett* 22:173-176.
- Song SY, Lee HJ, Kim JH, Oh SJ. 2015. Probiotic characteristics of lactic acid bacteria isolated from home-made kimchi and infant feces. *Curr Top Lactic Acid Bac Probio* 3:76-80.
- Tok E, Aslim B. 2010. Cholesterol removal by some lactic acid bacteria that can be used as probiotic. *Microbiol Immunol* 54:257-264.
- Wali H, Rehman FU, Umar A, Ahmed S. 2019. Cholesterol degradation and production of extracellular cholesterol oxidase from *Bacillus pumilus* W1 and *Serratia marcescens* W8. *BioMed Res Int* 2019:1359528.