



ARTICLE

## Characterization of Novel Amylase-Sensitive, Anti-Listerial Class IId Bacteriocin, Agilicin C7 Produced by *Ligilactobacillus agilis* C7

Jeong Min Yoo, Ji Hoon Song, Robie Vasquez, In-Chan Hwang, Jae Seung Lee, and Dae-Kyung Kang\*

Department of Animal Biotechnology, College of Biotechnology and Bioengineering, Dankook University, Cheonan 31116, Korea



Received May 1, 2023  
Revised June 6, 2023  
Accepted June 7, 2023

\*Corresponding author : Dae-Kyung Kang  
Department of Animal Biotechnology,  
College of Biotechnology and  
Bioengineering, Dankook University,  
Cheonan 31116, Korea  
Tel: +82-41-550-3655  
Fax: +82-41-550-3655  
E-mail: [dkkang@dankook.ac.kr](mailto:dkkang@dankook.ac.kr)

\*ORCID  
Jeong Min Yoo  
<https://orcid.org/0000-0003-0787-3309>  
Ji Hoon Song  
<https://orcid.org/0000-0003-0027-7416>  
Robie Vasquez  
<https://orcid.org/0000-0002-2878-9125>  
In-Chan Hwang  
<https://orcid.org/0000-0001-8268-989X>  
Jae Seung Lee  
<https://orcid.org/0009-0000-0907-4901>  
Dae-Kyung Kang  
<https://orcid.org/0000-0001-9241-1250>

**Abstract** Among various biological agents, bacteriocins are important candidates to control *Listeria monocytogenes* which is a foodborne pathogen. In this study, a novel bacteriocin, named agilicin C7, was isolated from *Ligilactobacillus agilis* C7 showing inhibitory activity against *L. monocytogenes*. Agilicin C7 biosynthesis gene was characterized by bioinformatics analyses and heterologously expressed in *Escherichia coli* for further study. The anti-listeria activity of recombinant agilicin C7 (r-agilicin C7) was lost by proteases and  $\alpha$ -amylase, suggesting that agilicin C7 is a glycoprotein. r-Agilicin C7 has wide pH and thermal stability and is also stable in various organic solvents. It destroyed *L. monocytogenes* by damaging the integrity of the cell envelope. These properties of r-agilicin C7 indicate that agilicin C7 is a novel amylase-sensitive anti-listerial Class IId bacteriocin. Physicochemical stability and inhibitory activity against *L. monocytogenes* of r-agilicin C7 suggest that it can be applied to control *L. monocytogenes* in the food industry, including dairy and meat products.

**Keywords** anti-listerial, bacteriocin, *Ligilactobacillus agilis*

### Introduction

*Listeria monocytogenes* is a Gram-positive pathogenic bacterium found in humans and animals (McLauchlin et al., 2004; Yoon and Kim, 2022). It has a high infection ability owing to host-pathogen interactions (Tsai and Chen, 2020). In humans, it is mainly transmitted through food and can cause listeriosis, which is especially fatal to children, elderly people, and immune-deficient patients (Gandhi and Chikindas, 2007). Listeriosis is caused by consuming food contaminated with *L. monocytogenes*, such as non-pasteurized milk, cheese, undercooked meat, and smoked ducks (Aygün and Pehlivanlar, 2006; Jang et al., 2021; Park et al., 2021). In particular, its ability to grow at low temperatures, even refrigerated temperatures, prevents its use in human foodstuffs (Possas et al., 2022).

Many biological agents that prevent contamination by or inhibit the growth of *L. monocytogenes* have been studied, and bacteriocins are essential candidates for controlling *L. monocytogenes* (Perumal et al., 2019). Bacteriocins are ribosomally synthesized proteinaceous substances produced by bacteria that are active against other bacteria (Cotter et al., 2005; Lee et al., 2021) and have been proposed as an alternative to antibiotics to inhibit the growth of pathogenic microorganisms (Cotter et al., 2013). Bacteriocins with antilisterial activity have been isolated from diverse lactic acid bacteria (LAB) such as *Enterococcus* spp., *Lactobacillus* spp., *Leuconostoc* spp., and *Weissella* spp. (Dong et al., 2022; Ghalfi et al., 2006; Todorov, 2010; Woo et al., 2021; Yang and Moon, 2021). Among them, nisin, an antibiotic bacteriocin produced by *Lactococcus lactis*, is the only bacteriocin used in the food industry in many countries to improve food safety and extend the shelf life of foods (Churklam et al., 2020; van Kuyjk et al., 2012). However, nisin-resistant *L. monocytogenes* mutants have also been reported (Sabaté and Audisio, 2013; Vadyvaloo et al., 2004), suggesting that new biological agents should be developed to overcome nisin resistance.

*Ligilactobacillus agilis* is a motile lactic acid bacterium generally isolated from the gastrointestinal tract of mammals (Baele et al., 2001; Kajikawa et al., 2018). Several *L. agilis* have been shown to inhibit *L. monocytogenes* (Gudiña et al., 2015; Shi et al., 2021). However, more information must be given regarding the antilisterial substances from *L. agilis*. Here, we report the isolation of *L. agilis* C7, which inhibits *L. monocytogenes* that contaminates various foods including meat and dairy products, and characterize a novel bacteriocin from *L. agilis* C7, named agilicin C7, at the genetic and molecular levels by heterologous expression in *Escherichia coli*.

## Materials and Methods

### Isolation and identification of lactic acid bacteria (LAB) with anti-listerial activity

Feces from healthy pigs were used to isolate LAB with antilisterial activities. One gram of fecal sample was mixed with 30 mL of sterile phosphate buffer (pH 7.4) and diluted ten-fold. The dilutions were spread on the de Man, Rogosa, and Sharpe (MRS; BD, Franklin Lakes, NJ, USA) agar plates containing 0.02% (w/v) sodium azide (Sigma-Aldrich, St. Louis, MO, USA) and 0.02% (w/v) Bromocresol Purple (BCP; Sigma-Aldrich). The plates were incubated at 37°C overnight. Colonies were selected and streaked onto MRS agar. To screen for anti-listerial LAB, the isolates were cultured in MRS broth at 37°C for 24 h, and the cell-free supernatant (CFS) was collected by centrifugation (8,000×g, 10 min, 4°C). The CFS was adjusted to pH 7.0 and filtered using a 0.45 µm pore size syringe filter (Minisart®, Sartorius, Göttingen, Germany). The antilisterial activity of CFS was measured using an agar well diffusion assay, with *L. monocytogenes* ATCC 19114 as an indicator strain (Yoon and Kim, 2022).

The 16S rRNA gene sequencing was used to identify the isolate with antilisterial activity. The 16S rRNA gene was amplified using a PCR Thermal Cycler (Dice® Gradient, TaKaRa, Shiga, Japan) with primers 27F (5'-AGAGTTTGATCCTGGCTCAG-3') and 1492R (5'-TACGTTACCTGTACGACTT-3'). The PCR condition was 35 cycles of denaturation at 95°C for 30 sec, annealing at 55°C for 30 sec, and extension at 72°C for 1 min. A homology search was performed using the National Center for Biotechnology Information BLAST (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>).

### Kinetics of the anti-listerial substance production

The kinetics of bacterial growth and production of antimicrobial substances by *L. agilis* C7 inhibiting growth of *L. monocytogenes* were investigated during batch fermentation. A culture of *L. agilis* C7 was inoculated at a 2% volume into

200 mL MRS broth and incubated at 37°C for 72 h. Samples were collected at designated times over a 72 h period. Cell growth was determined by measuring the optical density at 600 nm using a spectrophotometer (Shimadzu, Kyoto, Japan). The production of anti-listerial substances was compared by measuring the size of the inhibition zone of the neutralized CFS against *L. monocytogenes*.

### **Mining and characterization of the gene encoding anti-listerial bacteriocin**

Bioinformatic analyses of the *L. agilis* C7 genome, previously reported by Yoo et al. (2022), were conducted for mining for the bacteriocin gene, which is responsible for the anti-listerial activity, using BAGLE4 (<http://bagel4.molgenrug.nl/>), BACTIBASE (<https://bactibase.hammamilab.org>), AntiSMASH (<https://antismash.secondarymetabolites.org>), and BLASTP (<https://blast.ncbi.nlm.nih.gov>). Signal P 5.0 (<https://services.healthtech.dtu.dk/service.php?SignalP-5.0>) and MultAlin (<http://multalin.toulouse-ra.fr/multalin/>) were used to predict signal peptides and for multiple sequence alignment with other bacteriocins, respectively.

### **Cloning of the bacteriocin candidate genes in *Escherichia coli***

Sequence analysis of the putative bacteriocin biosynthesis gene, agilicin C7, showed an Ala-Ser-Ala cleavage site at the N-terminus of the signal sequence (Biet et al., 1998; Worobo et al., 1995). Forward and reverse primers targeting the mature form of agilicin C7 were designed based on *L. agilis* C7 genome sequences (Yoo et al., 2022). PCR amplification was performed using forward (5'-AAA GAATTC CATATG AACTCGGGTTACTGTAAACC-3') and reverse (5'-AAA GTCGAC CCCCCT ATAACCAGGTTG-3') primers, and the PCR products were purified using the NucleoSpin Gel and PCR Clean-up Kit (Macherey-Nagel, Düren, Germany). After the purified PCR products and pET21b vector were digested with *NdeI* and *Sall* for 1 h at 37°C, ligation was carried out using T4 Ligase overnight at 16°C. The pET-21b expression vector containing the agilicin C7 gene, named pET-21b-*agiC7*, was transformed into *E. coli* BL21 (DE3; Hwang et al., 2018). Then, recombinant transformants were selected on the Luria-Bertani (LB, BD) agar containing ampicillin (100 µg/mL).

### **Heterologous expression and purification of recombinant agilicin C7 in *Escherichia coli***

Overnight culture of *E. coli* BL21 (DE3) harboring the plasmid pET-21b-*agiC7* inoculated 1% (v/v) to LB broth containing ampicillin (100 µg/mL) and then was grown at 37°C. After inducing protein overexpression by adding 1 mM isopropyl β-D-1-thiogalactopyranoside (IPTG; Sigma-Aldrich), the cells were recovered by centrifugation at 10,000×g for 10 min. The cells were then resuspended in TN buffer (20 mM Tris-Cl, 100 mM NaCl, pH 8.0) and disrupted using a sonicator (VCX750; Sonics, Oklahoma City, OK, USA). The insoluble cytoplasmic fraction and cell debris were collected by centrifugation (13,000×g, 10 min, 4°C) because recombinant agilicin C7 (r-agilicin C7) was expressed as an inclusion body (data not shown), and the resulting pellet was dissolved in TN buffer containing 8 M urea. The His-tagged r-agilicin C7 was purified using nickel-nitroloacetic acid (Ni-NTA) affinity chromatography and analyzed by Tricine sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and zymogram.

### **Effect of enzymes, temperature, pH, and solvents on anti-listerial activity**

The purified r-agilicin C7 was incubated for 2 h at 37°C after adding proteinase-K, α-amylase, α-chymotrypsin, pepsin, and lipase at a 5 mg/mL concentration, respectively (Yang and Moon, 2021). All the enzymes were purchased from Sigma-Aldrich. After boiling for 5 min at 80°C to inactivate enzymes, the anti-listerial activity of the samples was evaluated by the

agar well diffusion method (Yoon and Kim, 2022).

To examine the stability of the anti-listerial activity of r-agilicin C7 over a wide range of pH, the pH of r-agilicin C7-containing solutions was adjusted to pH 1.0 to 10.0 using sterile 1 N NaOH or 1 N HCl. After incubation at 37°C for 2 h, the samples were readjusted to pH 8.0 and tested using the agar well diffusion method (Hwang et al., 2018).

The effect of temperature on anti-listerial activity was tested by incubating at 4°C, 25°C, 37°C, 50°C, 60°C, 80°C, 100°C, and 121°C for 15 min. The treated samples were then cooled to room temperature, and their anti-listerial activity was evaluated using the agar well diffusion method (Hwang et al., 2018).

The effect of organic solvents on the anti-listerial activity of r-agilicin C7 was evaluated by adding 10% (final concentration, v/v) of acetone, acetonitrile, methanol, ethanol, chloroform, and isopropanol, respectively. After incubating at 37°C for 2 h, the anti-listerial activity of the samples was evaluated by the agar well diffusion method (Hwang et al., 2018).

### **Antimicrobial spectrum of r-agilicin C7**

The antimicrobial activity of r-agilicin C7 against indicator strains, including Gram-negative and Gram-positive bacteria, and 2 fungi (Table 1), was examined. After overlaying the soft agar (0.75% agar) containing indicator strains on agar media, the wells were punched into the agar plates using a sterile glass cylinder (Hwang et al., 2018). The wells were filled with 200 µL of r-agilicin C7 solution and incubated at the optimal growth temperature. The antimicrobial activity was measured based on a clear zone around the wells.

### **Tricine sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and zymogram**

Tricine SDS-PAGE was performed to confirm the molecular weight of the r-agilicin C7, as described by Hwang et al. (2018), using 4% acrylamide stacking gel, 10% acrylamide spacer gel, and 16.5% acrylamide separation gel. The samples were mixed with Tricine sample buffer (Bio-Rad Laboratories, Hercules, CA, USA). Electrophoresis was performed in an electrophoresis chamber (PowerPac 300; Bio-Rad Laboratories) at 30 V for 9 h. After running, one gel was stained with staining buffer (10% acetic acid, 0.025% Coomassie Brilliant Blue G-250 (Bio-Rad Laboratories), 45% methanol, and 45% distilled water) for 1 h and then destained with destaining buffer (10% acetic acid, 45% methanol, and 45% distilled water) for 12 h. The other gel was fixed with r-agilicin protein in a destaining buffer overnight and then washed in distilled water for 3 h. The washed gels were placed in the square dishes and overlaid with 0.75% (w/v) soft agar inoculated with *L. monocytogenes* ATCC 19114 (approximately  $10^5$  CFU/mL). The plate was incubated for 6 h at 37°C, and the gel was examined for the inhibition zone.

### **Field emission-scanning electron microscopy (FE-SEM)**

FE-SEM was used to observe morphological changes in *L. monocytogenes* ATCC 19114 treated with various concentrations of r-agilicin C7 in BHI broth. An overnight culture of *L. monocytogenes* ATCC 19114 (approx.  $10^4$  CFU/mL) was inoculated into BHI broth containing r-agilicin C7 (0, 160 AU/mL, 320 AU/mL, respectively) and incubated for 2 h at 37°C. The treated cells were collected by centrifugation at  $6,000\times g$  for 10 min, and the pellets were washed twice with 1 mM PBS buffer (pH 7.0). The cells were treated with 0.1 M sodium cacodylate buffer containing 4% glutaraldehyde (EMS, Hatfield, PA, USA) and washed with 0.05 M sodium cacodylate three times for 10 min. Post-fixation was carried out with 1% osmium tetroxide (EMS) and then washed with 0.05 M sodium cacodylate, as mentioned above. The fixed cells were resuspended in ethanol

**Table 1. Antimicrobial spectrum of recombinant agilicin C7**

Indicator organism	Inhibition zone	Media
Gram-negative		
<i>Pseudomonas aeruginosa</i> KCCM 11266	–	NB
<i>Salmonella gallinarum</i> KVCC-BA0700722	–	NB
<i>Shigella flexneri</i> KCCM 40414	–	NB
<i>Salmonella typhimurium</i> KCCM 40253	–	NB
<i>Escherichia coli</i> K88	–	LB
<i>Klebsiella pneumoniae</i> KCCM11418	–	NB
<i>Salmonella enteritidis</i> KCCM12021	–	NB
<i>E. coli</i> DH5 $\alpha$ (Cloning host)	–	LB
<i>E. coli</i> BL21 (Cloning host)	–	LB
Gram-positive		
<i>Listeria monocytogenes</i> ATCC 19114	++	BHI
<i>Listeria monocytogenes</i> KCTC 3710	++	BHI
<i>Listeria grayi</i> KCTC 3433	++	BHI
<i>Listeria innocua</i> KCTC 3586	++	BHI
<i>Listeria welshimeri</i> KCTC 3587	++	BHI
<i>Enterococcus faecium</i> KCTC 13225	+	BHI
<i>Enterococcus faecalis</i> KCTC 3206	+	BHI
<i>Staphylococcus aureus</i> KFRI 00188	–	NB
<i>Bacillus licheniformis</i> KCCM 12145	–	NB
<i>Bacillus cereus</i> KCTC1661	–	LB
<i>Lactobacillus mucosae</i> LM1	–	MRS
<i>Lactobacillus plantarum</i> SK156	–	MRS
<i>Lactobacillus johnsonii</i> PF01	–	MRS
<i>Lactococcus garvieae</i> KCTC 5618	–	MRS
<i>Ligilactobacillus agilis</i> KCTC 3158	–	MRS
Fungi		
<i>Aspergillus flavus</i> KCTC6901	–	PDA
<i>Saccharomyces cerevisiae</i> KCCM7919	–	YM

Antimicrobial activity was measured by agar well diffusion. Inhibition zone was expressed in degrees: –, no inhibition; +, up to 1 mm; ++, up to 5 mm. KCCM, Korean Culture Center of Microorganisms; NB, nutrient broth (BD, Franklin Lakes, NJ, USA); KVCC, Korea Veterinary Culture Collection; LB, Luria-Bertani (BD); ATCC, American Type Culture Collection; BHI, brain heart infusion (BD); KCTC, Korean Collection for Type Culture; KFRI, Kerala Forest Research Institution; MRS, de Man Rogosa and Sharpe (BD); PDA, potato dextrose agar (BD); YM, yeast alt (BD).

(50%, 75%, 90%, 95%, and 100% for 20 min each) and treated with 100% isoamyl acetate (Tokyo Chemical Industry, Tokyo, Japan) to dehydrate. The morphology of *L. monocytogenes* ATCC 19114 cells treated with r-agilicin C7 was observed using a SIGMA 500 FE-SEM (Carl Zeiss, Oberkochen, Germany).

## Results

### Isolation of a bacterial strain with anti-listerial activity


Bacterial strains with a yellow halo on the BCP-MRS medium were isolated from pig fecal samples. Their CFSs were screened for antimicrobial activity against *L. monocytogenes* ATCC 19114 using an agar well diffusion assay. The isolate with the strongest inhibitory activity was selected for further experiments. 16S rRNA gene sequencing showed a close phylogenetic similarity to *L. agilis* LGM7-7 (99.93%), then it was identified as an *L. agilis* species and designated *L. agilis* C7. The genome sequence of the *L. agilis* C7 strain was deposited in the GenBank database under accession no. JAMGEC000000000 (Yoo et al., 2022).

The kinetics of anti-listerial substance production in *L. agilis* C7 indicated that it was associated with cell growth, as shown in Fig. 1. The anti-listerial substance was detected in the logarithmic growth phase, and its activity was stably maintained during the stationary growth phase.


### Mining and characterization of the gene encoding anti-listerial bacteriocin

To unravel the genetic traits relevant to the anti-listerial activity of the C7 strain, bioinformatic analyses of the C7 genome were performed, which revealed a complete operon highly homologous to an operon for garvicin Q biosynthesis. It was selected for further investigation because Garvicin Q family class II bacteriocins inhibit *L. monocytogenes* (Tosukhowong et al., 2012). Fig. 2A shows the organization of the C7 gene cluster, which is highly homologous to the Garvicin Q bacteriocin. It consists of four genes encoding, ABP transporter permease, histidine kinase, bacteriocin (named “agilicin C7”), and bacteriocin immunity protein. The nucleotide and deduced amino acid sequences of agilicin C7 are shown in Fig. 2B. Amino acid sequence analysis of agilicin C7 revealed an Ala-Ser-Ala cleavage site at the N-terminus of its signal sequence, which is also found in several class II d bacteriocins (Worobo et al., 1995). The Ala-Ser-Ala site at the N-terminus of divergicin A serves as a signal peptide for the cell's general export system, such as the *E. coli* *sec* pathway (Biet et al., 1998).

The putative amino acid sequence of agilicin C7 showed high homology with those of Class II d bacteriocins (Fig. 2C). It



**Fig. 1.** Kinetic growth production of an anti-listerial substance from *Ligilactobacillus agilis* C7 (A) and anti-listerial activity of the culture supernatant of *L. agilis* C7 (B). (A) Black circle, the growth curve of *L. agilis* C7; Gray bar, production of the anti-listerial substance. (B) Anti-listerial activity was detected by the agar well diffusion method.




**Fig. 2. Genetic organization (A), nucleotide sequence (B), and multi-alignment (C) of the agilicin C7.** (A) Genetic map of the predicted agilicin C7 operon, (B) the deduced amino acid sequences of agilicin C7 gene are presented below the nucleotide sequence. The putative -35 and -10 promoter regions and putative RBS are underlined. (C) Multiple sequence alignment of known class II d bacteriocins. Similar amino acids are shaded blue, and identical amino acids are shaded red. RBS, ribosome binding site.

shared homology with bovicin 255 (69%) from *Streptococcus bovis* and garvicin Q (52%) from *Lactococcus garviae*. Agilicin C7 has the N-terminal NGY and central YxVTK motifs, a typical feature of Class II d bacteriocins.

### Heterologous expression of agilicin C7 in *Escherichia coli*

We attempted to express agilicin C7 in *E. coli* for characterization heterologously. Because sequence analysis of agilicin C7 showed an Ala-Ser-Ala cleavage site at the N-terminus of the signal sequence (Biet et al., 1998; Worobo et al., 1995), mature sequences of the agilicin C7 gene without signal sequences were amplified from the genomic DNA of *L. agilis* C7 using PCR. The PCR product of agilicin C7 was inserted into the *Nde*I-*Sal*I site of the pET-21b expression vector, and the resulting plasmid was introduced into *E. coli* BL21 (DE3) as an expression host by electroporation. Recombinant agilicin C7 was expressed as an inclusion body under IPTG induction and purified using a His-tag purification column (data not shown). Figs. 3A and B shows that r-agilicin C7 exhibits anti-listerial activity. In addition, tricine-SDS-PAGE and zymography (Fig. 3A) showed that approximately 10 kDa of r-agilicin C7 was expressed, confirming its inhibitory activity against *L. monocytogenes*.



**Fig. 3. Tricine SDS-PAGE and zymogram of recombinant agilicin C7 (A) and agar well diffusion assay (B).** (A) Lane 1, prestained protein standards marker; Lane 2 and 3, SDS-PAGE of recombinant agilicin C7; Lane 4 and 5, zymogram of recombinant agilicin C7. (B) Agar well diffusion assay shows that recombinant agilicin C7 has inhibitory activity against *Listeria monocytogenes* ATCC 19114. SDS-PAGE, sodium dodecyl sulfate polyacrylamide gel electrophoresis.

### Antimicrobial activity spectrum

Since r-agilicin C7 inhibited the growth of *L. monocytogenes*, it was tested whether it showed antimicrobial activity against other bacteria and fungi. As shown in Table 1, r-agilicin C7 inhibited the growth of other listerial species, including *Listeria grayi*, *Listeria innocua*, *Listeria welshimeri*, and *Enterococcus* spp. However, it did not show antimicrobial activity against the other gram-positive and gram-negative bacteria tested or fungi.

### Effect of enzymes, temperature, pH, and solvents on anti-listerial activity

The stability of the r-agilicin C7 against enzymes, heat, pH, and organic solvents was also tested. As shown in Table 2, the anti-listerial activity of r-agilicin C7 was lost by pepsin,  $\alpha$ -chymotrypsin, protein K, and trypsin, indicating that agilicin C7 is a proteinaceous substance. Interestingly, its activity lost by  $\alpha$ -amylase also lost its activity. These results suggested that agilicin C7 is a glycoprotein. However, SDS-PAGE analysis did not show decrease in the molecular weight of r-agilicin C7 after  $\alpha$ -amylase treatment (data not shown), perhaps due to small change in molecular weight. It is necessary to detect the molecular weight change of r-agilicin C7 by  $\alpha$ -amylase treatment using liquid chromatography-mass spectrometer in the future, to support that agilicin C7 is glycoprotein.

The anti-listerial activity of r-agilicin C7 was maintained, even after treatment at 100°C for 15 min. In addition, the inhibitory activity was stable in the pH range of 1–10. Furthermore, the anti-listerial activity of r-agilicin C7 was stable after treatment with several organic solvents. These results indicate that agilicin C7 has wide pH and thermal stability, which are features of Class II bacteriocins (Negash and Tsehai, 2020).

### Action mode of r-agilicin C7 bacteriocin

To understand the action mode of r-agilicin C7, FE-SEM analysis was performed to observe cell morphological changes of

**Table 2.** Effect of enzyme, pH, temperature, and solvent treatment on the activity of the recombinant agilicin C7

Treatment	Residual activity (%) <sup>1)</sup>
Temperature (°C)	
4	100
25	100
37	100
50	100
60	100
80	90
100	80
121	ND
pH	
1	100
2	100
3	100
4	100
5	100
6	100
7	100
8	100
9	95
10	95
Enzyme (5 mg/mL)	
$\alpha$ -Amylase	ND
$\alpha$ -Chymotrypsin	ND
Proteinase-K	ND
Trypsin	ND
Pepsin	ND
Lipase	90
Solvent	
Acetone	90
Acetonitrile	90
Methanol	90
Ethanol	95
Chloroform	100
Isopropanol	100

<sup>1)</sup> Residual activity (%) is measured by the diameter of the inhibition zones compared to untreated control. ND, not detected.

*L. monocytogenes* as an indicator strain. Intact rod-shaped cell structures were observed before r-agilicin C7 treatment (Fig. 4A). The cells appeared extensively wrinkled and porous after treatment with r-agilicin C7 for 2 h, and some cells were lysed entirely, extruding the intracellular material (Figs. 4B and C). Additionally, the degree of cell lysis was dose dependent. Based on these results, r-agilicin C7 destroyed *L. monocytogenes* by damaging the integrity of the cell envelope, which is a typical property of class II bacteriocins.


## Discussion

In this study, we isolated a novel class IId bacteriocin, agilicin C7, from *L. agilis*. Several *L. agilis* strains have been reported to exhibit inhibitory activities against foodborne pathogens (Gudiña et al., 2015; Shi et al., 2021). However, information on antimicrobial substances on the biochemical and molecular basis needs to be included. This is the first study to characterize the genetic and molecular properties of an anti-listerial bacteriocin isolated from *L. agilis*.

Comparative genomic analysis of *L. agilis* C7 revealed a complete operon for putative bacteriocin biosynthesis, which is highly homologous to the operon for garvicin Q biosynthesis. The agilicin C7 biosynthesis gene contains the N-terminal NGY and central YxVTK motifs, a typical feature of class IId bacteriocins (Tymoszevska et al., 2017; Tymoszevska et al., 2020). Our data show that r-agilicin C7 remained active over a broad pH range and was thermally stable. It was also stable under several organic solvent treatments, all typical properties of class II bacteriocins (Negash and Tsehai, 2020).

Class II bacteriocins have been reported to have bactericidal and pore-forming modes of action (Simons et al., 2020). Class IId is a heterogeneous group of linear, unmodified, non-pediocin-like peptides (Iwatani et al., 2011). Tymoszevska et al. (2017) reported that garvicin Q, a class IId bacteriocin, targets Man-PTS in target cells. The nonpediocin-like class IId bacteriocin, lactococcin A, destroyed target cells through the pore formation mode of action (van Belkum et al., 1991). In this study, *L. monocytogenes* cells were extensively wrinkled and porous in response to r-agilicin C7. Some cells were lysed entirely, extruding intracellular material, suggesting that agilicin C7 has the same or a similar mode of action. To date, multiple Man-PTS systems have been discovered in *L. monocytogenes* (Aké et al., 2011), and they control not only sugar transport activity but also the expression of virulence genes (Gaballa et al., 2019). The host specificity of bacteriocins is determined by their outer membrane receptors and inner membrane proteins, which act as scaffolds for bacteriocin insertion and pore formation (Etayash et al., 2016).

Interestingly, the antimicrobial activity of r-agilicin C7 was abrogated by amylase. To date, several amylase-sensitive



**Fig. 4.** Field emission scanning electron microscopy images of *Listeria monocytogenes* ATCC 19114. (A) Untreated, (B) after treatment with recombinant agilicin C7 (160 AU/mL), (C) after treatment with recombinant agilicin C7 (320 AU/mL). Red arrow indicates the pore-forming.

bacteriocins have been reported (Kang and Kim, 2010; Keppler et al., 1994; Seo et al., 2014), most of which are classified as class IV (Heredia-Castro et al., 2015; Lee and Kim, 2010; Todorov, 2010) or unclassified (Hernández et al., 2005; Seo et al., 2014). Glycosylated bacteriocin is termed “glycocin” (Stepper et al., 2011), which is a cysteine-containing glycoactive peptide in which the O- or S-linked glycan appears essential (Balciunas et al., 2013). However, agilicin C7 does not have these properties, nor does it show sequence homology with glycocin. Therefore, agilicin C7 can be recognized as a novel subgroup of class IId bacteriocins, although the classification of bacteriocins is still controversial. It is necessary to compare the molecular weight change of r-agilicin C7 before and after  $\alpha$ -amylase treatment to support that agilicin C7 is glycoprotein, and which amino acid of agilicin C7 is glycosylated in the future.

Many bacteriocins from LAB have been characterized, and these studies have indicated potential applications of bacteriocins in food preservation and safety. Agilicin C7 has physicochemical stability and solid inhibitory activity against *L. monocytogenes*, suggesting that it can be applied to control *L. monocytogenes* in the food industry, including dairy and meat products (Yang and Moon, 2021). In the present study, successful cloning and expression of agilicin C7 in *E. coli* may facilitate mass production of this bacteriocin in high concentration and permit its use as safe antimicrobial food additive. However, whether agilicin C7 can inhibit *L. monocytogenes* in food matrices need to be investigated in the future because the efficacy and stability of a bacteriocin in foodstuffs are relevant to the chemical and physical properties of foods (Gálvez et al., 2007).

## Conflicts of Interest

The authors declare no potential conflicts of interest.

## Acknowledgements

This work was supported by the Korea Institute of Planning and Evaluation for Technology in Food, Agriculture, and Forestry (IPET) through the High Value-added Food Technology Development Program, funded by the Ministry of Agriculture, Food and Rural Affairs (MAFRA) (grant no. 321035052HD040).

## Author Contributions

Conceptualization: Kang DK. Data curation: Yoo JM, Song JH, Hwang IC. Formal analysis: Yoo JM, Song JH, Vasquez R, Lee JS. Methodology: Yoo JM, Vasquez R, Hwang IC, Lee JS. Software: Yoo JM. Validation: Lee JS, Kang DK. Investigation: Vasquez R, Kang DK. Writing - original draft: Yoo JM, Kang DK. Writing - review & editing: Yoo JM, Song JH, Vasquez R, Hwang IC, Lee JS, Kang DK.

## Ethics Approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

## References

Aké FMD, Joyet P, Deutscher J, Milohanic E. 2011. Mutational analysis of glucose transport regulation and glucose-mediated

- virulence gene repression in *Listeria monocytogenes*. Mol Microbiol 81:274-293.
- Aygun O, Pehlivanlar S. 2006. *Listeria* spp. in the raw milk and dairy products in Antakya, Turkey. Food Control 17:676-679.
- Baele M, Devriese LA, Haesebrouck F. 2001. *Lactobacillus agilis* is an important component of the pigeon crop flora. J Appl Microbiol 91:488-491.
- Balciunas EM, Martinez FAC, Todorov SD, de Melo Franco BDG, Converti A, de Souza Oliveira RP. 2013. Novel biotechnological applications of bacteriocins: A review. Food Control 32:134-142.
- Biet F, Berjeaud JM, Worobo RW, Cenatiempo Y, Fremaux C. 1998. Heterologous expression of the bacteriocin mesentericin Y105 using the dedicated transport system and the general secretion pathway. Microbiology 144:2845-2854.
- Churklam W, Chaturongakul S, Ngamwongsatit B, Aunpad R. 2020. The mechanisms of action of carvacrol and its synergism with nisin against *Listeria monocytogenes* on sliced bologna sausage. Food Control 108:106864.
- Cotter PD, Hill C, Paul Ross R. 2005. Bacteriocins: Developing innate immunity for food. Nat Rev Microbiol 3:777-788.
- Cotter PD, Paul Ross R, Hill C. 2013. Bacteriocins: A viable alternative to antibiotics? Nat Rev Microbiol 11:95-105.
- Dong B, Lin Y, Wang J, Du W, Sun C, Fu S, Wu T. 2022. Antibacterial activity of antimicrobial peptide gcDefb1 against foodborne pathogenic bacteria and its application in pork storage. Food Sci Biotechnol 31:597-605.
- Etayash H, Azmi S, Dangeti R, Kaur K. 2016. Peptide bacteriocins: Structure activity relationships. Curr Top Med Chem 16:220-241.
- Gaballa A, Guariglia-Oropeza V, Wiedmann M, Boor KJ. 2019. Cross talk between SigB and PrfA in *Listeria monocytogenes* facilitates transitions between extra- and intracellular environments. Microbiol Mol Biol Rev 83:e00034-19.
- Gálvez A, Abriouel H, López RL, Omar NB. 2007. Bacteriocin-based strategies for food biopreservation. Int J Food Microbiol 120:51-70.
- Gandhi M, Chikindas ML. 2007. *Listeria*: A foodborne pathogen that knows how to survive. Int J Food Microbiol 113:1-15.
- Ghalfi H, Allaoui A, Destain J, Benkerroum N, Thonart P. 2006. Bacteriocin activity by *Lactobacillus curvatus* CWBI-B28 to inactivate *Listeria monocytogenes* in cold-smoked salmon during 4°C storage. J Food Prot 69:1066-1071.
- Gudiña EJ, Fernandes EC, Teixeira JA, Rodrigues LR. 2015. Antimicrobial and anti-adhesive activities of cell-bound biosurfactant from *Lactobacillus agilis* CCUG31450. RSC Adv 5:90960-90968.
- Heredia-Castro PY, Méndez-Romero JI, Hernández-Mendoza A, Acedo-Félix E, González-Córdova AF, Vallejo-Cordoba B. 2015. Antimicrobial activity and partial characterization of bacteriocin-like inhibitory substances produced by *Lactobacillus* spp. isolated from artisanal Mexican cheese. J Dairy Sci 98:8285-8293.
- Hernández D, Cardell E, Zárate V. 2005. Antimicrobial activity of lactic acid bacteria isolated from Tenerife cheese: Initial characterization of plantaricin TF711, a bacteriocin-like substance produced by *Lactobacillus plantarum* TF711. J Appl Microbiol 99:77-84.
- Hwang IC, Oh JK, Kim SH, Oh S, Kang DK. 2018. Isolation and characterization of an anti-listerial bacteriocin from *Leuconostoc lactis* SD501. Korean J Food Sci Anim Resour 38:1008-1018.
- Iwatani S, Zendo T, Sonomoto K. 2011. Class IId or linear and non-pediocin-like bacteriocins. In Prokaryotic antimicrobial peptides: From genes to applications. Drider D, Rebuffat S (ed). Springer, New York, NY, USA. pp 237-252.
- Jang YS, Moon JS, Kang HJ, Bae D, Seo KH. 2021. Prevalence, characterization, and antimicrobial susceptibility of *Listeria monocytogenes* from raw beef and slaughterhouse environments in Korea. Foodborne Pathog Dis 18:419-425.
- Kajikawa A, Suzuki S, Igimi S. 2018. The impact of motility on the localization of *Lactobacillus agilis* in the murine gastrointestinal tract. BMC Microbiol 18:68.

- Kang TK, Kim WJ. 2010. Characterization of an amylase-sensitive bacteriocin DF01 produced by *Lactobacillus brevis* DF01 isolated from dongchimi, Korean fermented vegetable. *Korean J Food Sci Anim Resour* 30:795-803.
- Keppler K, Geisen R, Holzapfel WH. 1994. An  $\alpha$ -amylase sensitive bacteriocin of *Leuconostoc carnosum*. *Food Microbiol* 11:39-45.
- Lee HJ, Kim WJ. 2010. Isolation and characterization of anti-listerial and amylase sensitive enterocin producing *Enterococcus faecium* DB1 from *Gajami-sikhae*, a fermented flat fish in Korea. *Food Sci Biotechnol* 19:373-381.
- Lee JS, Park SW, Lee HB, Kang SS. 2021. Bacteriocin-like inhibitory substance (BLIS) activity of *Enterococcus faecium* DB1 against biofilm formation by *Clostridium perfringens*. *Probiotics Antimicrob Proteins* 13:1452-1457.
- McLauchlin J, Mitchell RT, Smerdon WJ, Jewell K. 2004. *Listeria monocytogenes* and listeriosis: A review of hazard characterisation for use in microbiological risk assessment of foods. *Int J Food Microbiol* 92:15-33.
- Negash AW, Tsehai BA. 2020. Current applications of bacteriocin. *Int J Microbiol* 2020:4374891.
- Park E, Ha J, Oh H, Kim S, Choi Y, Lee Y, Kim Y, Seo Y, Kang J, Yoon Y. 2021. High prevalence of *Listeria monocytogenes* in smoked duck: Antibiotic and heat resistance, virulence, and genetics of the isolates. *Food Sci Anim Resour* 41:324-334.
- Perumal V, Yao Z, Kim JA, Kim HJ, Kim JH. 2019. Purification and characterization of a bacteriocin, BacBS2, produced by *Bacillus velezensis* BS2 isolated from meongge jeotgal. *J Microbiol Biotechnol* 29:1033-1042.
- Possas A, Hernández M, Esteban-Carbonero O, Valero A, Rodríguez-Lázaro D. 2022. *Listeria monocytogenes* survives better at lower storage temperatures in regular and low-salt soft and cured cheeses. *Food Microbiol* 104:103979.
- Sabaté DC, Audisio MC. 2013. Inhibitory activity of surfactin, produced by different *Bacillus subtilis* subsp. *subtilis* strains, against *Listeria monocytogenes* sensitive and bacteriocin-resistant strains. *Microbiol Res* 168:125-129.
- Seo SH, Jung M, Kim WJ. 2014. Antilisterial and amylase-sensitive bacteriocin producing *Enterococcus faecium* SH01 from Mukeunji, a Korean over-ripened kimchi. *Food Sci Biotechnol* 23:1177-1184.
- Simons A, Alhanout K, Duval RE. 2020. Bacteriocins, antimicrobial peptides from bacterial origin: Overview of their biology and their impact against multidrug-resistant bacteria. *Microorganisms* 8:639.
- Shi S, Cheng B, Gu B, Sheng T, Tu J, Shao Y, Qi K, Zhou D. 2021. Evaluation of the probiotic and functional potential of *Lactobacillus agilis* 32 isolated from pig manure. *Lett Appl Microbiol* 73:9-19.
- Stepper J, Shastri S, Loo TS, Preston JC, Novak P, Man P, Moore CH, Havlíček V, Patchett ML, Norris GE. 2011. Cysteine S-glycosylation, a new post-translational modification found in glycopeptide bacteriocins. *FEBS Lett* 585:645-650.
- Todorov SD. 2010. Diversity of bacteriocinogenic lactic acid bacteria isolated from boza, a cereal-based fermented beverage from Bulgaria. *Food Control* 21:1011-1021.
- Tosukhowong A, Zendo T, Visessanguan W, Roytrakul S, Pumpuang L, Jaresitthikunchai J, Sonomoto K. 2012. Garvieacin Q, a novel class II bacteriocin from *Lactococcus garvieae* BCC 43578. *Appl Environ Microbiol* 78:1619-1623.
- Tsai YH, Chen WL. 2020. Host lipid rafts as the gates for *Listeria monocytogenes* infection: A mini-review. *Front Immunol* 11:1666.
- Tymoszevska A, Diep DB, Wirtek P, Aleksandrak-Piekarczyk T. 2017. The non-lantibiotic bacteriocin garvicin Q targets man-PTS in a broad spectrum of sensitive bacterial genera. *Sci Rep* 7:8359.
- Tymoszevska A, Walczak P, Aleksandrak-Piekarczyk T. 2020. BacSJ—Another bacteriocin with distinct spectrum of activity that targets man-PTS. *Int J Mol Sci* 21:7860.
- Vadyvaloo V, Snoep JL, Hastings JW, Rautenbach M. 2004. Physiological implications of class IIa bacteriocin resistance in *Listeria monocytogenes* strains. *Microbiology* 150:335-340.

- van Belkum MJ, Kok J, Venema G, Holo H, Nes IF, Konings WN, Abee T. 1991. The bacteriocin lactococcin A specifically increases permeability of lactococcal cytoplasmic membranes in a voltage-independent, protein-mediated manner. *J Bacteriol* 173:7934-7941.
- van Kuijk S, Noll KS, Chikindas ML. 2012. The species-specific mode of action of the antimicrobial peptide subtilisin against *Listeria monocytogenes* Scott A. *Lett Appl Microbiol* 54:52-58.
- Woo C, Jung S, Fugaban JII, Bucheli JEV, Holzapfel WH, Todorov SD. 2021. Bacteriocin production by *Leuconostoc citreum* ST110LD isolated from organic farm soil, a promising biopreservative. *J Appl Microbiol* 131:1226-1239.
- Worobo RW, Van Belkum MJ, Sailer M, Roy KL, Vederas JC, Stiles ME. 1995. A signal peptide secretion-dependent bacteriocin from *Carnobacterium divergens*. *J Bacteriol* 177:3143-3149.
- Yang JM, Moon GS. 2021. Partial characterization of an anti-listerial bacteriocin from *Enterococcus faecium* CJNU 2524. *Food Sci Anim Resour* 41:164-171.
- Yoo JM, Mendoza RM, Hwang IC, Kang DK. 2022. Whole genome sequence analysis of *Ligilactobacillus agilis* C7 isolated from pig feces revealed three bacteriocin gene clusters. *J Anim Sci Technol* 64:1008-1011.
- Yoon SH, Kim GB. 2022. Inhibition of *Listeria monocytogenes* in fresh cheese using a bacteriocin-producing *Lactococcus lactis* CAU2013 strain. *Food Sci Anim Resour* 42:1009-1019.